

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



United States
Department of
Agriculture

Human Nutrition
Information
Service

Home Economics
Research Report
Number 44

Foods Commonly Eaten by Individuals:

Amount Per Day and Per Eating Occasion

Eleanor M. Pao
Kathryn H. Fleming
Patricia M. Guenther
Sharon J. Mickle

FOODS COMMONLY EATEN BY INDIVIDUALS: AMOUNT PER DAY AND PER
EATING OCCASION, Eleanor M. Pao, Kathryn H. Fleming,
Patricia M. Guenther and Sharon J. Mickle
Consumer Nutrition Center, Human Nutrition Information Service,
U.S. Department of Agriculture,
Hyattsville, Maryland 20782.
Home Economics Research Report Number 44

ABSTRACT

Data on 3-day intakes of 200 commonly used foods and food groups by individuals in the Nationwide Food Consumption Survey 1977-78 are reported. The survey was conducted by the U.S. Department of Agriculture from April 1977 to March 1978 and included about 38,000 individuals in the 48 conterminous States. Data are tabulated by 16 sex-age categories and for total individuals. The 200 tables include the percentage of individuals using the food, the number of eating occasions at which the food was consumed, and the quantities (grams) of the food consumed by users per day and per eating occasion. Foods included in each table are explained briefly in the text. The number of eating occasions and quantities of food consumed by users are described by averages, standard deviations, and maximum values. Quantities consumed per day and per eating occasion are further described by values at seven specified percentiles.

Keywords: Dietary survey, eating occasion, food intake.

USDA, National Agricultural Library
NAL Bldg
10301 Baltimore Blvd
Beltsville, MD 20705-2351

Trade names are used in this publication solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the products by the U.S. Department of Agriculture or an endorsement of the Department over other products not mentioned.

Although data collection in the NFCS 77-78 was completed in mid-1978, this survey is the only source of national information on food intakes of individuals, by item and quantity, since 1965 when the first USDA nationwide survey of individuals was conducted. The results are still valid and will be used in coming years in the evaluation of many issues relating to food safety and product development, consumer demand, and nutrition education.

Supersedes Portion Sizes and Day's Intakes of Selected Foods, ARS-NE-67, issued December 1975.

CONTENTS

	Page
LIST OF TABLES.....	iii
INTRODUCTION.....	1
METHODS AND PROCEDURES.....	1
Data collection.....	1
Sample.....	2
Data handling.....	2
Selection of foods for tables.....	3
Content of tables.....	3
DESCRIPTION OF FOODS.....	4
Beverages.....	4
Eggs.....	5
Fats and oils.....	5
Fruits and fruit juices.....	6
Citrus fruit.....	6
Noncitrus fruit.....	6
Grain products.....	8
Yeast breads.....	8
Quick breads and tortillas.....	9
Cakes, cookies, pies, doughnuts, and sweet rolls..	9
Crackers and snacks.....	10
Pasta, pasta mixtures, and pizza.....	10
Rice and cooked cereals.....	11
Ready-to-eat cereals.....	11
Legumes and nuts.....	12
Meat, poultry, and fish.....	12
Meat, poultry, and fish.....	12
Processed meat.....	14
Mixtures.....	15
Milk and milk products.....	16
Fluid milk and milk beverages.....	16
Cheese.....	16
Milk products.....	17
Sugar and sweets.....	17
Vegetables.....	18
White potatoes.....	18
Raw vegetables.....	19
Cooked vegetables.....	19
Condiments.....	20
LITERATURE CITED.....	21
TABLES.....	22
APPENDIXES	
A -- System for classifying foods based on the first three digits of seven-digit food codes.....	423
B -- Table notes.....	431

LIST OF TABLES

	Table No.	Page
<u>Beverages</u>		
Coffee.....	1.01	22
Decaffeinated coffee.....	1.02	24
Tea.....	1.03	26
Carbonated soft drinks.....	1.04	28
✓ Cola-type carbonated soft drinks.....	1.05	30
Artificially sweetened carbonated soft drinks.....	1.06	32
Noncarbonated soft drinks.....	1.07	34
Fruit ades and drinks.....	1.08	36
Beer and ale.....	1.09	38
Wine.....	1.10	40
Distilled liquors.....	1.11	42
<u>Eggs</u>		
✓ Eggs.....	2.01	44
Boiled and poached eggs.....	2.02	46
Fried eggs.....	2.03	48
Scrambled eggs.....	2.04	50
<u>Fats and oils</u>		
✓ Butter.....	3.01	52
✓ Margarine.....	3.02	54
Whipped butter and margarine.....	3.03	56
✓ Salad dressings and mayonnaise.....	3.04	58
Mayonnaise.....	3.05	60
Salad dressings.....	3.06	62
Mayonnaise-type salad dressing.....	3.07	64
French dressing.....	3.08	66
Italian dressing.....	3.09	68
<u>Fruits</u>		
Citrus fruit:		
✓ Fresh grapefruit.....	4.01	70
✓ Grapefruit juice.....	4.02	72
✓ Fresh oranges.....	4.03	74
✓ Orange juice.....	4.04	76
Noncitrus fruit:		
Raw apples.....	4.05	78
Applesauce and cooked apples.....	4.06	80
Applejuice.....	4.07	82
Bananas.....	4.08	84
Cantaloup.....	4.09	86
Fruit cocktail.....	4.10	88

	Table No.	Page
Grapejuice.....	4.11	90
Raw peaches.....	4.12	92
✓ Cooked, canned, and frozen peaches.....	4.13	94
Raw pears.....	4.14	96
Cooked and canned pears.....	4.15	98
Cooked and canned pineapple.....	4.16	100
Raisins.....	4.17	102
Raw strawberries.....	4.18	104
 <u>Grain products</u>		
Yeast breads:		
Yeast breads.....	5.01	106
✓ White breads.....	5.02	108
White rolls.....	5.03	110
French, vienna, and italian breads.....	5.04	112
✱ Whole and cracked wheat breads.....	5.05	114
✓ Rye and pumpernickel breads.....	5.06	116
High fiber, wheat germ, bran, oatmeal, and multigrain breads.....	5.07	118
English muffins.....	5.08	120
Quick breads and tortillas:		
Biscuits other than refrigerator type.....	5.09	122
Refrigerator-type biscuits.....	5.10	124
✓ Cornbread.....	5.11	126
✱ Pancakes.....	5.12	128
Waffles.....	5.13	130
Tortillas.....	5.14	132
Cakes, cookies, pies, doughnuts, and sweet rolls:		
Cakes and cupcakes.....	5.15	134
Chocolate cakes and cupcakes.....	5.16	136
✱ Cookies.....	5.17	138
Chocolate chip cookies.....	5.18	140
Oatmeal cookies.....	5.19	142
Peanut butter cookies.....	5.20	144
Sugar cookies.....	5.21	146
Pies.....	5.22	148
Fruit pies.....	5.23	150
Cream and custard pies.....	5.24	152
Doughnuts.....	5.25	154
Sweet rolls.....	5.26	156
Crackers and snacks:		
✱ Crackers.....	5.27	158
Graham crackers.....	5.28	160
Saltines and soda crackers.....	5.29	162
Butter crackers.....	5.30	164
Popcorn.....	5.31	166

	Table No.	Page
Pretzels.....	5.32	168
Potato chips.....	5.33	170
Corn-based salty snacks.....	5.34	172
Pasta, pasta mixtures, and pizza:		
✓ Pasta.....	5.35	174
Spaghetti.....	5.36	176
Noodles.....	5.37	178
Macaroni.....	5.38	180
Spaghetti with tomato sauce.....	5.39	182
Macaroni and cheese.....	5.40	184
Lasagne.....	5.41	186
Pizza.....	5.42	188
Rice and cooked cereals:		
✓ Rice.....	5.43	190
Cooked cereals.....	5.44	192
Grits.....	5.45	194
Oatmeal.....	5.46	196
Wheat cereals.....	5.47	198
Ready-to-eat cereals:		
✓ Ready-to-eat cereals.....	5.48	200
Ready-to-eat cereals, 15-21 grams per cup, all sugar levels.....	5.49	202
Ready-to-eat cereals, 25-32 grams per cup, 30 percent or more sugar.....	5.50	204
Ready-to-eat cereals, 25-32 grams per cup, less than 15 percent sugar....	5.51	206
Ready-to-eat cereals, 35-45 grams per cup, 30 percent or more sugar.....	5.52	208
Ready-to-eat cereals, 35-45 grams per cup, less than 20 percent sugar....	5.53	210
Ready-to-eat cereals, 50-75 grams per cup, all sugar levels.....	5.54	212
Ready-to-eat cereals, 80 grams or more per cup, all sugar levels.....	5.55	214
 <u>Legumes and nuts</u>		
✓ Cooked dried beans.....	6.01	216
Baked beans.....	6.02	218
✓ Nuts.....	6.03	220
Peanuts.....	6.04	222
✓ Peanut butter.....	6.05	224
 <u>Meat, poultry, and fish</u>		
Meat, poultry, and fish:		
Meat, poultry, fish, and their mixtures...	7.01	226
Meat, poultry, and fish.....	7.02	228

	Table No.	Page
Beef, pork, lamb, and veal.....	7.03	230
Beef.....	7.04	232
Beef steaks.....	7.05	234
Beef roasts.....	7.06	236
Ground beef.....	7.07	238
Pork.....	7.08	240
Pork chops and steaks.....	7.09	242
Ham.....	7.10	244
Pork roasts.....	7.11	246
Bacon.....	7.12	248
Lamb.....	7.13	250
Veal.....	7.14	252
Poultry.....	7.15	254
Chicken.....	7.16	256
Turkey.....	7.17	258
Liver.....	7.18	260
Fish and shellfish.....	7.19	262
Finfish other than canned, dried, and raw.....	7.20	264
Canned tuna.....	7.21	266
Shrimp.....	7.22	268
Processed meat:		
Frankfurters.....	7.23	270
Bologna.....	7.24	272
Salami.....	7.25	274
Pork breakfast sausage.....	7.26	276
Sausage other than bologna, frank- furters, salami, and pork breakfast sausage.....	7.27	278
Mixtures:		
Beef stew with vegetables.....	7.28	280
Chili con carne.....	7.29	282
Meat loaf.....	7.30	284
Tuna salad.....	7.31	286
Pot pies.....	7.32	288
Frozen plate meals.....	7.33	290
Hamburger and cheeseburger sandwiches.....	7.34	292
Soups.....	7.35	294

Milk and milk products

Fluid milk and milk beverages:

Fluid milk, milk beverages, and milk- based infant formulas.....	8.01	296
Fluid milk.....	8.02	298

	Table No.	Page
Fluid whole milk.....	8.03	300
Fluid lowfat milk.....	8.04	302
Fluid skim milk.....	8.05	304
Reconstituted dry milk.....	8.06	306
Chocolate milk.....	8.07	308
Cocoa and hot chocolate.....	8.08	310
Cheese:		
✓ Natural and processed cheese.....	8.09	312
Cheddar cheese.....	8.10	314
Swiss cheese.....	8.11	316
Natural cheese other than cheddar, swiss, and parmesan.....	8.12	318
Processed cheese, cheese food, and cheese spread.....	8.13	320
Cottage cheese.....	8.14	322
Milk products:		
✓ Fluid cream.....	8.15	324
Liquid and frozen cream substitutes.....	8.16	326
Powdered cream substitutes.....	8.17	328
✓ Ice cream and ice milk.....	8.18	330
Nondairy whipped topping.....	8.19	332
Sour cream.....	8.20	334
Yogurt.....	8.21	336
<u>Sugar and sweets</u>		
White sugar.....	9.01	338
Sirup.....	9.02	340
Honey.....	9.03	342
✓ Jellies.....	9.04	344
Jams, preserves, and marmalades.....	9.05	346
Plain gelatin desserts.....	9.06	348
Gelatin desserts with fruit.....	9.07	350
Candy.....	9.08	352
<u>Vegetables</u>		
White potatoes:		
✓ White potatoes.....	10.01	354
Baked potatoes.....	10.02	356
Boiled potatoes.....	10.03	358
Creamed, scalloped, and au gratin potatoes.....	10.04	360
Mashed potatoes.....	10.05	362
Home-fried and hash-browned potatoes.....	10.06	364

	Table No.	Page
French fried potatoes.....	10.07	366
Potato salad.....	10.08	368
Raw vegetables:		
Cabbage salad and coleslaw.....	10.09	370
Raw carrots.....	10.10	372
Raw celery.....	10.11	374
Raw cucumbers.....	10.12	376
✓ Lettuce and tossed salads.....	10.13	378
Lettuce.....	10.14	380
Tossed salads.....	10.15	382
Raw mature onions.....	10.16	384
✓ Raw tomatoes.....	10.17	386
Cooked vegetables:		
✓ Cooked broccoli.....	10.18	388
Cooked cabbage.....	10.19	390
✓ Cooked carrots.....	10.20	392
✓ Cooked whole kernel corn.....	10.21	394
Cream-style corn.....	10.22	396
Cooked lima beans.....	10.23	398
Cooked mixed vegetables.....	10.24	400
Cooked cowpeas, field peas, and blackeye peas.....	10.25	402
✓ Cooked green peas.....	10.26	404
Cooked spinach.....	10.27	406
✓ Cooked string beans.....	10.28	408
Cooked summer squash.....	10.29	410
Cooked sweetpotatoes.....	10.30	412
Tomato juices.....	10.31	414
Condiments:		
Catsup.....	10.32	416
Mustard.....	10.33	418
Cucumber pickles.....	10.34	420

Foods Commonly Eaten by Individuals:

**Amount Per Day and
Per Eating Occasion**

Eleanor M. Pao
Kathryn H. Fleming
Patricia M. Guenther
Sharon J. Mickle¹

INTRODUCTION

This publication reports average intakes per day and per eating occasion for foods most commonly eaten by individuals who participated for 3 days in the Nationwide Food Consumption Survey (NFCS) conducted by the U.S. Department of Agriculture (USDA) in the 48 conterminous States April 1977 through March 1978. It updates and extends information for 1965 (10)² that has been widely used by business firms, trade associations, private researchers, journalists, Government agencies, and private individuals concerned with variations in the amounts of different kinds of foods consumed by individuals classified by sex and age. The data are also organized to meet the needs of nutrition specialists, consumer educators, and dietitians. The list of tables shows the commodity detail in which data are supplied to meet the requirements of market research and of educational and scientific applications.

This report provides information on many more foods than did the earlier publication and is derived from intakes of almost 38,000 individuals in 16 sex-age groups. Tables for 200 foods show the number and proportion of individuals who reported using each food, the number of occasions during the 3-day period on which they ate the food, and the quantities eaten per occasion and per day. Averages for actual users of foods are often quite different from the customary averages for whole population groups. An eating occasion was defined by the time at which one or more foods were consumed. Foods eaten at different times on the diet record were treated as separate eating occasions. The time of the eating occasion referred to the time eating began, and a few eating occasions, such as evening snacking, probably extended over a considerable time.

METHODS AND PROCEDURES

Data Collection

Information in the NFCS 1977-78 was collected during home visits by trained interviewers using a 1-day interviewer-recorded recall and a 2-day self-administered record. After the household respondent reported on food used in the household during the past week, the interviewer asked members of the household present to recall their intakes of food and beverage during the preceding day. Household members then were asked to record their intakes for the day of

¹The authors acknowledge the assistance of Brucy C. Gray, Jan L. Janiczek, and Michael S. Hamada, who developed the programs for data processing and table preparation.

²Underlined numbers in parentheses refer to Literature Cited, p. 21.

the interview and the next day, thus providing data for 3 consecutive days. Interviewers used a four-page leaflet of instructions to assist participants in describing foods and amounts consumed and left the leaflet for reference. A set of measuring cups and spoons and a ruler were given to each household for use in estimating quantities eaten. The household respondent usually provided information for children under 12 years and others unable to answer for themselves. If a household member was absent at the time of the interview but was expected to return within 2 days, forms were left to be completed. The interviewers returned to pick up the records and review them for clarity and completeness.

Sample

A stratified area probability sample of households was drawn in the 48 conterminous States in each of the four quarters from April 1977 through March 1978. About 15,000 households participated. All individuals in the households were asked to give food intake information in spring 1977. In the other three seasons all individuals under the age of 19 and one-half of those 19 years and over were asked to participate. In the three seasons other than spring, proportional representation was maintained among sex-age categories by treating each record for an individual 19 years and over as two records. In addition, a household weighting factor was applied to each individual to adjust for households included in the sample but providing no response. The unweighted number of individuals was 30,770. More than 90 percent of the individuals responding provided information for 3 days.

Data Handling

Interviewers mailed completed food intake forms to a central location for editing and coding. Coders assigned numerical codes to food items using a manual prepared at USDA (5). A seven-digit code was used to help classify foods into groups for tabulation. The first digit categorizes food items and beverages into 9 major food groups, the second digit identifies 47 major subgroups, and the third digit, 221 minor subgroups (see appendix A). The fourth digit further specifies minor subgroups. The remaining digits code information unique to the individual food. Foods with complete descriptions were assigned codes to preserve most of the detail specified. If the description of an item lacked detail, such as the type (e.g., type of cookie or meat), method of preparation (e.g., raw or cooked), or accompaniment (e.g., butter), the item was assigned a code for items lacking such specification.

Mixtures were usually classified in the food group of the major ingredient. Some mixtures such as salads, sandwiches, and frozen meals were reported either as a single item or by listing component parts. Some food items, such as peanut butter (table 6.05), are tabulated in this report only when they were reported separately, even though the food item might also have been a component of a sandwich or other mixture reported as a single unit. Therefore, the peanut butter in a sandwich reported as a unit is not captured, and peanut butter is underreported in the table.

Participants in the survey were asked to report amounts of foods they ate in measures most convenient to them, i.e., cups, tablespoons, dimensions, ounces. USDA nutritionists supplied the gram weight conversion factors in the coding manual (5) used by the coders. One ounce is equivalent to 28.35 grams. One cup, unless

otherwise noted in the text, refers to the standard 8-ounce measuring cup. The gram weights of common measures of food reported in the survey came mostly from Agricultural Handbook No. 456 (1), available revisions of Agricultural Handbook No. 8 (2-4), other USDA bulletins (7, 9), food industry, and actual determinations. If the amount eaten was inadequately specified, a standard portion size was assigned and identified. The weights in grams per food measure mentioned in the food descriptions in section III of this publication are those given in the coding manual. The number of times a food item is reported refers to the number of eating occasions in which it was consumed and not to the number of individual entries for that item. If a food item was entered more than once for a single eating occasion (such as milk on cereal and in coffee), the amounts on the dietary forms were summed and the total was the portion for that eating occasion. All maximum food consumption values were checked on the original dietary forms to be sure they were not errors caused by coding or transcription.

Selection of Foods for Tables

The food items and food groups selected for inclusion in this publication were those most frequently reported by survey participants. Most were mentioned on at least 1,000 occasions. The text provides a brief description of the foods included in each table. Several tables include all items in a major food group, for example, meat, poultry, and fish (table 7.02). Subsequent tables include items in a subgroup, such as beef, pork, lamb, and veal (table 7.03), which is followed by a table for each component--such as beef (table 7.04)--of the more cumulative table. Another level of subgrouping is provided in tables for beefsteaks (7.05), beef roasts (table 7.06), and ground beef (table 7.07). Unless specifically mentioned in the text, dietetic foods, such as artificially sweetened or low-sodium foods, were not included in the tables.

Content of Tables

For each table the numbers of individuals in total and in each of the 16 sex-age categories are identified. Five sets of intake data for users of the food or food group were computed. These intake data are based solely on persons who consumed the food item, and all averages reported are for the group of users only. The relevant formulas for the computations are given in appendix B. The first data set presents the percentages of individuals using the food on only 1 day, on only 2 days, and on all 3 of the 3 days for which data were collected. The second set includes the average number of eating occasions for the food, the standard deviation, and the maximum number of occasions at which the food was consumed during the 3 days.

In the third data set, the average quantity of food consumed per user per day and the standard deviation are given for individuals who used the food on only 1 day, on only 2 days, and on all 3 days reported. Also presented are the maximum quantity of the food item consumed on any 1 day and the maximum quantity consumed over 3 days. The average quantities of food per user per day are identified at specified percentiles in the distribution of the intakes of each food or food group. Percentile data for foods consumed by fewer than 50 individuals in a sex-age group are not reported. The fourth set of data consists of the average quantities, the standard

deviations, and the maximum quantities consumed by users per eating occasion. Also, the average quantities per eating occasion at specified percentiles in distributions of at least 50 individuals are reported. The last set of data has the number and the percentage of eating occasions for which portion sizes were assigned because amounts eaten were inadequately described. These assigned portion sizes are included in all calculations except in the fourth set of data, the average quantities consumed per eating occasion.

Additional information concerning foods consumed by individuals in the NFCS 1977-78 has been published (6, 11, 12).

DESCRIPTION OF FOODS

Beverages

Separate tables are included in the beverage section for coffee, tea, soft drinks, fruit ades and drinks, beer, wine, and distilled liquors. Milk and milk drinks are reported in the "Milk and Milk Products" section, fruit juices in "Fruits and Fruit Juices," and vegetable juices in the "Vegetables" section.

Coffee (table 1.01) includes liquid coffee made from ground roasted coffee, powdered instant coffee, and liquid coffee concentrate and coffee that was either acid-neutralized or decaffeinated. Usually sugar was reported separately, but some coffee included in this table had sugar added. Not included are turkish coffee; espresso; coffee made from instant cafe au lait; coffee combined with chicory, cereal, or cocoa; unreconstituted instant or liquid concentrate coffee; and coffee substitutes such as Postum. For all eating occasions with coffee, coffee prepared from ground roasted beans was specified for about 41 percent and coffee prepared from instant for 34 percent. Type of coffee was not specified for 25 percent of the eating occasions. About 10 percent of the eating occasions with coffee were for decaffeinated coffee (table 1.02). Of the occasions with decaffeinated coffee, 66 percent were for coffee prepared from powdered instant. One 6-fluid-ounce cup of coffee weighs 180-182 grams.

Tea (table 1.03) includes tea made from tea leaves (loose and in bags), powdered instant, and frozen concentrate and tea sold ready-to-drink in cans. On 46 percent of the occasions the type of tea was not specified. Tea for 34 percent of the occasions was identified as tea prepared from tea leaves and 20 percent as tea made from powdered instant. On less than 1 percent of the occasions tea was identified as ready-to-drink or prepared from frozen concentrate. Data in this table include tea with added cream or milk, sweeteners, or lemon but not the few reports of tea with added chocolate. Tea prepared from herbs, such as sassafras or mint, is not included. A 6-fluid-ounce cup of tea weighs 180-182 grams.

About two-thirds of the carbonated soft drinks (table 1.04) were identified as cola-type carbonated soft drinks (table 1.05) and 14 percent as artificially sweetened carbonated soft drinks (table 1.06). Of the noncarbonated soft drinks (table 1.07), most were prepared from powdered mixes containing sugar and fruit

flavor, including some with added vitamin C. No artificially sweetened noncarbonated soft drinks are included with the noncarbonated soft drinks (table 1.07). Carbonated and non-carbonated soft drinks weigh 356-372 grams per 12-ounce can or 237-248 grams per cup. Fruit ades and drinks (table 1.08) include ready-to-drink beverages as well as beverages made from powdered mixes and frozen concentrates but exclude low-calorie or low-sugar drinks. Fruit ades and drinks weigh 247-256 grams per cup.

Alcoholic beverages tabulated herein are beer and ale, wine, and distilled liquors that are not part of mixtures such as cocktails reported as a unit. Cocktails were not tabulated because of the great diversity in their alcoholic content. Beer and ale (table 1.09) include beer, light beer, and ale. Wine (table 1.10) includes table and dry wines, such as burgundy, claret, chianti, sauterne, rhine, and champagne and sweet or dessert wines such as marsala, port, tokay, madeira, muscatel, angelica, sherry, and sweet vermouth. Cooking wines are not included in this table. Distilled liquors (table 1.11) include brandy, whiskey, gin, vodka, and rum when reported separately but not in cocktails reported as a single item. Beer, wine, and distilled liquors weigh 28-30 grams per fluid ounce. One 12-ounce can or bottle of beer weighs 360 grams, and one glass of wine (3.5 fluid ounces) weighs 100 grams.

Eggs

Eggs (table 2.01) include eggs cooked by all methods, raw eggs, baby-food egg yolks, and mixtures having egg as the main ingredient such as omelets, creamed and deviled eggs, and egg salads. Excluded are meringues, egg soups and sandwiches, and egg substitutes. Intakes of the most popular forms in which eggs are prepared are reported in the following three tables. One large boiled or poached egg (table 2.02) weighs 50 grams, one large fried egg (table 2.03) 46 grams, and one large egg scrambled or made into a plain omelet (table 2.04) 64 grams (2).

Fats and Oils

Fats and oils tabulated include butter, margarine, and salad dressings when these items were reported separately. Fats in mixtures and fats used in cooking, such as deep-fat frying, are not included in these tables nor is salad dressing when a salad with dressing was reported as a unit, i.e., the amount of salad dressing was not reported separately by the respondent.

Butter (table 3.01) does not include whipped butter or butter-margarine blends. Margarine (table 3.02) includes unsalted margarine but does not include liquid, diet, whipped, or imitation margarine or butter-margarine blends. Butter and margarine weigh 14 grams per tablespoon or 5 grams per pat (1 inch square, 1/3 inch high).

Whipped butter and margarine (table 3.03) weigh 9 grams per tablespoon.

Salad dressings and mayonnaise (table 3.04) include all salad dressings and mayonnaise except the low-calorie types. Salad dressings and mayonnaise weigh 14-16 grams per tablespoon. About 42 percent of the reports of salad dressing and mayonnaise were

mayonnaise, 13 percent italian dressing, 11 percent french dressing, 9 percent mayonnaise-type salad dressing, 7 percent thousand island dressing, and 11 percent other types of salad dressing. The remaining 7 percent was not specified as to type.

Mayonnaise (table 3.05) includes regular mayonnaise only and does not include mayonnaise-type salad dressing, mayonnaise made with yogurt, and imitation mayonnaise. These three items are included with salad dressings and mayonnaise (table 3.04). Salad dressings (table 3.06) include all the items in table 3.04 except regular mayonnaise. The most frequently consumed salad dressings were the mayonnaise type (table 3.07), french dressing (table 3.08), and italian dressing (table 3.09). French dressing includes catalina, ranch style, and homemade but does not include low-calorie french dressing. Italian dressing includes california onion and green onion dressings but does not include low-calorie italian dressing.

Fruit and Fruit Juices

Citrus fruit

Only the most frequently reported citrus fruits and citrus fruit juices were tabulated. Fruit juice combinations and mixtures containing citrus fruit, such as gelatin desserts, fruit cup, and salads, were excluded. Fresh grapefruit (table 4.01) includes halves and fresh sections with juice. Almost all reports of grapefruit were for fresh halves. One-half medium grapefruit was calculated as yielding 134 grams of edible portion (9), and 1 cup of sections with juice weighs 230 grams (1). Grapefruit juice (table 4.02) encompasses fresh, reconstituted from frozen, and canned. For 25 percent of the occasions canned grapefruit juice was specified, for 16 percent fresh or frozen was specified, and for 58 percent the form was not specified. On 44 percent of the occasions the juice was specified as unsweetened. Fresh oranges (table 4.03) include all varieties. One medium orange was calculated to yield 145 grams of edible portion and 1 cup of sections to weigh 195 grams. Orange juice (table 4.04) covers all forms--fresh, reconstituted from frozen, and canned, sweetened and unsweetened. Of the reported orange juices, nearly one-half (48 percent) were reconstituted frozen, 17 percent were fresh, 4 percent were canned, and the rest were not specified. Nearly half (49 percent) of the reports specified unsweetened orange juice. One cup of orange juice weighs 248-250 grams, depending upon whether sugar is added.

Noncitrus fruit

Noncitrus fruits and fruit juices reported most often were tabulated. Raw apples (table 4.05) include all varieties, both peeled and unpeeled, but do not include apples in such mixtures as waldorf salad. A medium apple was calculated to provide an edible portion of 138 grams. Applesauce and cooked apples (table 4.06) include apples that are baked, fried, pickled, stewed, and prepared for babies, but most of the items (85 percent) were applesauce. One cup of applesauce with sugar weighs 255 grams and 1 cup unsweetened weighs 244 grams. Applejuice (table 4.07) excludes apple cider and juices that are only part applejuice, such as cranberry-applejuice. About 7 percent of

the reports of applejuice specified baby-food applejuice. One cup weighs 248 grams.

Bananas (table 4.08) include only raw bananas. The yield from one medium banana was calculated as 119 grams (9), and 1 cup of sliced bananas weighs 150 grams (1). Cantaloup (table 4.09) refers to fresh only and excludes the few reports of frozen balls. One-half cantaloup was calculated to yield 272 grams of edible portion (9), and 1 cup of fresh cubes to weigh 160 grams (1). Fruit cocktail (table 4.10) is mainly the canned or cooked type and includes canned fruit salad and fruit compote. Raw fruit cup and raw fruit cocktail are excluded. One cup of canned fruit cocktail with heavy sirup weighs 255 grams. Grapejuice (table 4.11) includes all forms, canned and frozen concentrate reconstituted with water, sweetened and unsweetened, with and without added vitamin C. Grapeade and grape drink are reported with fruit ades and fruit drinks (table 1.09). One cup of grapejuice weighs 253 grams.

Peaches were tabulated for both the raw and the cooked, canned, and frozen forms. Raw peaches (table 4.12) accounted for 35 percent of the occasions on which peaches were consumed. One medium raw peach (about 2-3/4 inches in diameter) was calculated to yield 152 grams of edible portion, and 1 cup of sliced peaches weighs 178 grams. Cooked, canned, and frozen peaches (table 4.13) include peaches not specified as to form (about 13 percent of those mentioned) and commercially prepared baby food (8 percent). On about 75 percent of the occasions entries were identified as canned or cooked and on about 2 percent as frozen. Of the peaches reported as canned or cooked (excluding the baby-food items), 27 percent were in light sirup, 9 percent in heavy sirup, 4 percent unsweetened, and the type of liquid was not mentioned for the rest. One cup of canned or cooked peaches with heavy sirup weighs 256 grams, 1 cup with light sirup 250 grams, 1 cup unsweetened 244 grams, and 1 cup drained halves 224 grams. Peaches in cobblers, pies, and other mixtures are not included in these two tables.

Pears, like peaches, were consumed raw almost as often as cooked or canned. Raw pears (table 4.14) comprised 46 percent of the pear items in this and the next table combined. One medium pear was calculated to provide an edible portion of 164 grams. Cooked and canned pears (also baby-food pears) (table 4.15) include 5 percent of the pear items for which the form (raw, cooked, or canned) was not specified. Of the cooked or canned items (excluding baby food), 71 percent lacked a description of the sirup, 19 percent had light sirup, 5 percent had heavy sirup, and 5 percent were specified as unsweetened. Pears in combination with other fruits such as the baby-food pears and pineapple are not included in these tables. One cup of canned pear halves with heavy sirup weighs 255 grams.

Pineapple was specified as cooked or canned in 64 percent of the reports of this food. Pineapple items were not specified as to type in 20 percent of the reports and were specified as raw for the remainder. Cooked and canned pineapple (table 4.16) includes all pineapple items not reported as raw. Of these cooked and

canned items, 10 percent were reported as cooked or canned in fruit juice, slightly more (12 percent) as unsweetened, and the remainder did not provide such information. One cup of canned pineapple (chunk, tidbit, or crushed) with heavy sirup weighs 255 grams.

Raisins (table 4.17) were the most frequently reported dried fruit with dried prunes reported about one-half as often and dates and apricots each less than one-tenth as often. One cup of raisins weighs 145 grams. Raisins in mixtures are not included in the table.

Among berries, strawberries were reported most often, followed by cranberries, which were mentioned about one-half as often, and blueberries and raspberries each mentioned about one-tenth as often. Of strawberry items reported (excluding mixtures), 77 percent were specified as raw, 19 percent as frozen, and the remaining 4 percent were cooked, canned, or not specified. Quantities for raw strawberries (table 4.18) only are tabulated. One cup of raw strawberries weighs 149 grams.

Grain Products

Yeast breads

Yeast breads (table 5.01) include all toasted and not toasted breads, rolls, and sweet rolls. Bread and rolls used in sandwiches with ingredients reported separately by the respondent are included here. Bread and rolls used in sandwiches with amounts of ingredients not reported separately are excluded. Also excluded from table 5.01 are such mixtures as french toast. Tables 5.02-5.08 treat separately several of the products included in table 5.01.

White breads (table 5.02) are plain and toasted white breads including sourdough, french, italian, portuguese, and vienna breads, and bread that was not identified as to type. About 80 percent were reported as "white bread" and 13 percent as "bread," i.e., type was not specified. Excluded are flat breads, cheese breads, egg breads, and other specialty breads. Most breads and toasts weigh 20-25 grams per medium slice. White rolls (table 5.03) are plain and toasted white rolls, including hard, soft, sourdough, french, vienna, garlic, submarine, and sandwich rolls, and rolls that were not specified as to type. Most medium-sized rolls weigh 25-37 grams, and a submarine roll 100 grams. Excluded are rolls that were used in sandwiches reported as a single unit (ingredients and amounts not listed individually by the respondent). Also excluded are cheese, egg, and sweet rolls, bagels, and bread sticks. The bread group identified as french, vienna, and italian bread (table 5.04) also includes toast, rolls, and portuguese, cuban, and spanish varieties. These breads weigh 20-30 grams per slice, and a medium roll weighs 45 grams.

Whole and cracked wheat breads (table 5.05) include plain and toasted, whole and cracked wheat bread and rolls. Excluded are flat bread and bagels. Table 5.06 has plain and toasted rye, pumpernickel, and black bread and rolls, excluding bagels. High fiber, wheat germ, bran, oatmeal, and multigrain breads (table

5.07) include plain and toasted breads made with other types of flours and grains such as barley, buckwheat, oatmeal, triticale, and soy or with added parts of grains such as bran and wheat germ. Breads and toast in tables 5.05-5.07 weigh 23-28 grams per slice.

English muffins (table 5.08) are those made with white, whole wheat, or cracked wheat flour; 95 percent were white or not specified as to type. One english muffin weighs 57 grams, and one toasted english muffin 50 grams.

Quick breads and tortillas

The next five tables report data for quick breads. Biscuits (table 5.09) exclude whole wheat and cheese biscuits and those made from refrigerated dough. One medium biscuit was assigned a weight of 38 grams. Information on all types of biscuits made from refrigerated dough is found in table 5.10. The weight of a regular-sized biscuit is 20 grams. Cornbread (table 5.11) includes cornbread (86 percent of the occasions) and corn muffins, sticks, and rounds. Excluded are other cornmeal products such as cornbread stuffing, corn pone, hush puppies, and spoonbread. One piece of cornbread (2-1/2 inches by 2-1/2 inches by 1-1/2 inches) weighs 78 grams.

All varieties of pancakes except crepes and chinese pancakes are included in table 5.12, and waffles are tabulated in table 5.13. One medium pancake weighs 27 grams, and one medium waffle 50 grams.

Tortillas (table 5.14) are only those that were reported separately and include corn, wheat flour, and whole wheat flour tortillas. Excluded are tortillas that were a part of mixtures reported as units such as tacos and burritos. Tortillas weigh 30-35 grams each.

Cakes, cookies, pies, doughnuts, and sweet rolls

Cakes and cupcakes (table 5.15) include fruitcakes, poundcakes, and shortcakes with fruit. Excluded are coffeecakes. Of the items reported, 80 percent were servings of cake and 20 percent were cupcakes. Of the cakes and cupcakes, 42 percent were chocolate, 11 percent were yellow, 9 percent were white, 35 percent were other flavors, and 3 percent were not specified as to flavor. Chocolate cakes and cupcakes (table 5.16) include chocolate, devil's food, and fudge-flavored cakes and cupcakes. One piece of cake (one-twelfth of a two-layer frosted cake) weighs 92 grams, and one frosted cupcake (2-1/2 inches diameter) 36 grams.

All varieties of cookies are combined in table 5.17. Graham crackers are not included but appear with crackers--tables 5.27 and 5.28. The four most frequently reported kinds of cookies are tabulated separately (tables 5.18-5.21). All five tables exclude dietetic cookies. Chocolate chip cookies (table 5.18) include those with and without such additional ingredients as nuts and raisins. One medium chocolate chip cookie was given a weight of 7 grams. Oatmeal cookies (table 5.19) include those with such additional ingredients as raisins and chocolate chips,

but oatmeal sandwich and oatmeal peanut butter cookies are excluded. One oatmeal cookie was given a weight of 13 grams. Peanut butter cookies (table 5.20) include peanut sandwich cookies and wafers and all other varieties of peanut cookies except those with oatmeal. One peanut butter cookie was given a weight of 12 grams. Sugar cookies (table 5.21) include those with such added ingredients as nuts, fruit, spice, sour cream, and jelly topping. Excluded are frosted sugar cookies and sugar wafers. One sugar cookie was given a weight of 16 grams.

Pies (table 5.22) include all types of pies and tarts. Fruit pies (table 5.23) include tarts, fried pies, and pies that were not specified as to kind by the respondent (3 percent of the reports of pies). Of the fruit pies, 56 percent were apple and 16 percent were cherry. Of the cream and custard-type pies and tarts (table 5.24), 27 percent were pumpkin and 18 percent were chocolate cream. Neither table 5.23 nor 5.24 includes chiffon, meringue, or pecan pies. A tart weighs 100 grams, a slice of fruit pie 135 grams, and one slice of cream or custard pie 130 grams.

All varieties of doughnuts are included in table 5.25 and sweet rolls, excluding danish pastries, in table 5.26. Most doughnuts weigh 42-57 grams, and one medium sweet roll 50 grams.

Crackers and snacks

Crackers (table 5.27) are of all types--graham crackers (plain, cinnamon, and chocolate covered), saltines, butter crackers, cheese crackers, toast thins, matzo, flat bread, and peanut butter sandwich crackers. Graham crackers (table 5.28) include the honey and cinnamon varieties but not the chocolate covered. One graham cracker square (or two small rectangular pieces) weighs 14 grams. Saltine and soda crackers (table 5.29) include those with seeds but exclude low-sodium crackers. One square saltine or soda cracker weighs 3 grams. With butter crackers (table 5.30) are all butter-type crackers except cheese. One round cracker weighs 3 grams and a rectangular cracker 4 grams.

Popcorn (table 5.31) includes plain, "buttered," salted, and popcorn with cheese, but not popcorn with sugar or caramel sirup. One cup weighs 6-9 grams. Pretzels (table 5.32) exclude soft pretzels and baby pretzels. One three-ring pretzel weighs 3 grams. Potato chips, flavored potato chips, and chips made from granulated potatoes are reported in table 5.33. One cup of potato chips weighs 18 grams. Corn-based salty snacks (table 5.34) include corn chips, tortilla chips, puffed corn, and toasted corn. One cup weighs 35-40 grams, depending upon the type.

Pasta, pasta mixtures, and pizza

Pasta (table 5.35) includes cooked macaroni, noodles (egg, whole wheat, and spinach), and spaghetti. Pasta may be enriched or unenriched and with or without added fat. Chow mein noodles, rice noodles, and such pasta mixtures as macaroni and cheese are not included. The pasta was spaghetti (table 5.36) on 48 percent of the occasions, noodles (table 5.37) on 30 percent, and

macaroni (table 5.38) on the other 21 percent. Cooked pastas weigh 140-160 grams per cup.

Three frequently reported pasta mixtures are tabulated. Spaghetti with tomato sauce (table 5.39) also includes macaroni and noodles mixed with tomatoes or tomato sauce with or without meat and cheese. One cup weighs 250 grams. Macaroni and cheese (table 5.40) also includes macaroni casserole and egg noodle with cottage cheese. One cup of homemade macaroni and cheese weighs 200 grams, and 1 cup from mix or canned 240 grams. Lasagne (table 5.41) weighs 250 grams per cup.

Pizza (table 5.42) includes all kinds except thick crust pizzas. A pizza 5 inches in diameter weighs 84-94 grams, and one 12 inches in diameter 486-543 grams.

Rice and cooked cereals

White rice (table 5.43) includes yellow rice but not sweet rice, brown rice, wild rice, oriental long rice (transparent noodles), or rice cooked with milk. Cream of rice and rice mixtures also are not part of this category. The majority of the reports of rice (84 percent) were not specified as to type, i.e., regular cooking, instant, or converted. Cooked rice weighs 165-175 grams per cup.

In the cooked cereals group (table 5.44) are grits, oatmeal, barley, cream of rice, bulgar, and wheat cereals. Cereals may be regular, quick cooking, or instant, made with water or milk, and with or without added fat. Not included are cornmeal mush, cooked cornstarch, and millet. Of the cooked cereals reported, 34 percent were grits or hominy (table 5.45), 43 percent were oatmeal (table 5.46), and 21 percent were wheat cereals (table 5.47) such as cream of wheat, whole wheat cereal, and cocoa wheat cereal. Wheat cereals do not include bulgar or wheat pilaf. The weights of 1 cup of cereal are as follows: grits, 220-245 grams; oatmeal, 240 grams; cream of rice, 245 grams; wheat cereals, 245 grams; and bulgar, 135 grams.

Ready-to-eat cereals

Ready-to-eat cereals are categorized by cup weight and by sugar content as a percent of dry weight. Data on sugar content were obtained from published reports (8) and information supplied by manufacturers. The general category of ready-to-eat cereals (table 5.48) includes all the cereals tabulated separately in tables 5.49 through 5.55 plus dry cereals not specified as to type and dry baby cereals. Baby cereals in jars and breakfast bars are not included.

Ready-to-eat cereals, 15-21 grams per cup, all sugar levels (table 5.49) encompass Corn Kix, Honey Combs, puffed corn, puffed oats, puffed rice, puffed wheat, Quisp, and Special K. In ready-to-eat cereals, 25-32 grams per cup, 30 percent or more sugar (table 5.50) are Alpha Bits, Apple Jacks, Boo Berry, Coco Pebbles, Cocoa Puffs, Cookie Crisp, Corny Snaps, Count Chocula, Crazy Cow, Frankenberry, Freakies, Frosty O's, Fruit Brute, Fruit Loops,

Fruity Pebbles, Golden Grahams, Kaboom, Lucky Charms, Sugar Pops, and Trix. Ready-to-eat cereals, 25-32 grams per cup, less than 15 percent sugar (table 5.51) include unprocessed bran, Cheerios, Corn Chex, corn flakes, Crispy Rice, Pep, Post Toasties, Rice Chex, Rice Krispies, Team, Total, and Wheaties.

Categorized as ready-to-eat cereals, 35-45 grams per cup, 30 percent or more sugar (table 5.52) are Captain Crunch, Captain Crunch with Peanut Butter, Cocoa Krispies, Country Crisp, Frosted Rice, Sugar Frosted Flakes, Sugar Smacks, and Super Sugar Crisp. Ready-to-eat cereals, 35-45 grams per cup, less than 20 percent sugar (table 5.53) include Bran Chex, Bran Flakes, Buc Wheats, 40% Bran, Grape Nut Flakes, Life, Product 19, and Wheat Chex. The group of ready-to-eat cereals, 50-75 grams per cup, all sugar levels (table 5.54) is comprised of All Bran, Bran Buds, Cinnamon Frosted Mini Wheats, Frosted Mini Wheats, 100% Bran, Raisin Bran, and Shredded Wheat. The group of ready-to-eat cereals, 80 grams or more per cup, all sugar levels (table 5.55) includes Alpen, Concentrate, Country Morning, cracked whole wheat, Cracklin' Bran, C. W. Post, Familia, Golden Harvest, Granola, Granola Puffs, Grape Nuts, Heartland, 100% Natural, Pernola, Proteinola, and wheat germ.

Legumes and Nuts

Of the cooked dried beans (table 6.01) reported, including canned, 46 percent were pinto beans, 19 percent kidney beans, 14 percent lima beans, and 13 percent white beans such as navy beans and great northern beans. The remaining beans reported were black, soy, mungo, or not specified as to type. This table does not include soy-derived products, soy-based infant formulas, or such bean mixtures as baked beans, stewed beans, and bean soups. Immature lima beans are reported with cooked vegetables (table 10.23). Cooked dried beans weigh 185-190 grams per cup.

Baked beans (table 6.02) are bean mixtures in which dried beans are cooked with sauce (tomato, chili, barbecue, sweet and sour), sugar, fat, or meat. Stewed beans and other bean mixtures are not included in this table. Baked beans weigh 255 grams per cup.

Nuts (table 6.03) include all nuts except coconut and nuts mixed with fruit or seeds, and peanuts accounted for 54 percent of the reports. Other frequently reported nuts were almonds (8 percent of the eating occasions), cashews (4 percent), pecans (6 percent), and walnuts (6 percent). The cup weight of nuts varies from 120 to 160 grams. Almonds, pecan halves, and peanuts weigh about 1 gram each, and cashews, butternuts, ginko nuts, and english walnut halves about 2 grams each.

Peanuts (table 6.04) include regular and dry roasted peanuts but not boiled peanuts. Peanut butter (table 6.05) does not include low-sodium or imitation peanut butter or the peanut butter portion of sandwiches when the peanut butter sandwich was reported as a unit. Peanut butter weighs 16 grams per tablespoon.

Meat, Poultry, and Fish

Meat, poultry, and fish

Among meat, poultry, fish, and their mixtures (table 7.01) are plain and breaded meat, poultry, and fish, and foods that are combinations of meat and grains or vegetables but have meat as

the main ingredient. Baby foods are included as are frozen plate meals when they were reported as a unit. Meat soups (table 7.35) and gravies are not in this group nor are mixtures having some meat but not as a major component. All foods tabulated in tables 7.02-7.27 are included in table 7.01 as are other meats and meat mixtures not separately tabulated.

Meat, poultry, and fish (table 7.02) include plain and breaded meat, poultry, and fish and baby-food meats. Food items that are combinations of meat and grain products or vegetables are omitted as are frozen plate meals (see table 7.33), soups (see table 7.35), and gravies.

Beef, pork, lamb, and veal (table 7.03) include all the foods listed in the individual beef, pork, lamb, and veal tables, which are described below. Excluded are organ meats, sausages, luncheon meats, spreads, and mixtures. Of the items reported in this table, 57 percent are beef and 41 percent are pork. Lamb and veal account for only 1 percent each.

In the beef category reported in table 7.04 are beef steaks, roasts, stew meat, ground beef, brisket, corned beef, pastrami, beef bacon, dried chipped beef, and other types of beef including oxtails, neckbones, shortribs, and baby-food and junior beef and beef sticks. Excluded are beef sausages, luncheon meats, meat spreads, organ meats, and mixtures. Separate tables are provided for beef steaks (table 7.05), beef roasts (table 7.06), and ground beef (table 7.07), which includes plain patties and plain meatballs and excludes meatloaf.

The cuts of pork in table 7.08 are chops, steaks, cutlets, and roasts; ham; bacon; spareribs; miscellaneous parts such as pigs' feet and ham hocks; ground, pickled, and dehydrated pork; and baby-food pork, ham, and pork sticks. The table does not include pork sausages, luncheon meats, meat spreads, organ meats, or mixtures.

Pork cutlets are included with pork chops and steaks in table 7.09. Of the pork items reported, 91 percent were pork chops. One medium pork chop was calculated to yield a cooked, edible portion of 80 grams. Ham (table 7.10) includes baby-food ham but excludes fresh ham, i.e., uncured. Most of the pork roasts reported in table 7.11 were fresh, but a few were cured. The few reports of fresh ham are not included in table 7.11. Bacon (table 7.12) includes pork breakfast strips but not fresh, uncured bacon, side pork, fatback, beef bacon, or canadian bacon. One medium slice of cooked bacon weighs 8 grams.

Data for lamb in table 7.13 include a few (less than 1 percent) reports of mutton. The item most frequently reported was lamb chops (62 percent). One medium lamb chop has a cooked, edible yield (lean and fat) of 109 grams. Tables of data on lamb and veal (table 7.14) include baby foods but exclude organ meats and mixtures.

Poultry (table 7.15) includes chicken, turkey, cornish game hen, duck, dove, squab, pigeon, quail, partridge, goose, and

pheasant. Excluded are poultry organ meats, luncheon meats, spreads, and mixtures. Of the reports of poultry, 88 percent were chicken and 11 percent were turkey. Chicken (table 7.16) includes all chicken parts, breaded and floured chicken, and baby-food chicken and chicken sticks, but not cornish game hen, chicken loaf, chicken liver, or chicken spread. Cooked edible yields of three parts were calculated as follows: one-half medium roasted breast, 90 grams; one fried whole leg, 112 grams; and one fried thigh, 61 grams.

Turkey (table 7.17) includes turkey roll, ground turkey, and baby-food turkey and turkey sticks. Excluded are turkey luncheon meats, organ meats, and mixtures.

Liver (table 7.18) includes beef, calf, pork, poultry, rabbit, venison, and baby-food livers. Excluded are fish liver, liver sausage, and such mixtures as chopped liver and liver pate. Of the reports of liver, 45 percent were beef, 18 percent poultry, and 5 percent pork, rabbit, or venison. The remaining 32 percent were not specified as to kind.

The fish and shellfish (table 7.19) include fresh, breaded, floured, canned, raw, and dried fish but not fish mixtures. Finfish (table 7.20) encompasses all varieties of finfish except canned, dried, and raw fish. Fish mixtures are also excluded. The most frequently reported item was fish sticks (13 percent of the finfish reports). Other popular finfish items were flounder (7 percent), haddock (6 percent), and perch (6 percent). Canned tuna (table 7.21) includes only plain tuna in water or in oil. One 7-ounce can of tuna, drained, weighs 169 grams. Tuna salad appears in table 7.31. Shrimp in table 7.22 is breaded and plain but not dried shrimp. A 1/2-cup serving of breaded, fried shrimp weighs 68 grams, and a 1/2-cup serving of steamed shrimp 64 grams.

Processed Meat

Frankfurters (table 7.23) are made from any type of meat or poultry and include a few that are breaded. Junior meat sticks are excluded. A medium frankfurter was assigned a weight of 44 grams. All types of bologna are included in table 7.24, and all types of salami in table 7.25. One slice of either weighs 28 grams. Meats that were part of sandwiches reported as a unit, i.e., the amount of meat was not reported separately by the respondent, are not included in these tables.

Included with pork breakfast sausage (table 7.26) are fresh bulk sausage, patties, and links (about 75 percent of the items reported); brown-and-serve sausage; and country-style and smoked sausage. Sausage that respondents did not identify as to type was assumed to be pork breakfast sausage and was also put in this table. Sausages other than frankfurters, bologna, salami, and pork breakfast sausage (table 7.27) include pork and beef sausage, beef sausage, smoked beef sausage, brown-and-serve beef sausage, smoked beef sausage sticks, blood sausage, bockwurst, bratwurst, capicola, cervelat, chorizos (spanish sausage), head cheese, knoblauch, knockwurst, mortadella, pepperoni, polish sausage (kielbasa), italian sausage, lup chong (chinese smoked pork sausage), mettwurst, souse, thuringer, and vienna sausage.

Polish and Italian were the most frequently reported of these sausages, representing 24 percent and 13 percent of the total number of items, respectively.

Mixtures

Of the many meat, poultry, and fish mixtures consumed by individuals, the most frequently reported were beef stew, chili con carne, meat loaf, tuna salad, pot pies, frozen plate meals, hamburger and cheeseburger sandwiches, and soups. The following weights were assigned: beef stew with vegetables (table 7.28), 245 grams per cup; chili con carne (table 7.29) with or without beans, 255 grams per cup; meatloaf (table 7.30), 11 grams per cubic inch; and tuna salad (table 7.31), 205 grams per cup.

Pot pies (table 7.32) include beef, ham, chicken, turkey, and tuna and weigh 227 grams per 8-ounce frozen pot pie. Of the pot pies reported, 69 percent were chicken or turkey and 28 percent were beef.

Frozen plate meals (table 7.33) include a wide variety of beef, pork, veal, poultry, fish, legume, and pasta entrees obtained frozen then heated to serve. Poultry entrees such as fried chicken or turkey slices with potatoes and mixed vegetables comprised 45 percent of the frozen plate meals reported. Beef entrees such as sliced or chopped beef with potatoes and/or other vegetables accounted for 32 percent of the reports, fish dinners 10 percent, and pasta-based frozen plate meals such as lasagne and macaroni and cheese 5 percent. Frozen plate meals were reported in a variety of package sizes from 7 to 18 ounces (198-510 grams).

Hamburger and cheeseburger sandwiches (table 7.34) include all hamburgers and cheeseburgers reported as a unit with or without condiments such as mustard, catsup, and special sauces and accompaniments such as lettuce, tomatoes, pickles, and bacon. Chiliburgers, tacoburgers, pizzaburgers, and sloppy joes are excluded. Of the total hamburger and cheeseburger sandwiches reported, 75 percent were hamburgers. One hamburger on a bun with regular condiments weighs 97 grams and with accompaniments (lettuce, tomato, pickles) 154 grams. A double hamburger (two patties, double-decker bun) with condiments was given a weight of 183 grams, and a hamburger with one-fourth pound (raw weight) of meat and accompaniments 266 grams. Addition of a slice of cheese increases the weight by 14 grams.

Of the soups reported (table 7.35) 37 percent were vegetable; 27 percent grain mixtures such as chicken noodle soup; 18 percent beef, poultry, or fish; 5 percent legume such as bean or pea soup; and 11 percent tomato, including tomato beef and tomato rice. The remaining 2 percent of soups include pork, lamb, potato, egg drop, cheese, and milk soups, and baby-food soups. Soups included were homemade or prepared from canned or dry mixes. Unreconstituted dry soup mixes and low-sodium soups are not included. One cup (8 fluid ounces) of soup weighs 227-250 grams.

Fluid milk and milk beverages

Fluid milk, milk beverages, and milk-based infant formulas (table 8.01) include whole, lowfat, skim, evaporated, condensed, and reconstituted dry milks; flavored milks such as chocolate milk, cocoa, and malted milk; milk-based infant formulas; buttermilk; and goat's milk. Excluded are eggnog, milk shakes, imitation milks such as soy milk, and meal replacements with milk such as instant breakfast. Of the total reports of fluid milk, milk beverages, and milk-based infant formulas, about 45 percent were whole milk, 16 percent lowfat milk, 4 percent skim milk, 1 percent reconstituted dry milk, 5 percent chocolate milk or cocoa, and 2 percent milk-based infant formulas. The remaining reports were evaporated milk, condensed milk, buttermilk, goat's milk (2 percent), and milk unspecified as to type (24 percent).

Under fluid milk (table 8.02) are whole, lowfat, and skim milk as well as milk not specified as to type. These types of milk are individually reported in tables 8.03, 8.04, and 8.05, respectively. Milk unspecified as to type is included with whole milk (table 8.03). Data for acidophilus and low-sodium whole milks are in table 8.02 but not in table 8.03. Buttermilk, evaporated milk, reconstituted dry milk, flavored milks, milk-based infant formulas, and goat's milk are not included with fluid milk (table 8.02). One cup (8 fluid ounces) of whole fluid milk weighs 244 grams.

Reconstituted dry milk (table 8.06) includes milk reconstituted from whole dry milk (3 percent of the eating occasions), lowfat dry milk (11 percent), and nonfat dry milk (48 percent). Dry milk not specified as to fat content was reported for 38 percent of the occasions. The weight of 1 cup (8 fluid ounces) of reconstituted dry milk is 245 grams.

With chocolate milk (table 8.07) and cocoa and hot chocolate (table 8.08) are beverages made with whole milk (27 percent of the occasions), lowfat milk (7 percent), skim or dry milk (4 percent), and milk not specified as to form and fat content (52 percent). Malted milk beverages are omitted. Chocolate milk, cocoa, and hot chocolate weigh 250-256 grams per cup (8 fluid ounces).

Cheese

Natural and processed cheeses (table 8.09) include natural cheeses such as blue, brick, camembert, brie, cheddar, gouda, edam, limburger, mozzarella, parmesan, and swiss, and processed cheeses such as cream cheeses, processed cheeses, cheese foods, and cheese spreads. Excluded are imitation cheese, cottage cheese, and cheese mixtures such as cheese cakes, fondues, dips, souffles, and soups. Of the reports of natural and processed cheese, 10 percent were cheddar cheese, 6 percent swiss, 8 percent other natural cheeses, 3 percent cream cheese, 50 percent processed cheese, and 23 percent cheeses not specified as to type. Separate tables are provided for cheddar cheese (table 8.10), swiss cheese (table 8.11), natural cheeses other than

cheddar, swiss, and parmesan (table 8.12), and processed cheese, cheese food, and cheese spread (table 8.13). One cubic inch of most cheeses weighs 17 grams.

Of the reports of cottage cheese (table 8.14), 29 percent were for creamed cottage cheese, 10 percent lowfat, 4 percent cottage cheese with fruits and vegetables and baby-food cottage cheese, and 57 percent were not specified as to type. Uncreamed, dry curd cottage cheese is not included in this table. One-half cup of cottage cheese weighs 113 grams.

Milk products

Under fluid cream (table 8.15) are data for light cream (8 percent of the eating occasions), half and half (28 percent), and heavy cream (2 percent), but more than half of the items (62 percent) were not specified as to type. Excluded from this table are powdered cream, whipped cream, and cream substitutes. Of the reports of liquid and frozen cream substitutes (table 8.16), 14 percent were for liquid and 28 percent for frozen; 58 percent were not specified as to type. One tablespoon of fluid cream, liquid cream substitute, or frozen cream substitute weighs 15 grams. Powdered cream substitutes (table 8.17) weigh about 6 grams per tablespoon, 3 grams per packet.

Ice cream and ice milk (table 8.18) include the so-called "hard serve" ice cream and ice milk but exclude "soft serve" ice cream, dietetic and imitation ice cream, and ice cream bars, sandwiches, cones, and sundaes. Of the reports of ice cream and ice milk, 96 percent were for ice cream and 4 percent for ice milk. Data on flavors indicated 16 percent were chocolate ice cream or ice milk, 70 percent flavors other than chocolate, but 14 percent were not specified as to flavor. One cup of ice cream weighs 133 grams, and 1 cup of ice milk 131 grams.

Of reported nondairy whipped toppings (table 8.19), 60 percent were frozen, 15 percent were made from powdered mixes, 4 percent were products in pressurized cans, and 21 percent were not specified as to type. Nondairy whipped toppings weigh 70-80 grams per cup.

Sour cream (table 8.20) includes a few reports of sour half and half but does not include imitation (nondairy) sour cream, filled sour cream, or sour cream in mixtures such as dips. Sour cream weighs 12 grams per tablespoon, 230 grams per cup (2).

Data for all types of yogurt except frozen are in table 8.21. Of reported yogurts, 60 percent were fruit varieties, only 4 percent were vanilla or coffee flavors, and 36 percent were not specified as to flavor. An 8-ounce (avoirdupois) container of yogurt weighs 227 grams and 1 cup 245 grams.

Sugar and Sweets

The sugar and sweets tabulated are those that were most commonly reported: white sugar, sirup, honey, jellies, jams, gelatin desserts, and candies. Under white sugar (table 9.01) are both the granular and lump forms as well as sugar not further specified, rock sugar, and rock candy. Table 9.01 does not include

confectioner's or powdered sugar. One teaspoon of granular white sugar weighs 4 grams, one individual packet 6 grams.

Sirup (table 9.02) includes cane, corn, fruit, maple, and sorghum sirups singly or in combination, buttered blends, and sirups not further specified as to type. Chocolate sirup, caramelized sugar, liquid brown sugar, sugar and water sirup, and dietetic sirups are excluded. One tablespoon of sirup or honey (table 9.03) weighs 20 grams.

Jellies (table 9.04) and jams, preserves, and marmalades (table 9.05) include all fruit flavors but exclude dietetic jellies, jam, and preserves. Jellies weigh 19 grams per tablespoon; jams, preserves, and marmalades 20 grams per tablespoon.

Plain gelatin desserts (table 9.06) and gelatin desserts with fruit (table 9.07) do not include gelatin desserts with added whipped cream, cream cheese, or nuts when desserts with these ingredients were reported as a unit. Dietetic gelatin desserts also are excluded from these tables. One-half cup of gelatin dessert, plain or with fruit, weighs 120 grams.

Candy (table 9.08) includes all types of candy except those specified as dietetic. Of the reports of candy, 59 percent specified chocolate candies or candies containing or coated with chocolate; 14 percent specified hard candies; 6 percent caramels (not chocolate), gumdrops, or taffy; and 4 percent marshmallows. Among other reports of candy were a variety of other kinds and those not specified as to type. One cup of Hershey Kisses, chocolate covered peanuts, or M&M's was assigned a weight of 170 grams, 1 cup of jelly beans (75 pieces) 220 grams, and 1 roll of Life-savers 22 grams. Most medium candy bars weigh 34-51 grams.

Vegetables

Only the most frequently reported vegetables were tabulated. Vegetable mixtures were excluded from tabulations with the exception of white potato mixtures, tomato juices, traditional mixed vegetables, and cabbage, lettuce, and tossed salads.

White potatoes

White potatoes (table 10.01) include all reports of white potatoes except potato chips and sticks (table 5.33), which are classified under crackers and snacks, and potato soups. Among the reports of white potatoes, 23 percent each were for mashed potatoes and french fried potatoes. Baked potatoes accounted for 15 percent, boiled potatoes 11 percent, and potato salad 6 percent of the occasions. Creamed, scalloped, and au gratin potatoes; home-fried potatoes; and hash-browned potatoes accounted for 5, 8, and 2 percent, respectively. The data for baked potatoes (table 10.02) pertain to all white potatoes baked in skin except stuffed baked potatoes, which were mentioned only a few times. One medium baked potato yields 93-98 grams of edible weight, excluding peel. Under boiled potatoes (table 10.03) are those with and without added fat and with and without peel but not canned potatoes. A medium boiled potato yields 122-127 grams edible portion without the peel. Creamed, scalloped, and au gratin potatoes (table 10.04) include potatoes cooked in sauce

with and without cheese. One cup weighs 235-245 grams. Mashed potatoes (table 10.05) weigh 210 grams per cup. Information on home-fried and hash-browned potatoes (table 10.06) is for german fried, o'brien, cottage style, and fried from raw as well as hash browned from fresh, frozen, and dry mix. One cup was assigned a weight of 155 grams for hash-browned and 170 grams for home fries. A fast food order of french fried potatoes (table 10.07) weighs about 70 grams. Potato salad (table 10.08) includes german style, and 1 cup weighs 250 grams.

Raw vegetables

Raw vegetables (tables 10.09-10.17) reported separately most often were cabbage, carrots, celery, cucumbers, lettuce, onions, and tomatoes. Other vegetables also were eaten raw but on too few occasions to be tabulated individually.

Raw cabbage salad and coleslaw (table 10.09) include a few salads with apples, raisins, or pineapple, and 1 cup weighs 120-135 grams. Chinese and red cabbage salad or coleslaw and plain cabbage are not included although they were reported a few times. Raw carrots (table 10.10) exclude carrot salads. One medium carrot strip weighs 4 grams. One large outer stalk of raw celery (table 10.11) weighs 40 grams. Raw cucumbers (table 10.12) do not include the few cucumber salads that were reported. One-half cup of cucumber slices weighs 70 grams.

Lettuce and tossed salads (table 10.13) include all the types treated separately in tables 10.14 and 10.15 as well as tossed salads with avocado, cheese, and/or eggs added to lettuce or assorted vegetables. Only data for plain lettuce with and without dressing are in table 10.14. Tossed salads (table 10.15) include salads with lettuce and other assorted vegetables, with and without salad dressing. One cup of plain lettuce weighs 55 grams, lettuce with dressing 65 grams, and tossed salad 60-108 grams because of some heavier ingredients such as tomatoes and cucumbers.

Raw mature onions (table 10.16) include onions when they were reported separately such as slices of onion mentioned as part of sandwiches and as garnishes, but the onions in salads and mixtures often were not reported separately. Green onions were seldom reported and are not included. One slice of onion weighs 18 grams. Raw tomatoes (table 10.17) include cherry, italian, and plum tomatoes. One slice of tomato weighs 30 grams, one cherry tomato 15 grams. Undoubtedly, many of the reports are for slices of tomatoes eaten as part of a sandwich or a salad. For many salads and some sandwiches with tomatoes, however, ingredients were not reported separately; thus raw tomatoes were used more often than shown in the table.

Cooked vegetables

Data on cooked vegetables, including canned, appear in tables 10.18-10.30. Included in all of these tables are vegetables that were eaten with or without added fat, e.g., butter or margarine. Baby-food and junior vegetables are included also. Excluded are

vegetables served with a sauce; creamed, dried, and pickled vegetables; and vegetable souffles and casseroles.

Broccoli (table 10.18) weighs 85-88 grams per 1/2-cup serving. Cabbage (table 10.19) includes white cabbage and cabbage that was not specified as to type. A 1/2-cup serving weighs about 75 grams. Carrots (table 10.20) exclude glazed. One-half cup weighs about 78 grams. Whole kernel corn (table 10.21) includes white and yellow corn on and off the cob. A 1/2-cup serving weighs 83-85 grams. Cream-style corn (table 10.22) includes white and yellow corn, and a 1/2-cup serving weighs 128 grams. Immature lima beans (table 10.23) weigh 85-88 grams per 1/2 cup. Mixed vegetables (table 10.24) include only the traditional mixture of corn, lima beans, peas, green beans, and carrots. Other vegetable combinations are excluded. One-half cup of mixed vegetables weighs 91-94 grams. Crowder peas are included with cowpeas, field peas, and blackeye peas, not dried (table 10.25). A 1/2-cup serving weighs 85-88 grams. For green peas (table 10.26) the weight is 83-85 grams per 1/2 cup.

Spinach (table 10.27) weighs 93-103 grams per 1/2 cup. String beans (table 10.28) include green string beans and pole beans but exclude yellow (wax) and italian beans. A 1/2-cup serving weighs 67-70 grams. With plain cooked and canned sweetpotatoes (table 10.30) are candied sweetpotatoes. One medium baked sweetpotato weighs 114-119 grams, and one medium boiled 151-156 grams. A 1/2-cup serving of canned sweetpotatoes weighs 113-115 grams.

Along with tomato juices (table 10.31) are tomato juice cocktails and combination juices such as tomato with other vegetable juices. One-half cup weighs 121-122 grams.

Condiments

The last three tables cover the most frequently reported condiments: catsup (table 10.32), mustard (table 10.33), and cucumber pickles (table 10.34), both sweet and sour, but not pickle relish. One tablespoon of catsup or mustard weighs 15 grams, and one sour pickle spear 30 grams.

LITERATURE CITED

- (1) Adams, C. F.
1975 Nutritive value of American foods in common units. U.S. Dept. Agr. Agr. Handb. 456, 291 pp.
- (2) Consumer and Food Economics Institute.
1976 Composition of foods: Dairy and egg products; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-1, 157 pp.
- (3)

1978 Composition of foods: Baby foods; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-3, 231 pp.
- (4)

1979 Composition of foods: Poultry products; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-5, 356 pp.
- (5)

1979 Manual of food codes and conversions of measures and gram-weight for use with individual food intake data from Nationwide Food Consumption Survey 1977-78, U.S. Dept. Agr. CFE (Adm.) 352, 569 pp.
- (6) Consumer Nutrition Center.
1980 Food and nutrient intakes of individuals in 1 day in the United States, spring 1977. Nationwide Food Consumption Survey 1977-78, U.S. Dept. Agr. Prelim. Rpt. No. 2, 121 pp.
- (7) Fulton, L., Matthews, E., and Davis, C.
1977 Average weight of a measured cup of various foods. U.S. Dept. Agr. Home Econ. Res. Rpt. No. 41, 26 pp.
- (8) Li, B.W., and Schuhmann, P. J.
1980 Gas-liquid chromatographic analysis of sugars in ready-to-eat breakfast cereals. J. Food Science. 45:138-141.
- (9) Matthews, R., and Garrison, Y. J.
1975 Food yield summarized by different stages of preparation. U.S. Dept. Agr. Agr. Handb. 102, 136 pp.
- (10) Pao, E. M., and Burk, M. C.
1975 Portion sizes and day's intakes of selected foods. U.S. Dept. Agr. ARS-NE-67, 70 pp.
- (11) Pao, E. M.
1981 Changes in American food consumption patterns and their nutritional significance. Food Technology. 35:43-53.
- (12) Rizek, R. L., and Jackson, E. M.
1980 Current food consumption practices and nutrient sources in the American diet. 35 pp. Consumer Nutrition Center, Science and Education Administration, U.S. Dept. Agr., Hyattsville, Md.

TABLE 1.01.--COFFEE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	51.1	0	1.0	1.2	2.0
ON ONLY 1 OF 3 DAYS.....PCT.....	5.3	0	.3	.7	1.3
ON ONLY 2 OF 3 DAYS.....PCT.....	6.4	0	.4	.5	.4
ON ALL 3 OF 3 DAYS.....PCT.....	39.4	0	.3	.1	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	4.9	0	1.9	1.7	1.5
STANDARD DEVIATION.....DO.....	3.4	0	.8	1.0	.8
MAXIMUM.....DO.....	64.0	0	3.0	6.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	521	0	106	86	107
STANDARD DEVIATION.....GM.....	436	0	73	54	98
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	80				
25TH.....GM.....	240				
50TH.....GM.....	400				
75TH.....GM.....	681				
90TH.....GM.....	1,021				
95TH.....GM.....	1,281				
99TH.....GM.....	2,083				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	311	0	73	161	180
STANDARD DEVIATION.....GM.....	199	0	78	70	90
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	396	0	177	168	208
STANDARD DEVIATION.....GM.....	260	0	46	45	127
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	619	0	195	173	246
STANDARD DEVIATION.....GM.....	445	0	26	100	128
MAXIMUM IN ANY 1 DAY.....GM.....	11,580	0	360	360	600
MAXIMUM OVER 3 DAYS.....GM.....	27,900	0	720	840	1,440
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	317	0	166	155	209
STANDARD DEVIATION.....GM.....	171	0	75	61	118
PERCENTILES: <u>1/</u>					
5TH.....GM.....	180				
25TH.....GM.....	240				
50TH.....GM.....	240				
75TH.....GM.....	480				
90TH.....GM.....	480				
95TH.....GM.....	720				
99TH.....GM.....	961				
MAXIMUM.....GM.....	2,880	0	360	240	600
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	654.1	0	0	1.0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	2.9	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.7	4.1	16.4	17.1	53.9	53.0	84.2	82.4	84.7	85.0	85.1	81.7
2.6	2.2	5.0	6.6	9.5	9.6	5.5	5.0	4.5	4.1	3.0	4.0
1.0	1.1	4.5	4.1	10.6	10.2	8.5	8.5	6.2	6.0	6.3	5.5
1.2	.8	6.8	6.4	33.8	33.2	70.2	68.9	74.1	74.9	75.9	72.2
1.8	1.7	2.6	2.8	4.3	4.2	5.7	5.3	5.2	4.7	5.2	4.7
1.1	.9	1.7	2.3	3.3	3.2	4.1	3.3	3.1	2.4	2.7	2.6
6.0	4.0	16.0	16.0	32.0	29.0	64.0	43.0	29.0	18.0	13.0	32.0
141	129	244	268	474	453	633	566	532	444	484	421
106	88	207	289	435	420	538	418	371	280	277	312
40	40	60	60	80	80	120	120	149	120	160	140
80	80	80	80	180	160	300	260	240	240	240	240
80	80	160	160	360	320	480	480	480	400	480	320
160	160	320	320	640	561	841	721	680	561	641	520
240	240	480	560	1,020	960	1,200	1,061	961	800	880	721
400	240	641	880	1,280	1,200	1,441	1,360	1,201	960	1,040	960
480	480	1,280	1,580	1,921	2,162	2,563	2,082	1,702	1,441	1,282	1,363
254	230	287	264	355	314	324	321	305	274	229	246
114	88	184	120	275	183	174	196	161	165	132	122
210	263	305	381	428	365	448	416	323	386	330	303
66	128	207	292	298	208	293	276	118	212	170	136
269	213	381	462	632	617	715	635	584	478	522	456
134	100	225	355	464	446	548	418	366	277	266	314
840	780	1,680	2,400	7,680	5,520	11,580	4,320	3,360	3,120	1,680	6,480
2,040	1,440	4,320	6,180	15,840	11,760	27,900	12,000	9,120	6,840	4,320	13,440
233	222	286	286	330	321	331	320	307	284	277	271
104	100	131	170	190	190	182	169	142	127	113	116
60	60	149	150	180	180	180	180	180	180	180	150
180	180	240	240	240	240	240	240	240	240	240	240
240	240	240	240	240	240	240	240	240	240	240	240
240	240	240	240	480	480	480	480	360	240	240	240
360	242	480	480	480	480	480	480	480	480	480	480
480	480	480	480	720	720	720	720	480	480	480	480
480	540	721	961	961	961	961	960	721	721	720	720
720	540	960	1,920	2,400	2,400	2,880	2,160	1,440	1,920	1,200	1,440
.8	5.8	4.7	5.8	56.3	99.9	185.1	173.5	29.3	59.5	12.3	20.1
.5	3.8	.8	.8	.6	.8	.8	.6	.6	.9	.5	.5

TABLE 1.02.--DECAFFEINATED COFFEE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.3	0	0	.1	.1
ON ONLY 1 OF 3 DAYS.....PCT.....	1.8	0	0	.1	.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.4	0	0	0	.1
ON ALL 3 OF 3 DAYS.....PCT.....	4.1	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	3.7	0	0	1.0	1.6
STANDARD DEVIATION.....DO.....	2.6	0	0	.0	.5
MAXIMUM.....DO.....	25.0	0	0	1.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	347	0	0	60	85
STANDARD DEVIATION.....GM.....	292	0	0	0	40
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	80				
25TH.....GM.....	160				
50TH.....GM.....	240				
75TH.....GM.....	480				
90TH.....GM.....	720				
95TH.....GM.....	911				
99TH.....GM.....	1,341				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	371	0	0	180	120
STANDARD DEVIATION.....GM.....	285	0	0	0	0
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	407	0	0	0	180
STANDARD DEVIATION.....GM.....	261	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	467	0	0	0	0
STANDARD DEVIATION.....GM.....	314	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	5,520	0	0	180	180
MAXIMUM OVER 3 DAYS.....GM.....	11,760	0	0	180	360
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	282	0	0	180	163
STANDARD DEVIATION.....GM.....	129	0	0	0	27
PERCENTILES: <u>1/</u>					
5TH.....GM.....	180				
25TH.....GM.....	240				
50TH.....GM.....	240				
75TH.....GM.....	240				
90TH.....GM.....	480				
95TH.....GM.....	480				
99TH.....GM.....	721				
MAXIMUM.....GM.....	1,440	0	0	180	180
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	41.5	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
.2	.3	1.2	1.3	2.9	4.4	11.0	13.6	15.1	20.1	20.7	21.3
.1	.2	.5	.7	.9	1.9	2.9	3.2	2.5	2.5	3.2	3.9
0	.1	.3	.5	.9	1.0	2.2	2.6	2.3	3.8	2.4	2.3
.1	0	.4	.2	1.1	1.5	6.0	7.7	10.4	13.8	15.1	15.1
2.0	1.3	2.1	2.2	2.6	2.7	3.8	3.9	4.1	3.8	4.1	3.8
1.0	.5	1.2	1.9	1.6	2.4	2.8	2.8	2.4	2.0	2.5	2.4
3.0	2.0	5.0	9.0	7.0	18.0	24.0	25.0	12.0	13.0	11.0	13.0
132	94	189	171	267	270	372	380	381	323	350	316
72	36	105	159	210	423	304	310	247	188	218	215
				60	60	80	80	80	80	80	80
				120	80	160	160	240	180	240	180
				240	160	240	320	320	240	320	240
				320	280	500	480	540	460	420	400
				561	560	800	800	641	561	720	561
				721	641	961	1,000	721	680	800	721
				881	1,922	1,521	1,361	1,364	880	1,081	1,280
206	222	345	214	352	330	407	392	383	324	308	401
54	43	133	105	297	207	330	330	225	142	149	212
0	208	230	295	324	329	439	457	418	374	376	378
0	43	54	93	132	159	244	343	176	185	119	177
201	0	313	440	438	517	516	509	464	382	418	373
36	0	87	225	230	654	325	319	245	183	211	222
242	240	480	720	1,440	5,520	2,400	3,360	1,920	1,920	1,620	1,680
724	480	1,200	2,160	2,640	11,760	5,760	7,440	4,095	4,320	3,240	3,840
202	215	265	234	305	301	295	291	279	258	255	252
44	44	89	63	156	181	132	141	107	84	74	94
				180	180	180	180	180	180	180	180
				240	240	240	240	240	240	240	240
				240	240	240	240	240	240	240	240
				360	240	300	240	240	240	240	240
				480	480	480	480	480	360	360	360
				720	720	480	480	480	480	480	480
				721	961	721	900	720	480	480	540
242	240	480	480	960	1,440	1,200	1,440	720	720	480	1,080
0	0	0	0	0	1.0	12.4	17.0	8.6	0	.8	1.8
0	0	0	0	0	.2	.6	.5	1.2	0	.2	.2

TABLE 1.03.--TEA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	38.5	4.9	19.9	22.4	24.5
ON ONLY 1 OF 3 DAYS.....PCT.....	12.0	3.1	9.3	9.5	10.1
ON ONLY 2 OF 3 DAYS.....PCT.....	10.1	.7	5.6	6.4	7.5
ON ALL 3 OF 3 DAYS.....PCT.....	16.4	1.1	5.0	6.5	6.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	3.1	3.1	2.4	2.5	2.3
STANDARD DEVIATION.....DO.....	2.3	3.0	1.8	2.1	1.5
MAXIMUM.....DO.....	36.0	13.0	12.0	18.0	11.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	315	118	136	175	190
STANDARD DEVIATION.....GM.....	278	127	109	161	141
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	80		30	40	60
25TH.....GM.....	121		60	80	80
50TH.....GM.....	240		100	160	160
75TH.....GM.....	401		200	221	240
90TH.....GM.....	640		282	360	363
95TH.....GM.....	801		362	403	440
99TH.....GM.....	1,362		480	646	720
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	330	232	200	243	276
STANDARD DEVIATION.....GM.....	200	260	130	151	139
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	379	98	207	243	275
STANDARD DEVIATION.....GM.....	218	68	95	99	104
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	504	264	261	327	340
STANDARD DEVIATION.....GM.....	310	146	121	211	161
MAXIMUM IN ANY 1 DAY.....GM.....	5,280	960	847	1,980	1,331
MAXIMUM OVER 3 DAYS.....GM.....	11,760	1,380	2,178	4,800	2,760
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	302	115	171	207	246
STANDARD DEVIATION.....GM.....	145	64	79	82	90
PERCENTILES: <u>1/</u>					
5TH.....GM.....	180		60	120	120
25TH.....GM.....	240		120	180	182
50TH.....GM.....	240		151	240	240
75TH.....GM.....	360		240	240	242
90TH.....GM.....	480		242	242	360
95TH.....GM.....	484		242	303	480
99TH.....GM.....	912		480	480	541
MAXIMUM.....GM.....	3,840	240	726	1,200	726
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	678.5	4.8	0	5.2	16.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	6.4	0	.5	1.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
27.4	29.3	30.1	33.5	38.3	46.8	41.0	48.4	39.5	50.5	32.6	47.3
11.2	11.2	10.7	12.1	13.2	14.9	11.9	13.1	9.4	11.7	9.2	11.5
7.5	8.9	8.4	10.5	11.0	11.5	10.4	11.7	9.7	13.2	7.6	11.7
8.7	9.2	10.9	10.9	14.1	20.4	18.7	23.6	20.5	25.6	15.8	24.2
2.5	2.4	2.8	2.7	3.0	3.3	3.2	3.4	3.2	3.4	3.1	3.4
1.6	1.6	1.9	2.1	2.4	2.6	2.3	2.4	2.1	2.3	2.0	2.3
10.0	10.0	13.0	24.0	36.0	27.0	16.0	22.0	11.0	17.0	10.0	20.0
258	230	332	272	359	332	359	336	329	308	284	296
209	169	287	257	346	294	302	281	277	222	202	214
80	80	80	80	80	80	80	80	80	80	60	80
101	100	160	110	160	121	160	160	160	160	140	160
200	161	240	210	280	240	250	240	240	240	240	240
320	302	440	360	480	441	480	441	440	400	400	400
540	443	645	521	720	661	721	641	641	564	600	561
721	560	921	681	968	840	960	803	841	721	721	641
769	887	1,450	1,200	1,762	1,442	1,441	1,442	1,441	1,120	960	1,041
360	315	395	309	384	331	354	323	328	307	272	325
272	148	254	139	244	198	253	151	129	143	100	161
355	341	434	375	473	403	416	371	313	342	329	330
147	163	236	185	330	231	218	193	119	143	115	164
452	383	561	480	610	530	558	506	487	444	427	421
237	186	328	332	408	323	322	301	300	224	191	218
2,520	2,040	3,360	3,600	4,620	5,280	4,356	3,540	2,400	2,400	1,200	1,920
4,080	3,540	6,120	10,320	11,160	11,760	6,292	9,600	5,760	4,860	3,120	5,040
310	282	362	298	364	298	335	295	305	269	275	262
140	112	188	117	212	131	168	124	118	101	101	83
180	180	180	180	180	180	180	180	180	180	180	180
240	240	240	240	240	240	240	240	240	240	240	240
240	240	300	240	300	240	240	240	240	240	240	240
360	300	480	360	480	360	480	360	360	300	242	240
480	480	484	480	484	480	480	480	480	480	480	360
484	480	721	480	721	484	720	480	480	480	480	480
960	720	968	721	1,080	726	961	721	720	600	541	480
1,440	1,452	1,920	960	3,840	1,920	1,694	1,920	720	960	1,080	968
9.0	11.7	17.2	20.8	55.5	114.3	100.5	183.9	39.1	68.8	4.6	26.4
.6	.8	1.5	1.5	1.2	1.4	1.5	1.6	2.7	2.3	.8	1.7

TABLE 1.04.--CARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	50.8	4.5	40.1	51.4	50.2
ON ONLY 1 OF 3 DAYS.....PCT.....	20.3	2.7	20.1	25.4	25.9
ON ONLY 2 OF 3 DAYS.....PCT.....	15.2	.7	12.5	16.6	14.5
ON ALL 3 OF 3 DAYS.....PCT.....	15.2	1.1	7.6	9.4	9.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	2.6	2.3	2.2	2.2	2.1
STANDARD DEVIATION.....DO.....	2.0	1.8	1.6	1.5	1.5
MAXIMUM.....DO.....	22.0	7.0	13.0	13.0	16.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	274	88	123	153	177
STANDARD DEVIATION.....GM.....	241	75	111	114	129
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	62		21	41	41
25TH.....GM.....	119		51	82	82
50TH.....GM.....	205		82	123	144
75TH.....GM.....	359		164	205	246
90TH.....GM.....	556		247	297	338
95TH.....GM.....	733		328	382	410
99TH.....GM.....	1,193		574	554	657
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	342	145	187	255	297
STANDARD DEVIATION.....GM.....	188	89	105	136	169
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	387	162	197	266	309
STANDARD DEVIATION.....GM.....	197	85	105	129	128
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	503	172	269	295	343
STANDARD DEVIATION.....GM.....	291	83	150	138	146
MAXIMUM IN ANY 1 DAY.....GM.....	5,228	679	1,538	1,107	1,845
MAXIMUM OVER 3 DAYS.....GM.....	10,824	775	3,198	2,276	3,444
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	310	115	169	209	248
STANDARD DEVIATION.....GM.....	124	61	74	89	95
PERCENTILES: <u>1/</u>					
5TH.....GM.....	124		62	93	123
25TH.....GM.....	246		123	124	185
50TH.....GM.....	296		154	185	246
75TH.....GM.....	369		246	246	308
90TH.....GM.....	492		246	310	369
95TH.....GM.....	492		308	369	372
99TH.....GM.....	739		372	492	496
MAXIMUM.....GM.....	1,968	248	496	984	984
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	371.3	0	2.5	13.3	2.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	.3	.7	.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
56.5	56.8	69.1	72.0	69.4	65.9	43.9	46.5	21.0	20.5	18.3	16.3
26.4	25.5	22.4	23.4	22.6	21.6	18.8	20.1	10.0	9.9	7.9	9.2
17.5	18.4	22.8	22.8	20.7	20.2	12.3	13.0	6.4	5.8	5.3	3.4
12.6	13.0	23.9	25.8	26.1	24.1	12.7	13.4	4.6	4.9	5.1	3.7
2.4	2.4	2.9	2.5	3.1	3.0	2.5	2.5	2.2	2.3	2.1	1.9
1.6	1.7	2.0	1.9	2.3	2.1	1.9	1.9	1.5	1.9	1.2	1.3
12.0	20.0	15.0	15.0	17.0	22.0	16.0	19.0	9.0	19.0	6.0	7.0
247	237	352	312	366	313	276	242	218	206	211	163
195	185	271	223	297	264	243	217	165	236	143	125
81	62	82	82	82	79	79	62	62	59	82	41
123	103	164	144	164	124	123	103	103	82	103	82
198	185	277	246	287	246	205	165	164	142	163	119
328	308	453	410	492	410	369	316	287	248	296	205
492	482	700	615	740	616	554	474	489	400	375	328
615	575	862	738	949	797	697	616	492	474	496	410
988	882	1,395	1,087	1,395	1,274	1,215	1,184	862	1,422	657	656
375	344	430	381	410	353	354	322	323	264	316	258
195	172	251	214	231	189	172	156	136	93	111	103
408	386	467	438	465	389	410	346	364	318	350	315
199	169	232	216	249	175	196	150	165	132	151	128
469	447	586	496	610	531	511	454	423	437	354	312
235	228	294	224	316	302	301	277	184	376	150	141
3,690	2,460	2,598	2,224	2,952	3,936	5,228	2,952	1,107	3,336	984	984
5,166	5,904	6,428	4,554	7,565	10,824	9,410	7,110	2,745	9,024	1,968	1,968
314	296	364	321	355	312	330	291	300	273	301	255
119	116	142	115	133	118	129	104	108	91	107	95
185	178	186	185	185	178	185	153	183	123	124	119
246	246	246	246	246	246	246	237	246	237	237	185
296	246	369	308	369	296	308	246	296	246	296	246
369	369	372	365	372	369	369	356	369	310	369	308
492	492	492	492	492	492	492	415	372	369	492	369
492	492	496	492	496	492	492	492	492	474	492	474
739	683	984	738	922	711	744	533	829	496	496	492
1,479	1,538	1,538	1,235	1,488	1,968	1,968	984	948	738	496	492
33.2	17.2	28.3	22.8	59.7	69.9	47.5	55.4	1.9	17.1	0	0
1.2	.6	1.0	.7	.7	.7	.9	.7	.4	2.1	0	0

TABLE 1.05.--COLA-TYPE CARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	36.1	2.2	31.0	37.8	38.2
ON ONLY 1 OF 3 DAYS.....PCT.....	16.7	1.4	17.7	20.9	22.0
ON ONLY 2 OF 3 DAYS.....PCT.....	10.5	.2	8.4	10.9	10.2
ON ALL 3 OF 3 DAYS.....PCT.....	9.0	.7	4.8	6.0	6.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.4	2.4	2.0	2.0	2.0
STANDARD DEVIATION.....DO.....	1.9	1.5	1.5	1.4	1.4
MAXIMUM.....DO.....	22.0	5.0	13.0	13.0	16.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	251	74	113	144	164
STANDARD DEVIATION.....GM.....	230	42	106	110	126
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	62		21	41	41
25TH.....GM.....	103		41	62	82
50TH.....GM.....	164		82	103	123
75TH.....GM.....	328		144	185	226
90TH.....GM.....	492		226	287	328
95TH.....GM.....	666		287	369	390
99TH.....GM.....	1,107		574	553	657
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	342	174	192	249	289
STANDARD DEVIATION.....GM.....	186	106	109	136	157
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	383	154	179	264	299
STANDARD DEVIATION.....GM.....	189	0	78	127	127
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	502	102	280	297	347
STANDARD DEVIATION.....GM.....	299	43	159	134	159
MAXIMUM IN ANY 1 DAY.....GM.....	5,228	400	1,538	984	1,845
MAXIMUM OVER 3 DAYS.....GM.....	10,824	492	3,198	2,276	3,444
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	312	95	170	213	248
STANDARD DEVIATION.....GM.....	121	56	75	87	93
PERCENTILES: <u>1/</u>					
5TH.....GM.....	154		62	123	123
25TH.....GM.....	246		123	123	185
50TH.....GM.....	308		153	185	246
75TH.....GM.....	369		246	246	308
90TH.....GM.....	492		246	308	369
95TH.....GM.....	492		308	369	369
99TH.....GM.....	738		369	492	492
MAXIMUM.....GM.....	1,968	246	492	984	984
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	222.8	0	.8	11.8	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	.1	.9	.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
44.9	44.4	54.8	54.8	57.3	46.1	30.3	26.4	12.8	10.4	9.2	8.2
24.0	22.4	21.6	21.1	21.5	19.5	14.4	13.2	7.1	5.4	4.8	5.3
12.3	13.9	18.1	17.7	16.6	13.5	8.6	7.2	3.5	2.6	2.8	1.5
8.6	8.1	15.1	16.0	19.2	13.1	7.3	6.0	2.3	2.4	1.6	1.5
2.1	2.2	2.6	2.6	2.9	2.7	2.3	2.2	2.0	2.0	2.0	1.8
1.5	1.5	1.9	1.7	2.2	2.1	1.7	1.7	1.4	1.2	1.3	1.1
12.0	20.0	12.0	11.0	17.0	22.0	15.0	19.0	7.0	6.0	6.0	6.0
220	213	314	278	339	278	255	219	204	170	218	153
169	163	243	203	288	264	229	203	149	115	169	128
82	62	82	82	82	62	82	62	62	41		41
103	82	123	123	123	123	123	82	103	82		67
164	164	246	246	246	205	164	164	164	164		103
287	267	410	369	451	369	328	267	246	246		205
431	410	656	534	738	574	492	410	492	328		308
534	534	779	657	903	739	656	554	492	390		410
821	780	1,232	985	1,395	1,209	985	985	616	512		657
371	344	435	379	401	343	363	318	334	252	311	244
186	188	248	190	228	173	168	147	140	115	132	106
387	377	467	430	449	382	394	346	372	321	413	342
196	163	240	211	214	171	172	148	163	105	147	102
434	417	558	469	603	544	508	452	420	319	467	332
201	209	270	217	324	341	310	287	153	90	193	165
3,690	2,460	2,214	2,091	2,952	3,936	5,228	2,952	1,107	738	984	984
5,166	5,904	4,428	3,813	6,704	10,824	9,410	6,888	2,214	1,845	1,968	1,968
308	292	359	321	355	312	333	294	302	256	327	257
108	116	135	112	127	114	126	99	90	79	111	102
185	185	185	185	185	185	185	185	185	123		123
246	246	246	246	246	246	246	246	246	200		185
246	246	369	308	369	246	308	246	308	246		246
369	369	431	369	369	369	369	369	369	308		308
492	492	492	492	492	492	492	492	369	369		369
492	492	492	492	492	492	492	492	492	369		492
615	616	739	677	739	615	739	492	492	492		492
1,107	1,538	1,538	1,230	1,476	1,968	1,968	984	677	738	492	492
19.5	10.6	16.5	14.0	42.6	44.7	25.8	32.7	0	2.2	0	0
1.0	.5	.8	.7	.7	.7	.8	.8	0	.6	0	0

TABLE 1.06.--ARTIFICIALLY SWEETENED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.1	.4	2.5	3.8	3.1
ON ONLY 1 OF 3 DAYS.....PCT.....	4.0	.4	1.6	2.6	2.0
ON ONLY 2 OF 3 DAYS.....PCT.....	2.1	0	.6	1.0	.8
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	0	.3	.3	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.4	1.0	1.6	1.7	1.6
STANDARD DEVIATION.....DO.....	2.0	.0	1.0	.9	.9
MAXIMUM.....DO.....	19.0	1.0	5.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	245	17	82	97	109
STANDARD DEVIATION.....GM.....	232	17	68	64	67
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	59			20	40
25TH.....GM.....	99			59	59
50TH.....GM.....	158			79	79
75TH.....GM.....	316			119	138
90TH.....GM.....	474			217	237
95TH.....GM.....	711			237	237
99TH.....GM.....	1,185			277	287
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	322	52	174	207	226
STANDARD DEVIATION.....GM.....	160	50	72	111	100
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	383	0	123	206	228
STANDARD DEVIATION.....GM.....	192	0	29	84	84
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	499	0	218	208	213
STANDARD DEVIATION.....GM.....	298	0	118	83	76
MAXIMUM IN ANY 1 DAY.....GM.....	2,844	119	533	711	474
MAXIMUM OVER 3 DAYS.....GM.....	8,532	119	1,244	830	860
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	306	52	150	168	200
STANDARD DEVIATION.....GM.....	117	50	56	75	81
PERCENTILES: <u>1/</u>					
5TH.....GM.....	178			59	59
25TH.....GM.....	237			119	147
50TH.....GM.....	296			178	178
75TH.....GM.....	356			237	237
90TH.....GM.....	474			237	356
95TH.....GM.....	474			296	356
99TH.....GM.....	712			356	474
MAXIMUM.....GM.....	1,896	119	356	356	474
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	52.8	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.0	4.7	3.4	11.4	5.2	15.7	7.0	14.2	2.2	5.1	2.2	4.1
2.1	2.7	2.2	6.6	2.8	6.2	3.4	6.8	.7	3.0	.7	2.7
.4	1.3	.6	2.5	1.1	4.4	1.8	3.7	.8	1.0	.5	.8
.5	.7	.5	2.2	1.2	5.1	1.7	3.8	.7	1.1	.9	.6
1.8	1.9	1.8	2.2	2.5	2.7	2.4	2.4	2.5	2.4	2.7	1.6
1.2	1.2	1.1	1.9	2.4	2.0	1.9	2.0	1.4	2.8	1.6	.9
6.0	7.0	5.0	12.0	15.0	12.0	12.0	19.0	6.0	18.0	6.0	4.0
172	189	185	222	273	287	258	243	267	249	193	141
130	141	125	201	257	243	239	228	187	375	87	99
40	59		59	79	59	59	59		79		
79	79		99	119	119	113	119		79		
119	138		158	158	237	158	158		119		
237	237		257	339	359	326	316		237		
435	375		474	612	593	592	474		415		
464	514		633	949	790	712	633		474		
513	633		1,265	1,302	1,224	1,068	1,185		2,844		
298	315	388	340	356	322	315	341	294	292	341	273
140	146	242	165	116	172	111	176	65	102	25	113
449	379	332	375	529	397	416	369	383	357	237	242
182	160	140	131	342	168	190	178	221	144	0	92
370	408	392	520	561	524	537	473	439	652	272	327
70	150	93	260	293	267	294	307	158	630	74	92
948	1,067	1,422	2,133	2,133	2,074	2,489	2,844	948	2,844	533	474
1,541	2,489	1,896	3,911	3,911	5,036	5,333	7,110	2,489	8,532	1,126	1,422
292	304	312	304	330	317	327	298	326	309	218	264
107	109	106	103	107	126	123	108	161	90	70	101
119	178		178	178	178	178	178		178		
237	237		237	237	237	237	237		237		
296	296		296	356	356	356	296		296		
356	356		356	356	356	356	356		356		
356	356		474	474	474	474	356		474		
474	474		474	474	474	474	474		474		
474	711		534	712	712	948	711		533		
948	1,067	474	948	1,067	1,896	1,067	948	948	533	356	474
0	2.2	0	3.1	3.1	12.6	3.5	14.3	0	13.9	0	0
0	1.2	0	.9	.6	.6	.4	.6	0	6.4	0	0

TABLE 1.07.--NONCARBONATED SOFT DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.9	4.2	25.4	30.4	27.1
ON ONLY 1 OF 3 DAYS.....PCT.....	5.5	1.8	11.1	13.3	12.5
ON ONLY 2 OF 3 DAYS.....PCT.....	3.1	1.8	7.3	9.4	8.0
ON ALL 3 OF 3 DAYS.....PCT.....	2.3	.6	7.0	7.6	6.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.2	2.2	2.5	2.6	2.4
STANDARD DEVIATION.....DO.....	1.6	1.2	1.9	1.8	1.6
MAXIMUM.....DO.....	12.0	5.0	12.0	12.0	10.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	216	95	160	192	200
STANDARD DEVIATION.....GM.....	190	70	150	151	154
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	62		41	51	62
25TH.....GM.....	83		62	83	83
50TH.....GM.....	165		103	145	165
75TH.....GM.....	269		207	248	248
90TH.....GM.....	413		331	413	413
95TH.....GM.....	559		434	496	496
99TH.....GM.....	993		826	662	744
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	339	144	198	268	299
STANDARD DEVIATION.....GM.....	219	125	114	128	138
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	360	159	219	290	308
STANDARD DEVIATION.....GM.....	218	74	94	148	127
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	434	214	325	368	383
STANDARD DEVIATION.....GM.....	238	32	184	162	180
MAXIMUM IN ANY 1 DAY.....GM.....	3,968	496	1,488	1,488	1,704
MAXIMUM OVER 3 DAYS.....GM.....	7,378	744	3,720	3,224	3,564
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	290	128	192	225	255
STANDARD DEVIATION.....GM.....	148	64	92	86	93
PERCENTILES: <u>1/</u>					
5TH.....GM.....	124		93	124	124
25TH.....GM.....	248		124	186	248
50TH.....GM.....	248		186	248	248
75TH.....GM.....	310		248	248	248
90TH.....GM.....	496		248	310	372
95TH.....GM.....	496		372	372	496
99TH.....GM.....	868		496	496	496
MAXIMUM.....GM.....	3,968	248	992	744	930
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	42.9	0	3.4	10.6	4.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.5	.8	.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
20.8	19.9	13.5	14.6	9.6	10.5	4.1	4.6	2.0	2.3	2.1	1.0
10.7	9.7	7.2	7.3	5.0	5.9	2.3	2.8	.7	1.3	.7	.6
6.3	6.1	3.9	4.3	2.5	2.8	1.4	1.2	.4	.4	.6	.2
3.9	4.1	2.4	3.0	2.1	1.9	.5	.7	.9	.6	.7	.2
2.2	2.2	2.3	2.1	2.2	2.1	1.8	1.8	3.3	2.1	2.9	1.7
1.6	1.5	1.6	1.4	1.7	1.5	1.3	1.2	2.3	1.6	2.0	.8
10.0	9.0	9.0	8.0	12.0	9.0	7.0	8.0	7.0	7.0	6.0	3.0
238	212	285	226	280	215	222	173	396	231	261	172
197	161	234	170	256	189	250	132	345	226	200	86
83	62	83	83	83	62	83	55				
88	83	103	83	124	83	83	83				
165	165	227	165	207	165	165	124				
310	248	372	289	331	248	248	217				
475	413	496	413	578	413	413	331				
620	579	827	496	703	496	537	413				
993	745	1,117	827	1,158	993	1,591	662				
383	342	458	363	463	341	350	305	325	287	248	392
205	191	305	202	402	174	178	144	93	94	0	244
432	334	520	373	455	393	426	313	316	593	248	403
289	149	287	135	182	183	429	108	52	474	0	0
461	429	585	449	551	463	558	396	713	437	516	201
220	176	266	204	358	267	340	159	293	177	92	41
2,790	1,984	2,480	1,488	3,968	1,984	3,968	1,240	1,736	1,984	930	806
3,906	2,976	3,906	3,224	6,448	3,968	7,378	2,232	2,976	2,728	1,860	806
320	286	371	316	385	310	365	283	356	333	273	298
145	126	181	126	217	127	250	96	190	146	30	122
186	124	218	186	186	186	248	155				
248	248	248	248	248	248	248	248				
248	248	310	248	310	248	279	248				
372	310	496	372	496	372	466	310				
496	496	496	496	496	496	496	496				
496	496	621	496	869	496	621	496				
745	744	993	745	993	745	993	496				
1,488	1,984	1,488	992	1,984	992	3,968	744	992	992	310	558
8.1	6.6	.8	3.2	1.4	2.4	0	1.8	0	0	0	0
.8	.7	.2	.7	.2	.2	0	.3	0	0	0	0

TABLE 1.08.--FRUIT ADES AND DRINKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	13.7	3.5	22.2	22.4	23.6
ON ONLY 1 OF 3 DAYS.....PCT.....	8.2	2.1	10.7	11.9	13.7
ON ONLY 2 OF 3 DAYS.....PCT.....	3.2	.8	6.2	6.2	5.7
ON ALL 3 OF 3 DAYS.....PCT.....	2.3	.7	5.4	4.3	4.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.9	1.8	2.3	2.2	2.0
STANDARD DEVIATION.....DO.....	1.3	.8	1.9	1.5	1.4
MAXIMUM.....DO.....	23.0	3.0	18.0	10.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	170	91	133	146	153
STANDARD DEVIATION.....GM.....	149	77	110	109	113
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	42		21	42	42
25TH.....GM.....	83		62	83	83
50TH.....GM.....	125		83	124	104
75TH.....GM.....	208		167	188	208
90TH.....GM.....	333		250	271	313
95TH.....GM.....	417		394	333	405
99TH.....GM.....	707		531	584	521
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	311	156	207	272	273
STANDARD DEVIATION.....GM.....	188	132	173	157	131
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	321	191	202	247	295
STANDARD DEVIATION.....GM.....	176	109	96	147	138
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	342	172	258	273	299
STANDARD DEVIATION.....GM.....	224	79	121	122	132
MAXIMUM IN ANY 1 DAY.....GM.....	4,464	500	1,250	1,125	1,000
MAXIMUM OVER 3 DAYS.....GM.....	10,416	688	1,625	2,460	2,223
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	269	150	171	202	233
STANDARD DEVIATION.....GM.....	144	80	79	87	101
PERCENTILES: <u>1/</u>					
5TH.....GM.....	125		63	95	125
25TH.....GM.....	188		125	125	186
50TH.....GM.....	250		155	188	249
75TH.....GM.....	310		249	250	250
90TH.....GM.....	496		250	250	372
95TH.....GM.....	500		250	375	496
99TH.....GM.....	750		500	500	500
MAXIMUM.....GM.....	3,000	250	750	750	871
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	41.1	0	3.2	1.1	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	.6	.1	.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
21.3	21.1	18.4	19.0	12.9	13.9	8.5	9.4	6.7	7.9	9.0	7.8
12.2	12.7	9.1	11.5	7.6	9.1	5.8	6.0	4.3	4.6	3.4	4.5
4.3	5.5	5.5	5.5	2.9	3.1	1.4	2.0	1.0	1.5	3.2	1.5
4.8	2.9	3.8	1.9	2.4	1.8	1.3	1.3	1.5	1.8	2.4	1.8
2.0	1.9	2.0	1.8	1.9	1.8	1.7	1.7	1.8	1.9	2.6	1.8
1.4	1.2	1.3	1.0	1.4	1.2	1.1	1.4	1.3	1.2	1.7	1.1
8.0	9.0	7.0	6.0	10.0	9.0	6.0	23.0	7.0	6.0	7.0	6.0
183	165	227	171	223	160	169	157	158	161	217	139
166	123	183	115	192	127	137	185	113	147	135	100
62	62	83	42	63	42	52	42	52	42		42
83	83	103	83	83	83	83	83	83	83		83
125	125	167	125	165	125	125	103	103	104		83
228	208	250	248	271	206	207	167	188	188		167
372	333	457	333	496	313	331	292	333	372		310
500	417	583	414	562	396	417	413	395	375		333
745	625	998	583	1,001	706	833	661	496	1,000		498
318	315	416	327	371	307	339	290	271	321	372	256
190	189	282	163	191	166	183	141	98	386	271	142
327	325	368	364	426	312	361	315	380	257	330	297
169	156	224	147	250	127	170	173	194	144	146	147
347	325	415	329	468	369	342	354	288	290	347	221
242	157	233	149	237	182	206	392	84	143	123	113
3,486	1,500	2,976	1,375	3,750	1,500	1,500	4,464	992	3,000	992	746
6,225	2,486	3,720	1,875	4,875	2,728	2,976	10,416	1,488	3,000	1,619	1,494
277	267	334	292	350	271	300	273	259	255	249	231
138	127	246	122	175	116	137	116	99	233	140	105
125	125	125	125	156	125	125	125	125	124		63
188	247	248	248	249	190	248	247	188	188		154
250	250	250	250	250	250	250	250	249	248		250
310	250	375	372	375	313	374	310	250	250		250
496	375	500	498	500	375	496	496	496	372		313
500	500	625	500	745	500	500	500	496	496		372
751	750	1,500	745	1,000	500	750	625	500	561		563
1,500	1,500	2,976	1,000	1,500	1,488	1,000	750	500	3,000	992	620
7.9	4.6	2.7	2.2	3.4	2.2	6.0	3.1	0	.9	0	2.2
.9	.5	.5	.5	.3	.2	.9	.3	0	.4	0	1.6

TABLE 1.09.--BEER AND ALE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.4	0	.3	.1	0
ON ONLY 1 OF 3 DAYS.....PCT.....	4.3	0	.2	.1	0
ON ONLY 2 OF 3 DAYS.....PCT.....	2.2	0	0	0	0
ON ALL 3 OF 3 DAYS.....PCT.....	1.9	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.4	0	1.6	1.0	0
STANDARD DEVIATION.....DO.....	2.3	0	.9	.0	0
MAXIMUM.....DO.....	27.0	0	3.0	1.0	0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	455	0	37	30	0
STANDARD DEVIATION.....GM.....	556	0	24	0	0
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	80				
25TH.....GM.....	120				
50TH.....GM.....	240				
75TH.....GM.....	600				
90TH.....GM.....	961				
95TH.....GM.....	1,321				
99TH.....GM.....	2,322				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	675	0	64	90	0
STANDARD DEVIATION.....GM.....	748	0	30	0	0
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	733	0	0	0	0
STANDARD DEVIATION.....GM.....	551	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	926	0	70	0	0
STANDARD DEVIATION.....GM.....	842	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	15,120	0	90	90	0
MAXIMUM OVER 3 DAYS.....GM.....	34,560	0	210	90	0
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	557	0	68	90	0
STANDARD DEVIATION.....GM.....	437	0	22	0	0
PERCENTILES: <u>1/</u>					
5TH.....GM.....	240				
25TH.....GM.....	360				
50TH.....GM.....	360				
75TH.....GM.....	720				
90TH.....GM.....	1,080				
95TH.....GM.....	1,441				
99TH.....GM.....	2,163				
MAXIMUM.....GM.....	8,640	0	90	90	0
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	25.5	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
0	(2)	3.9	1.4	26.8	7.8	21.5	5.4	10.8	1.7	7.4	1.3
0	(2)	2.7	1.1	12.8	5.1	9.7	3.3	4.2	.9	3.7	.7
0	0	.9	.4	7.9	1.7	6.1	1.0	2.2	.5	.8	.2
0	0	.4	0	6.2	.9	5.8	1.1	4.4	.3	2.8	.4
0	1.0	2.0	1.3	2.6	1.9	2.7	2.0	2.6	2.0	2.9	2.5
0	.0	1.8	.5	2.5	1.4	2.5	1.5	2.0	1.3	2.5	2.2
0	1.0	11.0	3.0	27.0	10.0	23.0	11.0	10.0	5.0	11.0	7.0
0	120	411	223	550	319	497	279	446	267	346	303
0	0	374	192	715	320	541	233	460	206	302	291
		120		120	80	100	70	110			
		120		160	120	160	120	120			
		240		360	220	360	240	300			
		601		720	420	601	360	661			
		961		1,081	721	1,081	600	921			
		1,200		1,441	960	1,482	721	1,201			
		1,681		2,761	1,441	2,524	1,082	2,641			
0	360	902	446	806	576	716	522	432	332	421	309
0	0	862	339	903	462	840	451	254	133	226	59
0	0	941	677	865	698	685	466	554	535	563	420
0	0	600	307	527	510	626	200	312	180	179	0
0	0	687	0	1,100	745	971	565	775	569	609	631
0	0	475	0	1,188	444	660	252	541	167	320	301
0	360	3,840	1,440	15,120	3,240	11,160	4,320	2,880	1,080	1,440	1,080
0	360	5,040	2,640	34,560	7,200	15,480	4,320	7,920	2,520	3,960	2,520
0	360	604	520	628	511	555	424	511	409	357	366
0	0	474	350	511	369	428	215	342	164	108	139
		240		240	209	240	210	210			
		360		360	360	360	360	360			
		360		360	360	360	360	360			
		721		721	720	720	360	720			
		1,440		1,082	961	1,080	721	1,080			
		1,441		1,621	1,441	1,262	721	1,440			
		2,940		2,164	1,922	2,163	1,082	1,800			
0	360	2,940	1,440	7,200	2,520	8,640	1,440	1,800	1,080	720	720
0	0	0	1.6	3.4	8.0	11.3	0	1.2	0	0	0
0	0	0	5.8	.1	1.0	.4	0	.4	0	0	0

TABLE 1.10.--WINE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	AGE (YEARS) AND SEX OF INDIVIDUAL				
	TOTAL	UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.8	0	0	.1	.2
ON ONLY 1 OF 3 DAYS.....PCT.....	3.1	0	0	.1	.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	0	0	0
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.7	0	0	1.0	1.0
STANDARD DEVIATION.....DO.....	1.1	0	0	.0	.0
MAXIMUM.....DO.....	9.0	0	0	1.0	1.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	113	0	0	56	45
STANDARD DEVIATION.....GM.....	118	0	0	33	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	19				
25TH.....GM.....	39				
50TH.....GM.....	77				
75TH.....GM.....	135				
90TH.....GM.....	232				
95TH.....GM.....	320				
99TH.....GM.....	618				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	221	0	0	167	136
STANDARD DEVIATION.....GM.....	211	0	0	99	80
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	221	0	0	0	0
STANDARD DEVIATION.....GM.....	181	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	237	0	0	0	0
STANDARD DEVIATION.....GM.....	173	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	3,213	0	0	232	232
MAXIMUM OVER 3 DAYS.....GM.....	4,634	0	0	232	232
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	200	0	0	167	136
STANDARD DEVIATION.....GM.....	163	0	0	99	80
PERCENTILES: <u>1/</u>					
5TH.....GM.....	58				
25TH.....GM.....	116				
50TH.....GM.....	174				
75TH.....GM.....	232				
90TH.....GM.....	348				
95TH.....GM.....	464				
99TH.....GM.....	721				
MAXIMUM.....GM.....	2,227	0	0	232	232
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	5.1	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
.3	.4	1.0	1.4	6.7	8.2	8.3	7.1	4.7	3.7	3.4	3.6
.3	.4	.8	1.3	5.0	5.9	4.6	4.2	2.1	2.5	.8	2.2
0	0	0	.1	1.1	1.6	2.3	1.5	1.3	.8	1.4	.6
0	0	.1	0	.6	.7	1.3	1.4	1.2	.4	1.2	.7
1.0	1.0	1.4	1.2	1.6	1.6	1.8	1.8	2.1	1.5	2.7	1.6
.0	.0	1.0	.5	1.1	1.0	1.1	1.2	1.3	.9	1.6	.8
1.0	1.0	4.0	3.0	8.0	9.0	7.0	8.0	5.0	5.0	6.0	3.0
18	32	79	62	145	109	126	97	108	76	249	61
14	19	94	55	157	115	125	86	94	62	268	34
				39	20	20	19	10	20		
				58	40	45	39	39	30		
				97	77	77	77	69	45		
				155	126	155	116	135	116		
				309	232	271	222	290	199		
				348	290	387	242	329	232		
				890	453	541	464	329	232		
55	96	155	183	330	237	209	173	139	177	268	129
41	57	206	170	339	196	173	132	79	148	53	48
0	0	0	116	308	219	260	171	188	124	162	136
0	0	0	0	292	111	222	110	110	61	66	20
0	0	249	0	328	280	244	197	194	183	529	89
0	0	0	0	218	262	137	102	109	68	291	48
116	203	928	812	2,036	3,213	1,856	1,044	522	696	812	232
116	203	928	812	3,518	4,634	2,784	1,856	986	696	2,436	522
55	96	167	156	279	211	209	163	158	150	276	116
41	57	178	104	247	162	165	103	110	107	158	55
				87	60	58	58	15	45		
				145	116	116	116	116	87		
				232	174	174	120	116	120		
				348	232	232	232	200	180		
				464	348	360	240	348	232		
				696	464	464	360	406	232		
				1,567	721	870	522	464	696		
116	203	928	464	1,856	2,227	1,856	928	464	696	480	232
0	0	0	0	0	5.1	0	0	0	0	0	0
0	0	0	0	0	.8	0	0	0	0	0	0

TABLE 1.11.--DISTILLED LIQUORS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.4	0	0	0	0
ON ONLY 1 OF 3 DAYS.....PCT.....	2.1	0	0	0	0
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	0	0	0
ON ALL 3 OF 3 DAYS.....PCT.....	.6	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.0	0	0	0	0
STANDARD DEVIATION.....DO.....	1.7	0	0	0	0
MAXIMUM.....DO.....	17.0	0	0	0	0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	46	0	0	0	0
STANDARD DEVIATION.....GM.....	49	0	0	0	0
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9				
25TH.....GM.....	14				
50TH.....GM.....	28				
75TH.....GM.....	56				
90TH.....GM.....	98				
95TH.....GM.....	140				
99TH.....GM.....	252				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	90	0	0	0	0
STANDARD DEVIATION.....GM.....	102	0	0	0	0
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	92	0	0	0	0
STANDARD DEVIATION.....GM.....	80	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	80	0	0	0	0
STANDARD DEVIATION.....GM.....	60	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,232	0	0	0	0
MAXIMUM OVER 3 DAYS.....GM.....	1,316	0	0	0	0
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	69	0	0	0	0
STANDARD DEVIATION.....GM.....	68	0	0	0	0
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28				
25TH.....GM.....	28				
50TH.....GM.....	56				
75TH.....GM.....	84				
90TH.....GM.....	126				
95TH.....GM.....	168				
99TH.....GM.....	364				
MAXIMUM.....GM.....	896	0	0	0	0
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	16.6	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
(2)	0	.3	.4	5.4	3.7	7.4	5.1	5.8	3.3	3.1	1.5
(2)	0	.3	.4	3.8	2.5	4.3	3.0	1.8	2.1	.3	.6
0	0	0	.1	1.1	.8	1.8	1.1	1.0	.6	.9	.6
0	0	0	0	.5	.4	1.2	1.0	3.0	.6	1.9	.3
1.0	0	1.0	1.1	1.9	1.8	2.0	2.0	2.8	1.8	3.2	1.8
.0	0	.0	.3	2.2	1.4	1.8	1.5	1.6	1.1	2.2	.7
1.0	0	1.0	2.0	17.0	9.0	15.0	11.0	6.0	5.0	9.0	3.0
149	0	9	9	44	40	55	41	60	26	48	32
0	0	0	4	46	48	56	44	47	15	47	28
				9	9	9	9	9	5		
				14	14	19	14	25	14		
				28	23	37	28	47	25		
				56	56	70	47	93	37		
				103	75	112	89	149	42		
				121	135	168	112	149	42		
				224	299	299	280	159	75		
448	0	28	24	89	81	116	81	55	58	19	32
0	0	0	6	107	111	121	74	27	34	0	20
0	0	0	28	127	83	107	76	67	55	41	43
0	0	0	0	82	60	87	84	36	16	10	18
0	0	0	0	73	94	92	71	90	40	63	84
0	0	0	0	34	75	77	52	45	14	54	0
448	0	28	28	896	896	1,232	448	224	126	168	84
448	0	28	56	896	980	1,316	896	476	224	504	252
448	0	28	25	71	67	83	62	64	44	44	54
0	0	0	6	68	73	81	59	33	23	22	27
				28	28	28	28	28	14		
				28	42	42	28	42	28		
				56	56	56	42	56	42		
				84	56	112	56	84	56		
				168	112	140	112	112	56		
				168	168	224	168	112	112		
				252	448	448	448	168	112		
448	0	28	28	896	896	896	448	224	112	112	84
0	0	0	0	2.3	3.3	10.0	0	0	.9	0	0
0	0	0	0	.6	.9	1.4	0	0	.9	0	0

TABLE 2.01.--EGGS

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	54.3	17.7	61.3	55.2	48.5
ON ONLY 1 OF 3 DAYS.....PCT.....	28.6	8.9	31.2	32.4	32.8
ON ONLY 2 OF 3 DAYS.....PCT.....	15.5	5.7	19.7	15.7	11.8
ON ALL 3 OF 3 DAYS.....PCT.....	10.3	3.1	10.4	7.1	4.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.7	1.7	1.7	1.6	1.4
STANDARD DEVIATION.....DO.....	.9	.8	.8	.8	.7
MAXIMUM.....DO.....	9.0	4.0	4.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	47	28	34	35	34
STANDARD DEVIATION.....GM.....	33	20	22	24	23
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15	5	11	13	15
25TH.....GM.....	21	11	17	17	17
50TH.....GM.....	40	23	29	29	29
75TH.....GM.....	61	43	43	43	43
90TH.....GM.....	92	50	64	64	64
95TH.....GM.....	109	73	73	82	79
99TH.....GM.....	165	81	116	128	110
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	85	47	62	67	72
STANDARD DEVIATION.....GM.....	50	35	33	34	42
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	86	55	59	69	73
STANDARD DEVIATION.....GM.....	41	26	22	32	28
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	83	46	61	69	68
STANDARD DEVIATION.....GM.....	38	19	25	28	32
MAXIMUM IN ANY 1 DAY.....GM.....	728	220	248	440	440
MAXIMUM OVER 3 DAYS.....GM.....	1,144	284	448	526	526
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	82	49	59	66	70
STANDARD DEVIATION.....GM.....	44	30	27	34	37
PERCENTILES: <u>1/</u>					
5TH.....GM.....	40	14	25	35	40
25TH.....GM.....	50	32	46	46	46
50TH.....GM.....	64	50	52	64	64
75TH.....GM.....	100	64	64	64	88
90TH.....GM.....	128	64	92	128	128
95TH.....GM.....	150	99	128	128	128
99TH.....GM.....	237	220	141	192	220
MAXIMUM.....GM.....	715	220	220	440	440
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	88.1	.9	1.6	4.4	2.3
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	.6	.1	.3	.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
49.1	44.3	52.3	44.4	54.8	51.1	62.0	56.7	66.7	57.4	71.7	57.4
31.2	28.2	27.0	27.5	27.3	29.1	27.1	29.5	25.6	28.9	21.3	29.5
12.6	11.1	15.6	13.0	16.6	15.4	18.0	16.0	19.2	16.3	21.6	14.2
5.4	5.0	9.6	3.9	10.9	6.6	16.9	11.3	21.9	12.2	28.8	13.7
1.5	1.5	1.7	1.5	1.8	1.6	1.9	1.8	2.0	1.8	2.2	1.8
.7	.8	.8	.7	.9	.8	.9	.9	.9	.9	1.0	.9
5.0	4.0	6.0	5.0	5.0	6.0	8.0	6.0	5.0	6.0	9.0	5.0
43	38	58	40	62	45	59	43	54	38	54	37
30	27	39	27	40	31	37	30	34	25	36	22
15	13	15	13	15	15	15	15	15	15	15	13
21	17	31	19	31	21	31	21	31	17	31	17
33	31	46	33	52	38	50	33	46	33	46	33
56	46	76	49	85	59	83	56	73	49	69	48
80	73	100	75	115	85	104	85	100	70	97	64
100	92	128	87	135	100	128	100	123	85	123	85
160	132	197	128	192	144	180	141	140	128	166	109
89	77	108	82	112	88	100	77	88	68	69	69
49	45	60	49	61	51	56	41	41	37	29	31
86	76	102	81	109	85	100	80	87	65	84	67
42	36	41	35	47	36	46	40	39	26	44	25
86	79	104	81	107	85	93	75	80	66	76	60
38	35	45	35	43	39	37	31	35	26	35	22
572	406	566	440	534	715	728	660	400	287	457	256
704	605	768	584	998	826	1,144	880	800	534	828	384
85	75	101	79	105	83	93	74	81	64	73	63
47	40	49	41	52	45	46	39	39	30	36	28
40	40	46	40	46	40	44	40	40	40	40	40
50	46	64	46	80	50	50	46	50	46	46	46
80	64	92	64	92	64	92	64	80	50	64	50
100	92	128	100	128	104	128	92	100	64	92	64
128	128	150	128	165	128	128	128	128	110	128	100
165	128	192	128	192	138	160	128	136	128	128	128
240	220	256	220	316	234	256	220	200	134	192	128
480	360	486	440	512	715	486	480	330	256	365	200
4.9	4.7	1.5	5.0	2.0	16.5	15.3	16.6	5.1	4.5	1.5	1.3
.3	.3	.1	.5	.1	.4	.3	.2	.3	.3	.2	.1

TABLE 2.02.--BOILED AND POACHED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	12.4	3.7	11.8	8.9	7.1
ON ONLY 1 OF 3 DAYS.....PCT.....	8.9	2.1	8.8	7.2	6.0
ON ONLY 2 OF 3 DAYS.....PCT.....	2.2	1.1	1.9	1.3	1.0
ON ALL 3 OF 3 DAYS.....PCT.....	1.3	.5	1.1	.4	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.6	1.4	1.3	1.2
STANDARD DEVIATION.....DO.....	.7	.7	.6	.5	.4
MAXIMUM.....DO.....	7.0	3.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	31	21	22	23	22
STANDARD DEVIATION.....GM.....	21	11	12	13	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		8	8	15
25TH.....GM.....	17		15	17	17
50TH.....GM.....	29		17	17	17
75TH.....GM.....	33		29	29	29
90TH.....GM.....	50		39	33	33
95TH.....GM.....	67		50	50	50
99TH.....GM.....	100		56	83	67
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	67	40	49	58	55
STANDARD DEVIATION.....GM.....	30	15	19	24	18
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	67	48	52	58	61
STANDARD DEVIATION.....GM.....	27	3	12	22	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	65	32	43	46	48
STANDARD DEVIATION.....GM.....	30	11	11	20	2
MAXIMUM IN ANY 1 DAY.....GM.....	400	50	132	150	100
MAXIMUM OVER 3 DAYS.....GM.....	850	150	167	261	200
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	65	41	48	55	56
STANDARD DEVIATION.....GM.....	28	13	16	22	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	44		25	25	44
25TH.....GM.....	50		44	44	50
50TH.....GM.....	50		50	50	50
75TH.....GM.....	88		50	50	50
90TH.....GM.....	100		58	100	100
95TH.....GM.....	100		88	100	100
99TH.....GM.....	150		100	116	100
MAXIMUM.....GM.....	250	50	100	150	100
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	29.6	0	0	1.0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	.5	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.9	7.5	4.0	7.4	7.8	12.5	13.1	17.7	19.3	22.6	22.2	24.2
5.0	6.2	3.4	5.8	6.1	9.2	9.4	12.6	10.7	15.2	12.9	13.8
.7	.8	.5	1.2	1.2	2.4	2.3	3.2	3.9	4.9	4.3	5.7
.1	.6	.1	.4	.5	.9	1.5	1.9	4.7	2.5	5.0	4.7
1.2	1.3	1.2	1.3	1.3	1.4	1.5	1.4	1.8	1.5	1.7	1.6
.4	.6	.6	.6	.6	.7	.8	.7	1.0	.7	.9	.9
3.0	3.0	4.0	3.0	4.0	6.0	7.0	6.0	5.0	4.0	4.0	5.0
26	26	34	29	37	31	35	29	42	27	36	32
15	17	24	18	25	20	27	17	34	17	18	19
11	15	8	13	8	15	15	15	15	13	15	15
17	17	17	17	17	17	17	17	17	17	17	17
17	17	33	25	33	29	33	25	33	17	33	29
33	33	33	33	45	33	39	33	50	33	50	44
42	50	59	50	67	51	67	50	83	50	59	50
59	58	67	67	96	67	88	67	100	59	67	72
67	100	133	100	117	100	117	100	133	92	100	100
68	63	86	67	82	69	75	64	70	57	73	58
30	29	54	26	42	31	35	25	30	24	28	23
67	56	84	79	92	68	76	62	72	61	78	59
20	15	46	34	37	25	31	23	29	22	22	20
77	66	85	58	86	71	78	59	80	54	53	58
25	21	17	17	22	26	45	17	44	14	12	18
200	174	400	200	232	250	350	232	400	200	150	116
300	300	400	350	400	500	850	348	800	300	300	300
68	62	82	67	84	68	73	61	72	56	64	57
28	27	39	27	39	29	33	23	32	23	23	21
38	38	33	38	25	44	44	44	44	44	44	44
50	50	50	50	50	50	50	50	50	50	50	50
50	50	100	50	88	50	50	50	50	50	50	50
100	88	100	100	100	100	100	76	100	50	100	50
100	100	150	100	132	100	100	100	100	100	100	100
100	100	150	100	150	100	132	100	116	100	100	100
150	174	200	150	200	152	174	116	200	116	116	116
200	174	200	150	232	200	250	184	200	200	116	116
.7	2.9	.9	1.0	1.0	6.0	5.7	7.7	0	0	1.5	1.3
.5	1.4	1.3	.7	.3	.7	.6	.4	0	0	.7	.3

TABLE 2.03.--FRIED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	24.3	2.1	20.7	21.7	20.8
ON ONLY 1 OF 3 DAYS.....PCT.....	15.9	1.8	13.5	15.3	16.1
ON ONLY 2 OF 3 DAYS.....PCT.....	5.5	0	5.9	5.2	3.5
ON ALL 3 OF 3 DAYS.....PCT.....	3.0	.3	1.3	1.2	1.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.3	1.4	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.7	.6	.6	.6
MAXIMUM.....DO.....	9.0	3.0	4.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	34	22	22	23	24
STANDARD DEVIATION.....GM.....	23	12	12	14	14
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	13		12	13	13
25TH.....GM.....	15		15	15	15
50TH.....GM.....	31		15	15	15
75TH.....GM.....	44		31	31	31
90TH.....GM.....	61		33	40	43
95TH.....GM.....	92		46	46	61
99TH.....GM.....	107		61	77	80
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	69	54	47	52	55
STANDARD DEVIATION.....GM.....	29	25	15	19	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	0	45	51	59
STANDARD DEVIATION.....GM.....	27	0	11	16	20
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	46	50	53	57
STANDARD DEVIATION.....GM.....	28	0	16	23	18
MAXIMUM IN ANY 1 DAY.....GM.....	368	92	104	184	138
MAXIMUM OVER 3 DAYS.....GM.....	828	138	312	368	276
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	69	51	46	51	56
STANDARD DEVIATION.....GM.....	28	21	15	18	21
PERCENTILES: <u>1/</u>					
5TH.....GM.....	40		23	40	40
25TH.....GM.....	46		40	40	46
50TH.....GM.....	52		46	46	46
75TH.....GM.....	92		46	46	80
90TH.....GM.....	92		52	92	92
95TH.....GM.....	104		92	92	92
99TH.....GM.....	138		104	104	104
MAXIMUM.....GM.....	368	92	104	138	138
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	14.8	0	1.6	0	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	.5	0	.3

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
24.3	19.9	27.7	19.0	28.9	19.8	33.7	24.1	31.2	19.9	36.4	17.8
17.9	15.2	17.1	14.3	17.3	14.7	19.3	16.2	15.3	13.0	14.7	10.5
4.5	3.5	7.2	3.8	7.8	3.5	8.7	5.2	8.5	3.6	9.2	3.7
1.9	1.2	3.4	.9	3.8	1.5	5.7	2.7	7.4	3.4	12.5	3.6
1.4	1.3	1.5	1.3	1.5	1.3	1.6	1.5	1.8	1.5	2.0	1.6
.7	.6	.7	.6	.7	.6	.8	.7	.9	.8	1.1	.8
5.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	9.0	3.0
32	26	43	27	45	29	43	30	40	28	44	30
20	17	26	15	27	17	25	18	25	16	32	17
13	13	15	13	15	13	15	13	15	13	13	13
15	15	31	15	31	15	27	15	15	15	27	15
31	15	31	27	31	31	31	29	31	23	31	31
31	31	53	31	61	31	61	31	52	32	61	40
61	46	88	46	88	53	92	57	77	46	92	52
77	61	92	61	92	61	92	69	92	61	92	61
100	92	123	88	128	92	132	92	123	82	123	92
73	58	84	62	89	67	81	62	68	56	66	58
35	21	27	23	32	26	29	24	27	20	27	25
70	63	84	65	88	66	81	63	73	55	62	55
27	27	27	25	27	23	27	23	26	15	24	18
73	59	88	56	89	64	78	61	66	53	73	52
24	23	31	18	30	23	26	21	26	17	36	16
368	323	230	172	276	230	276	184	200	184	276	184
412	415	690	276	644	322	450	385	400	368	828	276
71	59	85	62	88	66	79	61	68	54	65	55
32	24	29	23	31	25	27	23	26	18	25	21
40	40	46	40	46	40	40	40	40	40	40	40
46	46	80	46	80	46	46	46	46	46	46	40
80	46	92	46	92	46	92	46	52	46	46	46
92	80	92	92	92	92	92	92	92	52	92	52
92	92	104	92	138	92	104	92	92	92	92	92
104	92	138	92	138	92	138	92	92	92	92	92
160	104	184	120	184	138	156	120	138	96	138	92
368	323	230	138	276	184	184	140	146	104	138	184
1.8	0	0	0	0	3.2	3.5	1.0	2.1	0	0	0
.3	0	0	0	0	.2	.1	(2)	.3	0	0	0

TABLE 2.04.--SCRAMBLED EGGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	20.1	8.6	32.6	26.8	20.5
ON ONLY 1 OF 3 DAYS.....PCT.....	14.2	5.1	20.6	19.0	15.9
ON ONLY 2 OF 3 DAYS.....PCT.....	4.1	3.3	8.7	5.1	3.6
ON ALL 3 OF 3 DAYS.....PCT.....	1.8	.3	3.3	2.7	1.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.5	1.5	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.6	.7	.7	.6
MAXIMUM.....DO.....	4.0	3.0	4.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	49	31	36	37	38
STANDARD DEVIATION.....GM.....	32	19	22	24	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	21		11	19	21
25TH.....GM.....	21		21	21	21
50TH.....GM.....	43		21	21	32
75TH.....GM.....	64		43	43	43
90TH.....GM.....	85		64	64	64
95TH.....GM.....	128		73	85	85
99TH.....GM.....	157		128	128	147
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	109	59	75	81	92
STANDARD DEVIATION.....GM.....	52	41	34	37	49
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	104	70	71	79	79
STANDARD DEVIATION.....GM.....	46	20	28	28	23
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	102	53	73	81	92
STANDARD DEVIATION.....GM.....	40	0	24	29	42
MAXIMUM IN ANY 1 DAY.....GM.....	728	220	220	256	440
MAXIMUM OVER 3 DAYS.....GM.....	1,144	284	448	512	512
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	105	63	72	80	88
STANDARD DEVIATION.....GM.....	48	33	29	34	43
PERCENTILES: <u>1/</u>					
5TH.....GM.....	64		32	55	64
25TH.....GM.....	64		64	64	64
50TH.....GM.....	128		64	64	64
75TH.....GM.....	128		64	96	128
90TH.....GM.....	128		128	128	128
95TH.....GM.....	192		128	128	165
99TH.....GM.....	256		165	220	220
MAXIMUM.....GM.....	715	220	220	220	440
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	7.2	0	0	1.9	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	0	.3	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.1	16.7	21.5	17.8	20.7	20.6	20.5	18.8	19.9	18.5	18.3	17.7
14.3	12.4	15.3	13.7	14.5	15.0	13.5	13.5	12.2	12.8	9.3	13.5
3.5	3.3	4.0	3.1	4.3	4.3	4.6	3.7	4.1	3.4	4.8	2.4
1.3	1.0	2.2	1.0	1.9	1.3	2.3	1.6	3.6	2.2	4.3	1.8
1.3	1.3	1.4	1.3	1.4	1.3	1.5	1.4	1.6	1.4	1.7	1.3
.6	.6	.7	.6	.7	.6	.7	.7	.8	.7	.8	.7
4.0	3.0	3.0	3.0	3.0	4.0	4.0	4.0	3.0	4.0	3.0	3.0
48	42	60	45	63	48	60	45	58	42	59	39
29	26	37	28	37	30	35	28	33	25	41	22
21	21	21	21	21	21	21	21	21	21	21	21
21	21	43	21	43	21	43	21	43	21	21	21
43	43	43	43	43	43	43	43	43	43	43	37
64	43	73	43	85	61	85	43	79	43	85	43
85	73	107	85	128	85	107	85	128	85	128	64
107	85	147	110	147	90	128	97	128	85	128	85
165	128	192	147	192	171	192	128	149	128	192	128
115	96	133	105	141	110	132	103	116	90	91	89
56	45	58	52	58	50	53	42	39	38	35	34
101	94	139	103	136	103	123	98	118	86	96	87
41	34	51	45	54	36	52	43	36	27	37	26
101	95	111	103	125	112	114	93	101	85	113	81
44	37	49	28	40	46	36	36	37	27	38	24
568	348	440	440	512	715	728	660	256	256	198	256
704	605	768	512	714	715	1,144	880	576	384	576	384
108	95	129	104	135	107	124	99	111	87	102	86
51	39	55	47	54	48	47	41	38	34	39	31
64	64	64	64	64	64	64	64	64	64	64	64
64	64	64	64	128	64	128	64	64	64	64	64
128	64	128	128	128	128	128	85	128	64	128	64
128	128	128	128	128	128	128	128	128	128	128	128
165	128	192	128	192	128	192	128	136	128	128	128
220	165	220	192	256	192	192	158	192	128	192	128
256	220	320	256	320	256	256	220	192	192	198	158
440	256	440	440	512	715	468	440	220	256	198	158
1.1	0	0	1.1	0	2.0	1.0	0	0	0	0	0
.2	0	0	.3	0	.1	.1	0	0	0	0	0

TABLE 3.01.--BUTTER

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	31.4	5.2	29.2	28.7	31.7
ON ONLY 1 OF 3 DAYS.....PCT.....	16.7	3.8	17.4	15.9	17.2
ON ONLY 2 OF 3 DAYS.....PCT.....	8.4	1.0	7.2	7.7	8.6
ON ALL 3 OF 3 DAYS.....PCT.....	6.3	.3	4.6	5.1	5.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.1	1.5	1.9	2.0	2.0
STANDARD DEVIATION.....DO.....	1.4	.9	1.2	1.3	1.3
MAXIMUM.....DO.....	12.0	4.0	7.0	9.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	8	3	4	6	6
STANDARD DEVIATION.....GM.....	10	3	4	7	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	2		2	2	2
50TH.....GM.....	5		3	4	5
75TH.....GM.....	9		5	7	7
90TH.....GM.....	19		9	13	13
95TH.....GM.....	26		14	19	19
99TH.....GM.....	47		19	29	38
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	13	5	8	10	11
STANDARD DEVIATION.....GM.....	13	4	8	8	12
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	14	6	9	11	11
STANDARD DEVIATION.....GM.....	14	1	7	9	12
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	17	13	8	13	12
STANDARD DEVIATION.....GM.....	15	4	4	12	10
MAXIMUM IN ANY 1 DAY.....GM.....	341	28	57	128	227
MAXIMUM OVER 3 DAYS.....GM.....	454	50	57	298	236
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	12	6	7	9	10
STANDARD DEVIATION.....GM.....	13	4	6	10	11
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	5		5	5	5
50TH.....GM.....	7		5	7	5
75TH.....GM.....	14		7	10	10
90TH.....GM.....	28		14	14	14
95TH.....GM.....	28		14	28	28
99TH.....GM.....	57		28	43	57
MAXIMUM.....GM.....	341	14	57	128	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2890.4	4.2	89.4	157.6	186.1
PROPORTION OF EATING OCCASIONS.....PCT.....	11.6	10.7	15.2	16.0	16.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
32.4	30.9	32.4	32.6	34.1	30.8	35.6	31.4	30.0	27.2	29.9	29.4
17.2	18.3	18.3	20.2	18.1	16.7	18.1	16.8	13.5	12.3	9.1	12.4
9.3	8.3	8.1	8.4	9.9	9.1	8.8	8.3	7.9	7.4	7.0	6.5
5.9	4.2	6.0	3.5	6.2	5.1	8.8	6.3	8.6	7.5	13.9	10.5
2.0	1.8	1.9	1.7	2.0	2.0	2.2	2.1	2.5	2.3	3.2	2.8
1.3	1.2	1.3	1.1	1.4	1.2	1.6	1.4	1.7	1.5	2.2	2.0
9.0	8.0	7.0	8.0	9.0	8.0	9.0	9.0	9.0	8.0	12.0	9.0
8	6	9	7	10	8	10	7	9	7	13	10
10	6	11	11	11	10	13	8	11	8	14	11
2	2	2	2	2	2	2	2	2	2	2	2
2	2	3	2	3	2	3	2	3	3	3	2
5	4	5	5	6	5	6	5	6	5	8	6
9	8	10	9	12	9	13	9	11	9	17	13
19	13	19	14	20	19	24	16	20	14	28	21
24	19	26	19	29	27	32	24	30	20	40	35
52	28	52	80	44	47	59	38	52	36	69	66
13	11	14	12	15	13	15	12	13	13	11	12
14	10	15	11	14	15	15	13	12	12	14	15
15	11	16	15	18	15	17	13	15	11	15	14
17	7	13	15	18	14	16	11	12	7	14	9
16	13	19	19	19	18	21	15	17	12	20	17
15	9	16	23	14	16	19	11	16	11	15	14
298	123	200	170	227	341	227	184	142	199	85	113
313	170	370	288	397	364	454	213	312	220	208	203
12	10	14	13	14	13	14	11	11	9	12	10
15	9	12	14	13	15	14	10	11	11	9	10
4	2	5	5	5	2	5	2	2	2	2	2
5	5	5	5	5	5	5	5	5	5	5	5
9	7	9	9	10	7	9	7	9	5	9	7
14	14	14	14	14	14	14	14	14	10	14	14
28	15	28	28	28	28	28	27	28	14	28	15
28	28	42	43	43	28	43	28	28	28	28	28
71	43	57	57	57	57	71	57	57	43	43	57
298	114	114	114	170	341	199	160	114	199	43	113
147.5	171.5	127.9	127.9	297.1	379.3	473.9	466.8	65.1	105.9	28.7	61.5
10.8	14.0	14.5	15.9	11.0	11.7	12.0	10.0	7.8	9.6	5.6	7.6

TABLE 3.02.--MARGARINE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	43.1	8.5	43.8	46.1	42.9
ON ONLY 1 OF 3 DAYS.....PCT.....	18.7	4.9	21.0	21.9	18.9
ON ONLY 2 OF 3 DAYS.....PCT.....	13.2	2.4	14.0	14.2	15.2
ON ALL 3 OF 3 DAYS.....PCT.....	11.2	1.2	8.7	9.9	8.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	2.3	1.8	2.1	2.2	2.2
STANDARD DEVIATION.....DO.....	1.5	1.1	1.3	1.3	1.3
MAXIMUM.....DO.....	12.0	5.0	7.0	8.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	9	3	5	6	7
STANDARD DEVIATION.....GM.....	10	2	6	6	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	1	2
25TH.....GM.....	3		2	2	2
50TH.....GM.....	5		3	5	5
75TH.....GM.....	10		6	7	9
90TH.....GM.....	19		9	12	15
95TH.....GM.....	28		13	16	20
99TH.....GM.....	47		19	31	30
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	13	6	7	9	11
STANDARD DEVIATION.....GM.....	14	6	6	8	13
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	13	5	8	10	11
STANDARD DEVIATION.....GM.....	11	3	7	8	8
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	16	5	9	10	12
STANDARD DEVIATION.....GM.....	14	2	10	7	9
MAXIMUM IN ANY 1 DAY.....GM.....	227	27	113	114	113
MAXIMUM OVER 3 DAYS.....GM.....	383	28	269	119	184
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	11	5	6	8	9
STANDARD DEVIATION.....GM.....	11	4	6	8	8
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	5		3	5	5
50TH.....GM.....	7		5	5	5
75TH.....GM.....	14		7	9	14
90TH.....GM.....	28		14	14	14
95TH.....GM.....	28		14	21	28
99TH.....GM.....	57		28	28	43
MAXIMUM.....GM.....	227	27	71	114	113
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	1598.5	5.5	50.7	73.3	90.2
PROPORTION OF EATING OCCASIONS.....PCT.....	4.2	7.3	5.2	4.3	5.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
44.8	40.7	41.4	38.6	39.8	39.8	44.0	44.1	54.0	53.2	48.1	51.7
20.8	19.6	19.8	18.3	20.2	19.2	17.3	18.0	16.9	19.1	14.6	16.2
13.9	13.5	12.8	12.8	11.7	12.5	11.9	13.9	12.8	17.0	12.9	15.2
10.0	7.6	8.8	7.5	7.9	8.1	14.8	12.2	24.3	17.1	20.6	20.3
2.2	2.1	2.2	2.1	2.1	2.1	2.5	2.4	3.0	2.6	2.9	2.9
1.4	1.3	1.5	1.3	1.4	1.3	1.7	1.5	1.7	1.6	1.8	1.8
8.0	7.0	9.0	7.0	10.0	8.0	11.0	12.0	9.0	9.0	8.0	9.0
9	8	12	7	10	8	11	8	11	8	11	9
10	8	14	8	12	9	12	8	10	8	13	9
2	1	2	2	2	2	2	2	2	2	2	2
3	3	3	3	3	2	3	2	5	3	5	3
5	5	8	5	6	5	7	5	8	5	7	5
11	9	14	9	14	9	14	9	16	9	13	11
21	17	27	15	22	19	27	16	24	16	20	22
31	25	38	20	28	24	36	22	28	24	38	28
52	43	74	47	61	47	58	42	52	42	71	40
13	13	17	12	16	13	15	11	14	10	17	10
13	13	16	11	14	16	18	10	12	8	19	10
15	14	19	12	17	13	15	11	14	12	12	12
14	13	17	10	14	11	11	9	12	11	9	10
18	14	25	15	21	16	19	13	16	12	16	14
13	10	20	12	18	14	15	12	11	10	16	11
142	170	142	184	173	170	227	175	170	120	142	227
255	175	341	255	383	227	284	350	213	227	270	236
12	11	16	11	14	11	13	10	11	9	11	9
12	12	14	9	13	11	13	9	10	9	12	11
2	2	5	2	5	2	2	2	2	2	2	2
5	5	5	5	5	5	5	5	5	5	5	5
9	7	10	7	9	9	9	5	9	5	7	5
14	14	20	14	14	14	14	14	14	9	14	10
28	28	28	28	28	28	28	19	24	14	28	14
28	28	43	28	43	28	28	28	28	28	28	28
57	57	71	43	57	57	57	43	56	57	57	43
142	170	128	85	114	170	227	113	170	85	99	227
77.6	76.0	63.3	53.8	144.9	189.5	246.6	284.1	53.9	96.7	32.2	60.2
3.8	4.1	4.9	4.4	4.3	4.3	4.5	3.8	3.0	4.0	4.3	4.1

TABLE 3.03.--WHIPPED BUTTER AND MARGARINE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FCCD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.9	.5	1.8	2.0	1.7
ON ONLY 1 OF 3 DAYS.....PCT.....	1.1	.2	1.0	1.0	.9
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	0	.6	.6	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.3	.3	.2	.4	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.9	2.7	2.2	2.3	2.0
STANDARD DEVIATION.....CO.....	1.3	1.5	1.2	1.4	1.2
MAXIMUM.....CO.....	10.0	4.0	5.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	5	3	4	5	5
STANDARD DEVIATION.....GM.....	6	2	3	4	5
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1				
25TH.....GM.....	2				
50TH.....GM.....	3				
75TH.....GM.....	6				
90TH.....GM.....	13				
95TH.....GM.....	17				
99TH.....GM.....	31				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	11	2	6	7	10
STANDARD DEVIATION.....GM.....	12	0	4	5	9
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	10	0	8	8	7
STANDARD DEVIATION.....GM.....	9	0	6	8	4
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	10	4	5	9	14
STANDARD DEVIATION.....GM.....	10	0	2	4	7
MAXIMUM IN ANY 1 DAY.....GM.....	104	5	22	44	38
MAXIMUM OVER 3 DAYS.....GM.....	179	13	40	63	76
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	8	3	5	6	8
STANDARD DEVIATION.....GM.....	9	1	4	5	5
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2				
25TH.....GM.....	3				
50TH.....GM.....	6				
75TH.....GM.....	9				
90TH.....GM.....	19				
95TH.....GM.....	28				
99TH.....GM.....	38				
MAXIMUM.....GM.....	76	5	22	28	19
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	18.2	0	1.7	1.2	2.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.3	0	4.4	1.6	4.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.3	1.4	1.9	2.3	1.9	2.1	2.3	1.7	2.7	2.3	2.3	1.5
.6	1.1	1.2	1.5	1.2	1.4	1.2	.9	1.4	1.6	.6	.7
.3	.3	.4	.6	.4	.6	.8	.4	.7	.5	1.3	.5
.4	.1	.3	.2	.3	.1	.3	.3	.6	.2	.4	.3
2.3	1.4	1.8	1.5	1.9	1.7	2.1	1.9	2.3	1.6	2.6	2.2
1.4	.7	1.1	.7	1.3	1.1	1.3	1.4	1.4	1.0	1.0	1.3
5.0	4.0	6.0	3.0	6.0	6.0	6.0	10.0	5.0	4.0	4.0	4.0
7	4	5	5	6	5	7	4	8	6	9	5
10	4	5	5	5	5	6	4	13	9	11	4
				1	1	1	1				
				2	2	3	1				
				4	3	5	3				
				6	6	9	5				
				13	9	13	9				
				19	14	19	13				
				20	31	31	23				
9	9	7	13	14	11	14	10	11	11	8	5
7	6	5	15	13	12	16	9	11	16	4	4
8	13	11	10	9	10	14	6	11	23	7	13
3	10	7	8	7	10	9	5	7	17	6	4
15	5	12	6	9	8	10	7	18	4	31	6
14	2	3	2	6	3	7	6	23	1	0	1
57	38	28	57	57	76	85	38	104	57	38	19
142	66	47	57	60	94	94	69	179	113	94	38
9	9	8	10	9	8	10	7	10	12	10	7
10	7	5	12	9	9	8	7	12	14	10	5
				3	2	2	2				
				3	3	3	3				
				6	6	8	3				
				9	9	16	9				
				19	19	19	19				
				28	19	19	28				
				57	38	47	28				
57	28	19	57	57	76	47	38	38	57	38	19
0	0	0	0	0	4.2	2.0	4.2	0	0	2.3	0
0	0	0	0	0	2.2	.8	1.9	0	0	7.3	0

TABLE 3.04.--SALAD DRESSINGS AND MAYONNAISE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	44.7	1.1	22.6	34.2	39.1
ON ONLY 1 OF 3 DAYS.....PCT.....	25.5	1.1	16.4	21.7	24.7
ON ONLY 2 OF 3 DAYS.....PCT.....	13.5	0	4.8	9.9	10.4
ON ALL 3 OF 3 DAYS.....PCT.....	5.7	0	1.3	2.5	4.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.7	1.0	1.5	1.6	1.6
STANDARD DEVIATION.....DO.....	1.0	.0	.8	.8	.9
MAXIMUM.....DO.....	8.0	1.0	5.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	12	3	5	8	8
STANDARD DEVIATION.....GM.....	14	2	5	10	8
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	5		2	2	3
50TH.....GM.....	9		3	5	5
75TH.....GM.....	16		7	9	10
90TH.....GM.....	26		11	16	18
95TH.....GM.....	36		15	22	21
99TH.....GM.....	59		24	43	40
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	23	8	11	16	17
STANDARD DEVIATION.....GM.....	21	5	12	18	15
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	23	0	10	14	17
STANDARD DEVIATION.....GM.....	19	0	8	12	14
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	25	0	11	20	15
STANDARD DEVIATION.....GM.....	23	0	5	24	11
MAXIMUM IN ANY 1 DAY.....GM.....	768	15	83	165	125
MAXIMUM OVER 3 DAYS.....GM.....	1,536	15	83	289	189
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	21	8	10	15	15
STANDARD DEVIATION.....GM.....	22	5	10	18	14
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		2	2	2
25TH.....GM.....	9		5	5	5
50TH.....GM.....	15		5	10	14
75TH.....GM.....	29		14	15	16
90TH.....GM.....	44		28	30	31
95TH.....GM.....	60		30	41	32
99TH.....GM.....	94		41	83	63
MAXIMUM.....GM.....	768	15	83	165	125
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	1601.6	0	20.5	46.9	67.5
PROPORTION OF EATING OCCASIONS.....PCT.....	5.4	0	5.8	5.1	5.8

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
37.6	41.9	40.3	43.5	49.5	51.7	51.7	50.3	42.3	41.5	28.0	34.9
24.1	26.0	23.9	24.4	26.9	28.3	27.0	28.3	26.3	23.7	15.4	22.0
10.3	11.9	11.0	14.1	14.6	17.7	15.5	15.7	10.9	12.5	8.6	8.2
3.2	4.0	5.4	5.0	8.1	5.8	9.1	6.3	5.1	5.3	4.0	4.7
1.6	1.6	1.7	1.7	1.9	1.7	1.9	1.8	1.6	1.7	1.7	1.7
.9	.9	1.0	1.0	1.1	.9	1.1	1.0	.9	.9	.9	.9
6.0	6.0	6.0	6.0	6.0	6.0	8.0	6.0	5.0	6.0	4.0	6.0
10	10	13	13	15	13	15	12	12	9	12	10
10	10	12	16	15	12	19	12	12	9	14	9
2	2	2	2	2	2	2	2	2	2	2	2
3	5	5	5	5	5	5	5	5	3	5	4
7	6	10	9	10	10	10	10	9	6	9	6
12	12	18	16	20	18	20	15	14	11	16	12
21	20	29	27	31	26	31	25	26	20	24	21
29	27	40	37	41	36	41	36	39	26	29	26
46	46	51	61	73	60	63	61	56	44	56	41
22	19	27	23	28	24	26	23	20	19	21	18
26	17	26	24	25	21	22	21	16	17	16	14
20	22	26	25	27	25	27	22	24	17	24	20
13	20	19	33	20	18	19	17	21	14	27	15
22	18	24	26	28	25	28	26	28	18	25	19
14	12	16	17	22	17	37	18	16	12	14	11
279	270	151	470	278	245	768	245	169	201	250	78
279	297	340	940	495	360	1,536	311	250	256	365	184
19	18	23	22	24	22	24	21	21	16	22	17
19	18	20	27	22	20	29	19	18	16	17	14
5	5	5	5	5	5	5	5	5	3	5	5
7	5	11	9	11	9	10	9	10	5	14	5
14	14	15	15	16	15	16	15	15	14	15	15
28	28	29	30	31	30	31	29	30	21	29	28
32	32	46	44	55	46	47	44	32	31	46	32
59	49	63	60	63	61	63	60	59	41	59	47
83	83	118	94	118	94	118	83	113	64	83	63
250	256	139	470	250	245	768	245	125	188	125	78
75.0	76.2	58.4	78.5	226.0	261.5	282.7	286.9	43.0	42.2	7.4	29.2
6.1	5.2	6.0	7.1	6.3	5.4	5.9	4.6	5.6	3.4	2.9	5.1

TABLE 3.05.--MAYONNAISE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	21.4	.4	13.3	19.4	20.3
ON ONLY 1 OF 3 DAYS.....PCT.....	14.4	.4	10.0	13.7	13.9
ON ONLY 2 OF 3 DAYS.....PCT.....	5.3	0	2.8	4.2	4.7
ON ALL 3 OF 3 DAYS.....PCT.....	1.7	0	.6	1.5	1.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.0	1.4	1.4	1.5
STANDARD DEVIATION.....DO.....	.8	.0	.7	.7	.8
MAXIMUM.....DO.....	8.0	1.0	5.0	4.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	7	3	4	6	5
STANDARD DEVIATION.....GM.....	7	2	3	10	5
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	1	1
25TH.....GM.....	2		2	2	2
50TH.....GM.....	5		2	5	5
75TH.....GM.....	9		5	6	9
90TH.....GM.....	14		9	9	11
95TH.....GM.....	18		9	14	14
99TH.....GM.....	31		14	96	28
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	14	10	9	12	11
STANDARD DEVIATION.....GM.....	13	5	8	10	8
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	14	0	8	10	12
STANDARD DEVIATION.....GM.....	10	0	5	8	8
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	15	0	7	24	12
STANDARD DEVIATION.....GM.....	11	0	2	30	10
MAXIMUM IN ANY 1 DAY.....GM.....	206	14	41	165	69
MAXIMUM OVER 3 DAYS.....GM.....	289	14	41	289	165
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	13	10	8	13	11
STANDARD DEVIATION.....GM.....	12	5	7	17	8
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		2	2	2
25TH.....GM.....	5		5	5	5
50TH.....GM.....	14		5	9	9
75TH.....GM.....	14		14	14	14
90TH.....GM.....	28		14	28	28
95TH.....GM.....	28		28	28	28
99TH.....GM.....	55		28	83	41
MAXIMUM.....GM.....	193	14	41	165	55
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	774.6	0	9.5	28.7	42.9
PROPORTION OF EATING OCCASIONS.....PCT.....	6.4	0	5.0	6.0	7.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.7	20.6	21.0	21.9	25.6	25.4	24.1	22.2	15.9	18.4	12.6	13.3
12.7	13.7	13.4	14.7	16.1	17.6	15.1	15.4	11.7	11.8	7.8	10.0
3.9	5.4	5.1	5.7	6.8	6.4	6.2	5.4	2.8	5.4	3.2	2.4
1.0	1.5	2.5	1.4	2.7	1.4	2.9	1.3	1.4	1.3	1.5	.8
1.4	1.5	1.6	1.5	1.6	1.4	1.6	1.4	1.4	1.5	1.5	1.4
.8	.7	.9	.8	.9	.7	.9	.7	.7	.8	.7	.7
6.0	4.0	6.0	6.0	6.0	5.0	8.0	5.0	4.0	5.0	3.0	4.0
6	6	9	6	9	6	7	5	7	6	8	5
7	6	9	6	8	5	7	5	8	5	5	4
1	2	2	2	2	2	2	1	2	1	2	1
2	2	5	2	5	2	3	2	2	2	5	2
5	5	6	5	6	5	5	5	5	5	9	5
9	7	10	9	12	9	9	8	9	9	9	6
14	14	18	14	18	14	14	11	18	12	12	14
18	18	28	18	23	18	23	14	21	18	14	14
38	28	46	28	37	25	37	28	53	25	28	18
14	13	18	13	19	14	16	12	14	12	16	11
14	12	18	11	17	12	13	11	12	11	9	10
15	13	16	13	19	14	15	12	16	11	14	12
12	10	9	8	13	9	11	10	7	9	2	7
16	12	21	15	16	14	14	12	22	14	15	12
11	9	12	12	8	7	9	7	15	7	8	5
113	110	151	110	110	124	110	206	73	62	41	55
138	138	154	142	220	138	124	234	159	83	83	55
14	12	17	12	16	13	14	11	15	11	15	11
12	11	13	10	14	10	11	10	13	10	8	8
2	2	5	2	5	2	2	2	5	2	5	2
5	5	9	5	5	5	5	5	5	5	14	5
14	14	14	14	14	14	14	9	14	9	14	14
14	14	28	14	28	14	14	14	14	14	14	14
28	28	28	28	28	28	28	28	28	28	28	28
28	28	41	28	41	28	28	28	41	28	28	28
55	55	69	41	69	55	55	41	73	55	41	41
113	110	110	110	110	84	110	193	73	62	41	41
30.1	27.8	38.1	41.7	132.6	132.6	120.5	121.4	18.9	19.2	1.5	8.9
5.7	4.3	8.2	8.6	8.2	6.7	6.4	5.4	7.4	4.0	1.4	5.0

TABLE 3.06.--SALAD DRESSINGS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST CNCE IN 3 DAYS.....PCT.....	29.7	.6	11.0	18.6	22.8
CN ONLY 1 OF 3 DAYS.....PCT.....	19.2	.6	8.5	13.4	15.8
ON ONLY 2 OF 3 DAYS.....PCT.....	7.8	0	2.3	4.6	5.4
ON ALL 3 OF 3 DAYS.....PCT.....	2.7	0	.3	.6	1.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.4	1.4	1.5
STANDARD DEVIATION.....DO.....	.8	.0	.7	.6	.8
MAXIMUM.....DO.....	6.0	1.0	5.0	4.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	14	2	6	8	9
STANDARD DEVIATION.....GM.....	14	2	6	8	8
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		(2)	2	2
25TH.....GM.....	5		2	2	3
50TH.....GM.....	10		4	5	7
75TH.....GM.....	20		7	10	11
90TH.....GM.....	29		15	19	20
95TH.....GM.....	39		20	21	21
99TH.....GM.....	62		28	42	41
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	28	7	13	18	21
STANDARD DEVIATION.....GM.....	23	5	14	21	17
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	0	14	18	20
STANDARD DEVIATION.....GM.....	29	0	11	14	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	28	0	11	10	16
STANDARD DEVIATION.....GM.....	19	0	5	6	12
MAXIMUM IN ANY 1 DAY.....GM.....	768	15	83	128	125
MAXIMUM OVER 3 DAYS.....GM.....	1,518	15	83	128	189
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	26	7	12	17	19
STANDARD DEVIATION.....GM.....	25	5	12	18	16
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		2	3	5
25TH.....GM.....	15		5	5	5
50TH.....GM.....	16		7	15	15
75TH.....GM.....	31		16	16	29
90TH.....GM.....	59		30	31	32
95TH.....GM.....	63		32	59	59
99TH.....GM.....	123		59	123	64
MAXIMUM.....GM.....	768	15	83	128	125
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	829.6	0	11.0	18.1	24.6
PROPORTION OF EATING OCCASIONS.....PCT.....	4.7	0	6.8	4.1	4.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
23.9	26.4	24.7	27.4	31.2	35.2	35.9	36.2	28.9	27.4	16.8	24.4
16.9	18.9	17.2	17.0	19.6	23.1	21.4	23.0	18.4	17.0	8.5	15.6
5.6	6.2	5.5	8.2	7.9	9.1	10.2	9.9	7.2	7.6	6.7	5.2
1.5	1.3	2.0	2.3	3.7	3.0	4.3	3.3	3.3	2.8	1.7	3.6
1.4	1.4	1.5	1.6	1.6	1.5	1.6	1.6	1.6	1.6	1.7	1.7
.7	.7	.8	.8	.9	.8	.9	.8	.9	.8	.8	.9
5.0	6.0	5.0	5.0	6.0	6.0	5.0	6.0	5.0	5.0	4.0	6.0
11	11	14	15	17	14	16	14	13	10	15	11
11	11	13	19	15	13	21	12	13	9	16	9
2	2	2	2	3	2	3	2	2	2	2	2
5	5	5	5	7	5	7	5	5	5	5	5
9	9	10	10	10	10	11	10	10	9	11	10
15	12	20	20	21	20	21	20	15	11	20	15
21	21	30	31	32	28	31	27	30	21	21	23
29	30	41	40	42	40	42	40	44	30	46	26
57	55	54	74	82	61	77	62	56	45	122	52
26	24	31	30	35	30	32	28	24	22	26	21
28	22	25	26	26	23	24	22	20	18	19	15
22	26	33	33	33	31	35	28	26	20	26	25
14	21	22	42	22	21	51	20	24	16	29	17
23	22	24	31	33	28	29	29	30	20	34	19
16	15	20	15	26	19	18	18	15	13	16	11
279	256	147	470	250	245	768	245	169	188	250	78
279	256	340	940	495	360	1,518	308	250	243	365	184
23	23	28	29	31	28	30	26	24	20	26	20
22	21	23	32	25	22	35	21	19	18	20	15
5	5	5	5	5	5	5	5	5	5	5	5
10	10	15	15	15	15	15	15	15	8	15	8
16	16	21	29	29	29	29	16	16	15	16	16
30	30	31	31	32	31	31	31	31	29	31	30
47	47	61	59	61	59	61	59	44	32	46	32
63	63	64	63	66	63	63	63	59	47	59	59
90	120	125	125	125	120	123	120	125	104	125	64
250	256	128	470	250	245	768	245	125	188	125	78
44.9	48.3	20.2	36.8	93.4	131.4	162.1	165.4	24.1	23.0	5.9	20.3
6.3	6.0	4.0	5.9	4.7	4.5	5.6	4.1	4.6	3.0	3.9	5.0

TABLE 3.07.--MAYONNAISE-TYPE SALAD DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.9	.4	4.1	4.7	4.5
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	.4	3.3	3.4	3.0
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	.8	1.0	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	0	.3	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.0	1.3	1.4	1.5
STANDARD DEVIATION.....DO.....	.8	.0	.5	.7	.8
MAXIMUM.....DO.....	6.0	1.0	3.0	3.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	7	3	4	5	6
STANDARD DEVIATION.....GM.....	7	2	4	4	5
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2			1	2
25TH.....GM.....	3			2	3
50TH.....GM.....	5			3	5
75TH.....GM.....	10			6	10
90TH.....GM.....	15			10	13
95TH.....GM.....	20			15	15
99TH.....GM.....	39			15	21
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	17	9	10	11	14
STANDARD DEVIATION.....GM.....	17	5	9	9	10
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	16	0	11	10	10
STANDARD DEVIATION.....GM.....	12	0	11	6	6
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	14	0	0	8	11
STANDARD DEVIATION.....GM.....	10	0	0	3	7
MAXIMUM IN ANY 1 DAY.....GM.....	147	15	44	44	44
MAXIMUM OVER 3 DAYS.....GM.....	162	15	73	44	87
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	15	9	9	9	12
STANDARD DEVIATION.....GM.....	13	5	8	7	9
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2			2	2
25TH.....GM.....	5			5	5
50TH.....GM.....	15			5	10
75TH.....GM.....	15			15	15
90TH.....GM.....	29			15	29
95TH.....GM.....	44			22	29
99TH.....GM.....	59			29	44
MAXIMUM.....GM.....	120	15	30	29	44
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	102.3	0	3.0	3.5	1.9
PROPORTION OF EATING OCCASIONS.....PCT.....	3.7	0	5.5	3.0	1.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.2	4.6	3.5	4.1	5.9	5.1	5.6	4.7	4.5	4.9	3.3	5.9
3.4	3.4	2.7	3.3	3.3	3.4	3.6	3.4	3.8	3.3	2.7	4.9
1.2	1.0	.5	.8	1.8	1.3	1.1	1.2	.7	1.2	.6	.5
.5	.3	.3	0	.9	.3	.9	.2	0	.4	0	.5
1.5	1.5	1.5	1.3	1.7	1.5	1.6	1.4	1.2	1.6	1.2	1.5
.7	.8	.8	.6	1.0	.7	.8	.6	.4	.9	.4	1.2
4.0	5.0	4.0	4.0	5.0	3.0	5.0	3.0	2.0	5.0	2.0	6.0
7	7	11	8	10	7	8	7	7	8	8	6
6	7	12	6	8	6	8	6	6	10	7	5
2	2		2	2	2	2	2	2	2		2
3	3		3	5	2	3	2	3	3		2
5	5		5	9	5	5	5	5	5		5
10	8		10	15	10	10	10	10	10		7
15	15		20	20	15	16	15	15	20		10
20	24		20	29	20	20	20	20	39		20
34	39		24	34	29	39	29	29	44		20
15	17	25	19	20	17	17	15	20	20	24	17
9	19	30	16	16	15	17	15	18	26	21	14
17	15	32	19	20	14	19	16	18	9	9	6
13	14	24	8	11	9	13	12	9	8	6	1
15	11	18	0	17	14	13	10	0	22	0	10
8	7	7	0	12	12	8	5	0	14	0	(2)
66	118	147	59	88	88	120	118	88	118	60	60
110	118	162	73	162	118	120	118	88	132	60	60
15	14	22	18	18	15	16	14	18	15	19	12
10	14	18	14	13	12	14	13	12	20	16	11
5	5		5	5	2	5	2	5	2		5
5	5		10	10	5	5	5	10	5		5
15	10		15	15	14	15	15	15	10		7
15	15		28	28	15	15	15	29	15		15
29	29		44	30	29	29	29	29	29		29
29	29		44	44	30	44	39	44	44		30
59	59		59	59	59	59	59	59	118		59
66	118	88	59	88	73	120	118	59	118	59	59
8.1	3.7	2.2	6.4	15.4	9.6	21.8	12.4	1.3	3.4	2.5	7.2
5.0	2.4	2.9	8.3	3.8	2.4	4.9	2.7	2.1	2.5	11.3	8.2

TABLE 3.08.--FRENCH DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.6	0	1.5	3.9	5.9
ON ONLY 1 OF 3 DAYS.....PCT.....	5.2	0	1.3	3.5	5.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	.2	.4	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	0	1.1	1.1	1.2
STANDARD DEVIATION.....DO.....	.6	0	.3	.4	.5
MAXIMUM.....DO.....	4.0	0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	13	0	8	8	9
STANDARD DEVIATION.....GM.....	13	0	8	8	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2			2	2
25TH.....GM.....	5			3	3
50TH.....GM.....	10			5	5
75TH.....GM.....	16			10	10
90TH.....GM.....	21			21	21
95TH.....GM.....	32			21	21
99TH.....GM.....	48			42	21
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	31	0	22	21	25
STANDARD DEVIATION.....GM.....	34	0	22	21	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	0	26	18	18
STANDARD DEVIATION.....GM.....	20	0	4	20	8
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	27	0	0	16	11
STANDARD DEVIATION.....GM.....	17	0	0	0	6
MAXIMUM IN ANY 1 DAY.....GM.....	750	0	83	125	125
MAXIMUM OVER 3 DAYS.....GM.....	750	0	83	125	125
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	30	0	23	20	22
STANDARD DEVIATION.....GM.....	30	0	20	20	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5			5	5
25TH.....GM.....	16			5	10
50TH.....GM.....	30			16	16
75TH.....GM.....	31			30	31
90TH.....GM.....	63			63	47
95TH.....GM.....	63			63	63
99TH.....GM.....	125			125	63
MAXIMUM.....GM.....	750	0	83	125	125
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	103.2	0	.9	3.1	2.4
PROPORTION OF EATING OCCASIONS.....PCT.....	3.3	0	5.1	4.1	1.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.3	7.2	6.0	7.6	7.2	7.0	8.4	7.0	6.7	5.9	3.8	5.0
5.3	5.8	4.7	6.1	5.9	5.6	6.6	5.5	4.4	4.1	2.5	3.9
.9	1.3	1.2	1.2	1.2	1.1	1.4	1.3	1.9	1.0	1.2	.9
(2)	.2	.1	.3	.1	.3	.4	.2	.3	.7	0	.2
1.2	1.3	1.2	1.2	1.2	1.3	1.3	1.3	1.4	1.4	1.5	1.3
.5	.5	.5	.5	.5	.6	.6	.6	.7	.7	.8	.6
3.0	4.0	3.0	4.0	4.0	4.0	3.0	4.0	3.0	4.0	3.0	4.0
11	10	13	13	15	13	16	11	14	9	12	10
13	9	11	12	13	10	20	9	11	7	17	7
2	2	2	2	3	2	3	3	5	2		2
5	5	5	5	7	5	8	5	10	5		5
8	5	10	10	10	10	10	10	10	5		10
11	11	21	16	21	21	21	16	11	10		11
21	21	31	21	31	21	26	21	38	21		21
21	26	42	42	42	31	42	31	42	27		26
83	42	42	83	63	47	63	42	47	42		31
30	25	32	33	40	33	39	27	36	22	16	24
39	21	27	34	31	24	60	18	27	18	9	15
25	27	31	26	34	30	32	31	21	17	37	24
17	17	18	15	27	16	21	18	14	8	38	11
16	9	27	31	35	25	43	21	30	18	0	26
0	5	8	15	8	20	18	7	18	9	0	0
250	125	125	250	250	125	750	125	125	125	125	63
250	125	126	250	281	188	750	141	141	125	240	94
28	24	31	30	37	31	37	27	28	19	24	24
34	20	24	30	31	22	49	18	23	15	25	14
5	5	5	5	8	5	10	5	5	5		5
10	16	16	16	16	16	16	16	16	16		16
16	16	31	30	31	31	31	16	16	16		16
31	31	32	31	39	32	47	31	31	28		31
63	63	63	63	63	63	63	48	47	31		47
63	63	94	63	95	63	63	63	63	32		47
250	83	125	125	125	125	125	94	125	63		63
250	125	125	250	250	125	750	125	125	125	125	63
5.7	6.1	1.8	4.7	9.2	11.0	25.6	21.0	4.0	2.9	1.1	3.6
3.7	3.1	1.7	3.4	2.7	2.4	4.8	3.3	3.7	2.0	3.7	5.6

TABLE 3.09.--ITALIAN DRESSING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.1	0	1.3	3.9	5.5
ON ONLY 1 OF 3 DAYS.....PCT.....	6.1	0	1.3	3.0	4.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.7	0	0	.7	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	0	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	0	1.0	1.3	1.2
STANDARD DEVIATION.....DO.....	.6	0	.0	.6	.5
MAXIMUM.....DO.....	5.0	0	1.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	12	0	4	9	9
STANDARD DEVIATION.....GM.....	11	0	5	9	9
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2			1	2
25TH.....GM.....	5			2	5
50TH.....GM.....	10			5	7
75TH.....GM.....	15			10	10
90TH.....GM.....	20			20	20
95TH.....GM.....	34			29	20
99TH.....GM.....	52			39	49
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	0	12	19	23
STANDARD DEVIATION.....GM.....	22	0	15	17	21
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	28	0	0	28	20
STANDARD DEVIATION.....GM.....	23	0	0	17	13
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	0	0	7	49
STANDARD DEVIATION.....GM.....	17	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	470	0	59	59	118
MAXIMUM OVER 3 DAYS.....GM.....	940	0	59	118	146
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	28	0	12	21	22
STANDARD DEVIATION.....GM.....	21	0	15	18	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5			2	5
25TH.....GM.....	15			5	10
50TH.....GM.....	29			15	15
75TH.....GM.....	30			29	29
90TH.....GM.....	59			59	59
95TH.....GM.....	59			59	59
99TH.....GM.....	118			59	118
MAXIMUM.....GM.....	470	0	59	59	118
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	275.2	0	0	8.2	7.9
PROPORTION OF EATING OCCASIONS.....PCT.....	6.7	0	0	9.8	6.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.8	6.9	6.5	6.7	6.6	11.0	10.2	11.3	7.6	7.3	4.5	5.7
4.5	5.9	5.3	5.1	4.9	8.5	7.2	8.2	5.4	5.5	3.5	3.4
1.1	.8	.9	1.2	1.6	2.3	2.2	2.3	1.8	1.4	.9	1.7
.2	.2	.3	.4	.1	.2	.8	.8	.5	.4	.2	.5
1.3	1.2	1.3	1.3	1.3	1.3	1.4	1.4	1.4	1.3	1.3	1.6
.5	.4	.6	.7	.5	.5	.7	.7	.9	.6	.6	.7
4.0	3.0	4.0	4.0	3.0	4.0	5.0	5.0	5.0	4.0	3.0	3.0
12	10	12	17	13	13	15	12	11	9	11	12
11	9	10	30	10	10	12	9	9	8	10	11
2	2	2	2	3	3	3	2	2	2		2
5	5	5	10	5	5	7	5	5	5		5
10	10	10	10	10	10	10	10	10	7		10
15	10	15	20	20	15	20	15	15	10		15
20	20	26	29	20	20	30	20	20	20		29
39	20	39	44	39	29	39	29	29	20		39
65	39	59	78	49	49	59	39	49	49		52
25	25	31	31	33	32	33	28	22	20	24	22
23	18	23	19	26	26	24	18	11	14	15	14
29	31	34	58	29	28	30	25	29	24	32	30
11	26	25	95	17	13	18	13	18	20	21	22
48	25	22	45	35	23	35	27	24	14	29	16
19	5	8	21	9	17	17	17	13	7	0	6
118	118	118	470	206	235	147	118	103	118	78	78
195	235	176	940	206	235	235	294	147	146	122	155
28	26	30	39	31	30	31	26	23	20	25	23
21	20	23	53	22	20	21	17	15	17	18	18
5	5	5	10	5	5	7	5	5	5		5
15	15	15	15	15	15	15	15	10	10		15
29	28	29	29	29	29	29	28	21	15		15
29	29	29	44	30	30	44	29	29	29		29
59	59	59	59	59	59	59	59	44	30		59
59	59	78	78	59	59	73	59	59	59		59
118	118	118	470	118	118	118	78	59	118		78
118	118	118	470	118	146	118	118	59	118	78	78
14.8	16.4	2.9	6.4	21.4	46.8	67.0	59.1	9.3	9.1	0	6.1
9.6	9.4	2.6	4.8	6.3	6.1	9.4	5.3	7.6	5.3	0	6.9

TABLE 4.01.--FRESH GRAPEFRUIT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.7	0	1.1	1.0	1.5
ON ONLY 1 OF 3 DAYS.....PCT.....	2.6	0	1.1	.7	1.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	0	0	.2	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	0	.1	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER....	1.7	0	1.1	1.3	1.2
STANDARD DEVIATION.....DO.....	.9	0	.3	.6	.5
MAXIMUM.....DO.....	7.0	0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	90	0	55	66	64
STANDARD DEVIATION.....GM.....	58	0	24	31	39
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	38				
25TH.....GM.....	45				
50TH.....GM.....	89				
75TH.....GM.....	110				
90TH.....GM.....	165				
95TH.....GM.....	205				
99TH.....GM.....	268				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	163	0	164	159	155
STANDARD DEVIATION.....GM.....	67	0	73	71	55
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	162	0	0	139	195
STANDARD DEVIATION.....GM.....	59	0	0	15	75
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	167	0	0	134	0
STANDARD DEVIATION.....GM.....	65	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	804	0	268	330	402
MAXIMUM OVER 3 DAYS.....GM.....	1,876	0	268	402	670
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	159	0	145	149	158
STANDARD DEVIATION.....GM.....	58	0	57	56	64
PERCENTILES: 1/					
5TH.....GM.....	106				
25TH.....GM.....	134				
50TH.....GM.....	134				
75TH.....GM.....	165				
90TH.....GM.....	268				
95TH.....GM.....	268				
99TH.....GM.....	330				
MAXIMUM.....GM.....	660	0	268	330	402
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER....	4.9	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.6	2.4	2.2	2.3	2.7	3.7	6.0	7.4	8.9	12.5	7.9	10.4
1.3	1.7	1.7	1.5	2.0	2.1	3.2	3.8	3.5	5.7	2.9	5.6
.2	.5	.4	.4	.4	1.1	1.8	2.2	2.8	3.0	2.4	2.1
.1	.2	.1	.4	.3	.5	1.0	1.4	2.6	3.7	2.6	2.8
1.3	1.4	1.3	1.7	1.4	1.6	1.7	1.8	2.0	1.8	2.0	1.8
.9	.6	.5	1.0	.9	.7	.8	1.0	1.0	.9	.9	1.0
5.0	3.0	3.0	4.0	5.0	4.0	5.0	7.0	7.0	3.0	4.0	5.0
72	70	63	93	85	90	90	94	110	93	93	83
44	33	33	45	55	52	53	68	76	51	53	52
	35			45	35	45	37	45	38		22
	45			45	45	45	45	45	45		45
	55			55	89	89	89	89	89		55
	89			89	110	110	110	134	134		134
	110			155	165	165	160	268	165		165
	134			223	199	179	223	291	179		165
	179			268	268	268	330	330	231		244
169	165	151	200	183	182	160	162	148	149	142	133
63	63	55	75	69	103	61	60	35	50	23	43
162	150	154	196	197	176	170	158	166	152	133	132
58	38	56	74	78	64	61	58	77	40	8	13
186	134	134	142	187	149	160	183	194	153	149	157
44	0	0	13	62	46	49	91	69	34	56	33
330	330	536	469	495	660	460	804	660	330	536	330
670	536	620	603	804	804	804	1,876	990	804	1,072	732
160	153	150	159	178	169	160	160	167	152	141	136
56	50	68	57	67	77	61	58	59	42	30	28
	106			134	106	106	106	134	115		83
	134			134	134	134	134	134	134		134
	134			134	134	134	134	134	134		134
	134			230	165	165	165	165	165		134
	268			268	268	268	268	268	212		165
	268			330	330	268	268	330	268		165
	330			330	454	330	330	330	268		268
330	330	536	330	495	660	460	495	330	330	268	268
0	0	0	0	0	0	2.2	0	0	1.2	0	1.5
0	0	0	0	0	0	.4	0	0	.3	0	.8

TABLE 4.02.--GRAPEFRUIT JUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.6	.6	1.0	1.2	1.6
ON ONLY 1 OF 3 DAYS.....PCT.....	1.9	.3	.6	.6	1.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	.3	.5	.3
ON ALL 3 OF 3 DAYS.....PCT.....	.8	.3	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.9	2.8	1.5	1.6	1.6
STANDARD DEVIATION.....DO.....	1.4	2.0	.8	.6	1.1
MAXIMUM.....DO.....	14.0	5.0	3.0	3.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	131	133	79	95	100
STANDARD DEVIATION.....GM.....	115	103	66	44	70
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	41				
25TH.....GM.....	63				
50TH.....GM.....	83				
75TH.....GM.....	165				
90TH.....GM.....	250				
95TH.....GM.....	329				
99TH.....GM.....	658				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	219	123	110	205	185
STANDARD DEVIATION.....GM.....	121	61	60	119	59
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	223	0	212	183	239
STANDARD DEVIATION.....GM.....	114	0	29	38	86
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	245	247	226	146	214
STANDARD DEVIATION.....GM.....	167	0	0	0	77
MAXIMUM IN ANY 1 DAY.....GM.....	1,976	371	250	438	556
MAXIMUM OVER 3 DAYS.....GM.....	3,976	741	679	494	926
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	202	143	156	174	184
STANDARD DEVIATION.....GM.....	99	44	66	47	52
PERCENTILES: <u>1/</u>					
5TH.....GM.....	95				
25TH.....GM.....	125				
50TH.....GM.....	186				
75TH.....GM.....	247				
90TH.....GM.....	250				
95TH.....GM.....	375				
99TH.....GM.....	500				
MAXIMUM.....GM.....	1,500	217	250	250	309
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	4.9	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.3	1.5	1.7	2.2	2.5	3.8	5.0	6.4	5.0	5.4	4.2	3.7
.8	1.0	1.0	1.3	1.7	2.2	2.2	2.9	2.7	2.4	2.8	2.0
.3	.3	.6	.6	.7	1.0	1.5	1.6	1.4	1.7	.7	.8
.2	.2	.2	.3	.1	.6	1.3	1.9	.9	1.3	.7	.9
1.6	1.8	1.6	1.9	1.5	1.9	2.0	2.2	1.8	2.0	1.5	1.8
.9	1.4	.8	1.3	.6	1.4	1.3	1.6	.8	1.2	.8	1.1
3.0	8.0	3.0	5.0	3.0	9.0	10.0	14.0	3.0	7.0	3.0	6.0
106	103	136	133	116	145	137	141	110	118	90	109
68	80	100	95	66	158	110	121	61	103	49	90
				31	42	41	41	41	41		
				72	82	63	63	62	62		
				83	83	83	102	83	82		
				165	165	167	185	165	125		
				188	268	319	250	188	207		
				250	362	371	330	227	323		
				329	792	515	708	269	658		
194	230	294	238	262	256	208	208	235	173	175	185
76	162	307	91	139	125	99	108	154	58	48	59
249	204	263	224	250	230	253	207	203	183	237	210
100	148	100	82	96	112	158	103	69	56	25	107
182	191	223	353	200	338	219	255	162	229	148	183
55	82	72	30	41	296	129	152	54	152	31	128
500	813	1,500	556	750	1,976	988	1,235	682	741	313	741
750	993	1,500	1,235	988	3,976	2,161	2,717	806	1,976	563	1,616
194	173	248	210	234	229	202	196	187	179	181	178
73	72	202	66	104	107	108	93	82	92	53	72
				124	124	94	98	98	94		
				185	185	125	125	125	124		
				216	188	185	185	185	185		
				250	250	247	247	247	247		
				438	375	313	250	250	247		
				494	500	375	313	250	250		
				500	588	500	625	494	741		
371	438	1,500	500	500	750	988	988	494	741	313	494
0	0	0	0	4.9	0	0	0	0	0	0	0
0	0	0	0	3.4	0	0	0	0	0	0	0

TABLE 4.03.--FRESH ORANGES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.0	.9	8.1	10.0	12.6
ON ONLY 1 OF 3 DAYS.....PCT.....	5.7	.4	5.2	7.1	9.7
ON ONLY 2 OF 3 DAYS.....PCT.....	2.2	0	2.6	1.8	2.5
ON ALL 3 OF 3 DAYS.....PCT.....	1.1	.5	.3	1.1	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	2.2	1.5	1.5	1.4
STANDARD DEVIATION.....DO.....	.9	1.0	.7	.9	.9
MAXIMUM.....DO.....	8.0	3.0	6.0	6.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	78	63	57	68	63
STANDARD DEVIATION.....GM.....	54	25	30	46	48
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	24		24	21	22
25TH.....GM.....	48		33	48	48
50TH.....GM.....	48		48	48	48
75TH.....GM.....	97		67	82	73
90TH.....GM.....	145		97	145	115
95TH.....GM.....	180		120	169	145
99TH.....GM.....	290		145	265	338
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	147	110	125	140	135
STANDARD DEVIATION.....GM.....	66	36	40	47	57
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	158	0	123	143	159
STANDARD DEVIATION.....GM.....	63	0	42	56	62
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	167	82	120	161	212
STANDARD DEVIATION.....GM.....	61	8	34	50	97
MAXIMUM IN ANY 1 DAY.....GM.....	1,160	145	290	470	580
MAXIMUM OVER 3 DAYS.....GM.....	1,595	268	435	870	1,015
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	146	87	117	134	134
STANDARD DEVIATION.....GM.....	57	34	45	44	46
PERCENTILES: <u>1/</u>					
5TH.....GM.....	73		50	63	65
25TH.....GM.....	145		73	100	100
50TH.....GM.....	145		145	145	145
75TH.....GM.....	145		145	145	145
90TH.....GM.....	180		145	145	180
95TH.....GM.....	228		180	195	180
99TH.....GM.....	360		228	290	290
MAXIMUM.....GM.....	1,160	145	290	290	435
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	1.9	0	1.0	1.0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	(2)	0	.8	.4	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.7	11.2	8.9	9.4	7.3	8.0	9.1	8.9	7.1	9.4	10.7	10.7
7.1	7.7	5.5	6.8	4.4	5.3	4.6	5.7	4.8	4.8	4.8	6.0
3.1	2.5	2.4	1.9	1.9	1.9	2.9	1.9	1.1	2.9	1.8	2.5
.5	1.0	1.0	.8	1.1	.8	1.6	1.3	1.2	1.7	4.1	2.2
1.5	1.5	1.6	1.4	1.6	1.5	1.7	1.6	1.5	1.8	2.2	1.7
.7	.9	1.0	.8	.8	.8	.9	.9	.9	.9	1.4	1.1
5.0	6.0	6.0	6.0	5.0	5.0	5.0	5.0	5.0	4.0	6.0	8.0
75	69	87	68	85	79	88	78	79	88	103	78
47	50	76	43	56	53	52	52	73	59	90	49
30	17	24	24	33	30	30	24	24	30	8	10
48	48	48	48	48	48	48	48	48	48	48	48
48	48	60	48	60	56	67	48	48	67	73	48
97	97	97	97	97	97	120	97	97	108	102	98
145	145	145	120	153	145	157	145	145	145	290	145
168	152	196	167	180	150	180	193	277	193	337	178
205	242	420	205	290	242	242	242	360	338	362	180
155	134	149	145	158	159	160	146	133	141	137	133
68	66	53	63	54	77	95	60	42	44	58	52
164	156	182	143	171	172	150	159	137	165	121	143
53	60	119	36	78	89	40	50	22	57	29	31
184	167	209	169	167	161	160	166	207	166	178	151
73	56	98	43	51	37	46	53	95	66	103	41
580	580	1,160	580	540	1,080	900	456	540	435	505	300
1,160	1,085	1,595	870	900	1,440	1,015	905	1,080	1,080	1,085	800
150	137	158	142	157	156	151	145	155	150	141	136
51	49	84	51	54	72	57	52	74	51	60	41
73	50	73	73	84	84	72	73	74	90	73	73
145	145	145	145	145	145	145	145	145	145	90	145
145	145	145	145	145	145	145	145	145	145	145	145
145	145	145	145	145	145	180	145	145	145	180	145
180	180	180	180	218	180	180	180	180	180	180	180
228	180	290	180	290	290	200	228	360	290	290	180
293	290	435	290	360	360	360	290	435	360	360	228
580	540	1,160	580	390	1,080	900	456	435	435	360	290
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0

TABLE 4.04.--ORANGE JUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	35.5	20.9	40.9	41.7	43.7
ON ONLY 1 OF 3 DAYS.....PCT.....	13.9	9.8	14.3	17.1	17.4
ON ONLY 2 OF 3 DAYS.....PCT.....	9.3	5.6	10.5	11.5	11.8
ON ALL 3 OF 3 DAYS.....PCT.....	12.4	5.5	16.2	13.1	14.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.1	2.0	2.4	2.1	2.1
STANDARD DEVIATION.....DO.....	1.1	1.3	1.4	1.2	1.1
MAXIMUM.....DO.....	12.0	8.0	11.0	10.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	133	81	121	119	124
STANDARD DEVIATION.....GM.....	92	61	94	82	87
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	42	21	28	42	42
25TH.....GM.....	73	42	42	62	62
50TH.....GM.....	125	73	104	93	104
75TH.....GM.....	166	104	166	166	166
90TH.....GM.....	249	145	228	208	228
95TH.....GM.....	290	167	291	249	249
99TH.....GM.....	496	332	415	415	415
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	207	116	141	187	188
STANDARD DEVIATION.....GM.....	107	49	72	82	77
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	209	144	174	185	187
STANDARD DEVIATION.....GM.....	98	49	84	90	73
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	201	142	189	190	197
STANDARD DEVIATION.....GM.....	102	74	101	89	100
MAXIMUM IN ANY 1 DAY.....GM.....	1,992	498	996	1,058	1,245
MAXIMUM OVER 3 DAYS.....GM.....	3,237	1,370	2,739	1,994	3,237
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	190	122	153	167	178
STANDARD DEVIATION.....GM.....	84	51	70	73	68
PERCENTILES: <u>1/</u>					
5TH.....GM.....	95	62	62	93	95
25TH.....GM.....	125	93	125	125	125
50TH.....GM.....	187	125	125	125	187
75TH.....GM.....	249	131	187	187	248
90TH.....GM.....	249	187	249	249	249
95TH.....GM.....	311	249	249	249	249
99TH.....GM.....	498	250	374	498	374
MAXIMUM.....GM.....	1,240	250	500	747	747
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	126.0	0	2.9	8.3	4.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	.3	.5	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
39.4	41.0	37.3	36.6	32.9	31.8	32.0	35.1	32.7	40.4	28.6	36.7
16.7	16.4	15.2	18.3	14.8	13.8	12.1	12.7	9.5	11.0	8.0	10.2
9.7	11.8	10.0	9.4	8.7	9.0	8.0	9.4	7.9	8.5	5.2	9.2
13.0	12.8	12.0	8.9	9.4	9.1	11.9	13.0	15.3	20.9	15.4	17.2
2.1	2.1	2.0	1.9	2.0	2.0	2.1	2.1	2.3	2.4	2.3	2.3
1.1	1.1	1.0	1.0	1.0	1.2	1.0	1.1	1.1	1.1	.9	1.0
8.0	10.0	7.0	7.0	7.0	12.0	12.0	12.0	9.0	9.0	5.0	8.0
134	129	155	129	148	139	133	128	134	143	141	129
87	84	108	82	112	106	86	84	85	81	72	87
42	42	42	42	42	42	42	42	42	42	42	42
83	63	83	62	83	83	83	72	83	83	83	83
124	124	125	104	125	125	125	124	125	125	125	125
167	166	187	166	187	166	166	166	166	187	187	166
249	249	249	249	249	249	249	228	248	249	249	208
280	270	332	280	332	331	270	249	249	291	249	249
436	415	540	372	498	498	415	435	498	415	331	457
222	202	256	218	238	218	213	197	185	185	192	166
117	97	150	101	127	103	134	85	72	59	94	63
209	206	246	232	257	226	217	186	183	195	248	182
89	87	147	92	128	101	95	68	71	88	112	101
206	202	236	216	234	229	189	192	185	190	173	178
93	87	111	81	138	136	89	92	90	73	55	89
1,245	996	1,992	992	1,494	1,183	1,984	1,183	996	996	498	996
2,241	2,054	2,988	1,991	3,113	3,175	2,988	2,771	2,241	1,494	996	2,241
195	188	228	208	227	207	191	181	173	179	180	168
80	77	116	81	122	86	80	67	63	63	71	65
124	124	124	124	124	124	95	124	93	124	93	95
125	125	155	125	154	125	125	125	125	125	125	125
187	187	248	187	187	187	187	187	156	186	186	125
249	249	249	249	249	249	249	248	248	248	248	187
249	249	311	249	374	249	249	249	249	249	249	249
311	249	498	374	498	374	311	249	249	249	249	249
498	498	747	498	750	498	498	496	374	374	496	436
747	747	1,240	747	1,000	996	996	623	498	498	498	498
8.1	4.7	12.0	5.5	14.7	15.1	20.2	13.4	6.8	4.5	1.6	3.5
.5	.3	1.1	.6	.6	.4	.6	.3	.8	.3	.5	.4

TABLE 4.05.--RAW APPLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	18.2	1.7	23.6	23.8	25.8
ON ONLY 1 OF 3 DAYS.....PCT.....	11.3	1.3	14.9	16.8	16.0
ON ONLY 2 OF 3 DAYS.....PCT.....	4.8	.3	6.0	5.8	6.8
ON ALL 3 OF 3 DAYS.....PCT.....	2.2	0	2.6	1.2	3.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	1.2	1.6	1.4	1.6
STANDARD DEVIATION.....DO.....	.9	.4	.9	.7	.9
MAXIMUM.....DO.....	8.0	2.0	6.0	6.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	75	37	56	59	71
STANDARD DEVIATION.....GM.....	53	31	42	37	45
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	23		12	23	23
25TH.....GM.....	46		23	35	46
50TH.....GM.....	46		46	46	46
75TH.....GM.....	92		78	71	92
90TH.....GM.....	138		92	92	138
95TH.....GM.....	184		138	138	141
99TH.....GM.....	276		184	184	230
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	142	73	107	129	134
STANDARD DEVIATION.....GM.....	58	50	58	51	58
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	149	138	115	133	147
STANDARD DEVIATION.....GM.....	52	0	44	48	48
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	166	0	123	142	150
STANDARD DEVIATION.....GM.....	71	0	64	56	39
MAXIMUM IN ANY 1 DAY.....GM.....	1,272	138	848	552	700
MAXIMUM OVER 3 DAYS.....GM.....	3,816	276	1,272	966	976
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	141	94	105	124	132
STANDARD DEVIATION.....GM.....	49	51	44	39	41
PERCENTILES: <u>1/</u>					
5TH.....GM.....	69		35	69	69
25TH.....GM.....	138		69	106	106
50TH.....GM.....	138		106	138	138
75TH.....GM.....	138		138	138	138
90TH.....GM.....	212		138	138	138
95TH.....GM.....	212		138	212	212
99TH.....GM.....	276		212	212	276
MAXIMUM.....GM.....	636	138	276	424	530
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	4.0	0	0	0	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	(2)	0	0	0	.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
22.0	24.5	16.7	19.1	15.0	17.0	19.2	16.8	13.0	16.3	14.0	14.0
13.7	16.4	10.4	12.7	8.5	11.2	10.6	9.9	6.7	11.3	7.7	7.0
5.8	5.4	4.3	4.8	4.3	4.5	5.3	4.9	3.1	3.1	3.1	4.0
2.6	2.7	2.0	1.6	2.1	1.3	3.2	1.9	3.1	1.9	3.1	3.0
1.6	1.5	1.6	1.5	1.7	1.5	1.7	1.6	1.8	1.5	1.8	1.9
.9	.8	1.0	.8	.9	.8	1.0	.9	1.0	.9	1.0	.9
8.0	6.0	8.0	5.0	7.0	7.0	8.0	7.0	6.0	6.0	6.0	6.0
78	71	83	71	85	71	86	75	90	70	97	81
58	45	55	45	54	45	61	44	62	48	160	53
23	35	26	26	35	23	35	26	18	23	10	21
46	46	46	46	46	46	46	46	46	46	46	46
46	46	71	46	71	46	71	58	71	46	46	69
92	92	92	92	92	92	92	92	138	92	106	92
138	138	141	138	141	138	141	138	163	138	141	141
184	141	212	141	212	141	209	163	212	187	212	212
353	230	283	230	276	258	325	212	279	230	1,273	230
145	144	155	144	157	146	149	144	137	135	126	133
62	52	68	47	68	66	50	55	60	47	49	53
152	150	168	155	157	148	156	148	156	154	137	136
45	42	58	59	55	48	70	42	46	42	42	41
186	157	182	158	173	166	178	157	171	163	237	154
81	53	55	48	55	54	64	39	55	57	292	54
690	636	690	828	742	848	1,272	636	424	424	1,272	276
1,380	966	1,104	966	966	1,008	1,484	966	976	828	3,816	828
146	140	151	142	153	140	150	140	147	137	164	131
55	41	48	46	52	45	50	44	48	47	121	50
69	78	106	69	106	69	106	69	69	63	69	63
138	138	138	138	138	138	138	138	138	138	138	106
138	138	138	138	138	138	138	138	138	138	138	138
138	138	138	138	138	138	138	138	138	138	138	138
212	207	212	212	212	212	212	212	212	212	212	212
212	212	212	212	276	212	212	212	212	212	636	212
414	276	276	276	276	276	312	276	276	276	636	276
636	552	424	552	530	424	636	636	276	424	636	276
0	0	0	0	2.3	0	0	0	0	0	0	0
0	0	0	0	.2	0	0	0	0	0	0	0

TABLE 4.06.--APPLESAUCE AND COOKED APPLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.8	35.6	13.6	10.4	14.1
ON ONLY 1 OF 3 DAYS.....PCT.....	7.6	18.5	8.3	8.1	11.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.7	11.6	3.9	1.8	2.5
ON ALL 3 OF 3 DAYS.....PCT.....	.5	5.5	1.4	.5	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	2.2	1.7	1.3	1.2
STANDARD DEVIATION.....DO.....	.7	1.5	1.1	.7	.5
MAXIMUM.....DO.....	9.0	9.0	8.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	61	52	59	56	55
STANDARD DEVIATION.....GM.....	47	39	49	35	37
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11	9	9	11	16
25TH.....GM.....	38	23	21	41	32
50TH.....GM.....	43	43	43	43	43
75TH.....GM.....	85	66	85	85	83
90TH.....GM.....	127	99	128	85	85
95TH.....GM.....	160	135	170	128	128
99TH.....GM.....	253	209	255	213	170
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	146	102	109	135	136
STANDARD DEVIATION.....GM.....	95	68	79	75	80
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	139	95	114	125	130
STANDARD DEVIATION.....GM.....	78	51	63	41	68
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	137	89	141	139	140
STANDARD DEVIATION.....GM.....	88	52	62	45	30
MAXIMUM IN ANY 1 DAY.....GM.....	1,016	510	510	383	510
MAXIMUM OVER 3 DAYS.....GM.....	2,034	638	797	638	638
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	134	71	104	126	132
STANDARD DEVIATION.....GM.....	86	49	65	61	76
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28	14	16	32	32
25TH.....GM.....	64	42	57	65	64
50TH.....GM.....	128	57	97	128	128
75TH.....GM.....	130	92	128	128	128
90TH.....GM.....	255	128	213	255	255
95TH.....GM.....	255	135	255	255	255
99TH.....GM.....	488	255	255	255	480
MAXIMUM.....GM.....	765	510	255	383	510
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	32.6	.6	2.9	0	6.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	.2	1.2	0	1.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
13.6	11.1	10.2	7.7	5.1	5.8	8.1	9.5	10.7	11.5	15.1	17.4
11.8	9.2	8.5	6.7	4.3	5.1	6.8	7.5	7.5	7.9	10.5	9.2
1.5	1.7	1.5	.9	.8	.6	1.2	1.7	2.5	2.9	2.2	5.3
.2	.3	.2	.1	(2)	.1	.2	.4	.7	.7	2.4	2.9
1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.3	1.4	1.5	1.7	1.8
.4	.5	.5	.4	.4	.5	.5	.6	.6	.7	.9	1.0
3.0	4.0	4.0	4.0	3.0	4.0	3.0	5.0	4.0	4.0	4.0	6.0
58	55	69	57	68	51	65	59	56	62	89	85
53	42	73	33	46	40	45	47	30	41	67	57
10	11	15	16	21	10	11	10	11	11	21	16
31	28	32	41	43	21	41	32	40	41	40	43
43	43	43	43	53	43	43	43	43	43	85	84
83	80	85	85	85	71	85	85	85	85	127	112
128	85	149	85	128	85	128	127	85	127	202	170
128	138	170	128	169	128	166	168	95	129	213	198
340	213	255	170	339	170	255	254	128	170	340	255
142	137	157	158	184	129	165	142	147	139	207	170
92	86	110	86	130	82	99	97	74	84	130	90
187	130	184	129	154	156	179	135	108	132	136	157
129	78	90	54	67	56	94	73	56	52	54	88
222	158	376	150	127	146	72	153	77	131	174	141
106	80	235	0	0	121	47	92	23	56	98	63
765	510	1,016	383	1,016	510	638	638	255	454	636	510
1,275	765	2,034	510	1,016	893	893	1,020	510	760	1,020	765
151	134	171	146	171	130	158	134	121	128	161	141
107	82	125	73	98	79	92	84	71	67	89	80
32	26	48	48	64	16	31	23	21	32	30	26
84	64	96	122	128	64	120	64	64	65	113	90
128	128	128	128	128	128	128	128	128	128	128	128
191	128	244	191	199	128	255	130	128	128	255	191
255	255	255	255	255	255	255	255	255	255	255	255
255	255	510	255	383	255	255	255	255	255	255	255
512	383	510	383	508	284	510	383	255	255	383	383
765	510	762	383	510	510	510	610	255	386	383	383
2.6	1.5	1.0	0	3.7	3.0	3.1	2.4	.9	0	0	4.6
.8	.5	.6	0	1.5	.8	.6	.3	.6	0	0	1.5

TABLE 4.07.--APPLEJUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.8	19.2	13.1	8.5	5.5
ON ONLY 1 OF 3 DAYS.....PCT.....	2.2	9.3	5.1	3.8	3.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	4.5	4.6	2.8	1.6
ON ALL 3 OF 3 DAYS.....PCT.....	.6	5.4	3.5	2.0	.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.8	2.2	2.6	2.3	1.8
STANDARD DEVIATION.....DO.....	1.4	1.5	2.4	1.8	1.1
MAXIMUM.....DO.....	15.0	8.0	15.0	11.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	116	91	128	132	115
STANDARD DEVIATION.....GM.....	98	71	117	141	88
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	41	21	31	41	41
25TH.....GM.....	62	42	62	62	62
50TH.....GM.....	83	78	103	83	83
75TH.....GM.....	145	124	165	155	145
90TH.....GM.....	227	176	227	238	207
95TH.....GM.....	289	248	372	331	289
99TH.....GM.....	496	331	651	889	516
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	219	128	165	175	201
STANDARD DEVIATION.....GM.....	130	50	81	81	91
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	223	164	200	193	209
STANDARD DEVIATION.....GM.....	113	90	104	71	83
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	223	157	230	279	239
STANDARD DEVIATION.....GM.....	152	76	162	223	128
MAXIMUM IN ANY 1 DAY.....GM.....	1,302	496	868	1,302	1,178
MAXIMUM OVER 3 DAYS.....GM.....	3,224	1,240	2,356	3,224	1,550
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	191	125	148	170	193
STANDARD DEVIATION.....GM.....	101	56	64	65	87
PERCENTILES: <u>1/</u>					
5TH.....GM.....	63	47	62	93	124
25TH.....GM.....	124	93	124	124	124
50TH.....GM.....	186	124	124	155	186
75TH.....GM.....	248	131	186	186	248
90TH.....GM.....	248	217	248	248	248
95TH.....GM.....	372	248	248	248	248
99TH.....GM.....	496	250	250	372	496
MAXIMUM.....GM.....	992	250	496	496	992
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	14.4	0	2.9	3.7	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	.8	1.1	.9

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.0	4.0	2.7	3.1	2.4	3.7	2.3	2.7	1.6	2.8	3.8	3.1
2.0	2.8	1.9	2.4	1.7	2.6	1.4	1.7	.9	1.3	2.0	1.5
.9	.8	.5	.7	.5	.7	.6	.6	.4	.6	1.0	.6
.1	.3	.4	.1	.2	.3	.4	.4	.3	.9	.8	1.1
1.6	1.5	1.6	1.4	1.4	1.5	1.7	1.7	1.8	2.0	1.7	1.9
.8	.7	.8	.9	.8	.9	.9	1.2	1.1	1.2	.8	1.0
5.0	5.0	3.0	5.0	4.0	7.0	4.0	10.0	4.0	5.0	3.0	4.0
98	99	138	112	126	118	120	118	131	102	110	97
63	67	124	88	128	86	76	89	124	60	68	46
41	41	41	41	41	41	52	41	31	41	41	41
62	62	62	62	62	62	83	62	41	62	41	41
83	83	83	83	83	83	83	83	62	103	93	83
114	124	165	103	145	155	165	165	186	124	145	124
165	186	227	248	289	248	186	207	413	186	227	165
181	248	434	331	331	310	248	248	413	248	248	165
393	372	662	413	662	413	413	569	413	310	248	186
219	211	267	264	237	249	260	244	125	197	189	167
90	98	124	183	159	152	121	148	25	87	113	55
199	229	371	289	291	274	198	228	393	178	250	194
87	94	198	172	164	119	58	89	199	39	75	58
237	243	241	118	419	263	213	223	226	146	154	136
129	65	207	92	228	92	113	136	30	76	59	22
496	558	992	992	992	992	744	992	744	496	496	248
1,178	1,116	1,984	1,240	1,984	1,488	1,240	1,736	1,240	930	744	558
190	204	259	236	262	233	216	211	215	153	196	152
69	74	180	139	152	114	94	102	91	81	100	49
124	124	124	41	124	124	94	124	93	30	93	124
124	155	186	155	186	186	124	124	124	124	124	124
186	186	186	248	248	248	186	186	248	186	186	124
248	248	310	248	248	248	248	248	248	186	248	186
248	248	496	372	496	372	372	248	248	248	248	248
248	372	496	496	496	496	372	310	496	248	434	248
496	496	992	992	992	744	496	558	496	496	496	248
496	496	992	992	992	992	620	992	496	496	496	248
1.1	.9	0	0	0	0	2.2	0	0	2.0	0	0
1.1	.7	0	0	0	0	1.2	0	0	2.0	0	0

TABLE 4.08.--BANANAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	16.1	9.0	23.7	18.1	17.5
ON ONLY 1 OF 3 DAYS.....PCT.....	10.4	4.6	13.1	11.8	12.2
ON ONLY 2 OF 3 DAYS.....PCT.....	3.9	2.0	7.9	4.8	4.7
ON ALL 3 OF 3 DAYS.....PCT.....	1.9	2.4	2.8	1.4	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	2.0	1.8	1.5	1.4
STANDARD DEVIATION.....DO.....	.9	1.3	1.0	.8	.6
MAXIMUM.....DO.....	7.0	7.0	6.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	55	40	51	51	50
STANDARD DEVIATION.....GM.....	34	30	35	30	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		15	20	20
25TH.....GM.....	40		25	32	40
50TH.....GM.....	40		40	40	40
75TH.....GM.....	79		65	71	63
90TH.....GM.....	108		96	91	79
95TH.....GM.....	119		119	119	99
99TH.....GM.....	170		170	136	119
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	113	63	90	107	110
STANDARD DEVIATION.....GM.....	44	34	38	46	32
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	110	76	102	105	111
STANDARD DEVIATION.....GM.....	37	35	49	35	28
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	115	67	99	109	127
STANDARD DEVIATION.....GM.....	40	35	38	28	24
MAXIMUM IN ANY 1 DAY.....GM.....	544	179	357	276	272
MAXIMUM OVER 3 DAYS.....GM.....	816	476	714	476	595
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	106	61	86	99	109
STANDARD DEVIATION.....GM.....	37	31	34	34	30
PERCENTILES: <u>1/</u>					
5TH.....GM.....	50		30	48	56
25TH.....GM.....	95		60	60	95
50TH.....GM.....	119		95	119	119
75TH.....GM.....	119		119	119	119
90TH.....GM.....	136		119	119	136
95TH.....GM.....	136		119	136	136
99TH.....GM.....	238		136	238	224
MAXIMUM.....GM.....	476	136	272	276	238
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	32.4	.9	0	.9	5.9
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	1.1	0	.2	1.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.6	14.6	12.8	12.4	11.7	11.1	17.9	15.8	24.4	26.4	28.7	26.1
10.5	10.4	9.1	9.3	7.7	8.0	10.8	10.3	13.5	15.7	15.0	12.6
3.4	3.2	2.8	2.5	2.6	2.1	4.4	3.8	5.6	5.9	9.1	6.7
.6	1.0	.9	.6	1.4	.9	2.7	1.7	5.2	4.8	4.6	6.8
1.4	1.4	1.4	1.3	1.6	1.4	1.6	1.6	1.7	1.7	1.7	1.9
.7	.6	.7	.6	.9	.7	.9	.9	.9	.9	.8	1.0
5.0	4.0	5.0	4.0	6.0	5.0	6.0	7.0	6.0	5.0	4.0	6.0
53	52	57	48	65	50	62	53	61	57	57	62
30	27	35	26	43	31	36	34	36	35	36	38
20	20	20	13	32	20	20	20	20	20	20	20
40	40	40	40	40	40	40	40	40	40	40	40
40	40	40	40	40	40	45	40	45	40	40	40
79	71	79	48	79	60	79	65	79	79	79	79
79	79	119	79	119	85	119	95	119	103	95	119
119	119	136	95	136	119	125	119	119	119	119	127
159	125	181	159	238	159	181	182	198	198	136	159
119	121	124	110	131	110	121	109	113	108	103	105
44	46	62	36	56	37	51	39	37	36	38	34
118	110	130	108	124	103	116	110	105	102	104	98
39	34	38	33	35	36	37	38	25	28	30	37
128	103	129	113	149	122	120	112	111	111	108	108
34	29	26	32	51	32	30	45	36	39	45	33
397	408	456	272	476	357	544	357	271	272	272	285
595	544	595	476	812	598	714	714	714	680	816	714
114	110	121	109	125	106	113	102	106	103	102	98
38	32	50	35	49	35	37	32	30	33	37	31
60	60	59	48	60	39	60	48	57	48	57	60
119	95	119	95	119	95	95	68	95	68	68	60
119	119	119	119	119	119	119	119	119	119	119	119
119	119	119	119	119	119	119	119	119	119	119	119
136	136	136	136	136	136	136	119	119	136	119	136
136	136	140	136	238	136	136	136	136	136	136	136
238	238	456	238	357	238	238	152	190	238	272	136
357	357	456	272	476	357	476	285	266	238	272	238
5.2	0	0	0	0	1.7	5.1	3.6	1.6	3.6	1.6	2.2
1.2	0	0	0	0	.2	.4	.2	.3	.5	.6	.5

TABLE 4.09.--CANTALOUPE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.3	.2	1.1	1.5	2.2
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	.2	.9	1.0	2.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.2	.5	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.0	1.1	1.4	1.1
STANDARD DEVIATION.....CO.....	.8	.0	.4	.5	.3
MAXIMUM.....DO.....	6.0	1.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	81	45	26	57	49
STANDARD DEVIATION.....GM.....	61	0	11	29	31
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	21				
25TH.....GM.....	45				
50TH.....GM.....	68				
75TH.....GM.....	91				
90TH.....GM.....	181				
95TH.....GM.....	181				
99TH.....GM.....	272				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	180	136	79	164	133
STANDARD DEVIATION.....GM.....	107	0	31	91	75
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	181	0	35	92	148
STANDARD DEVIATION.....GM.....	87	0	19	37	64
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	192	0	0	0	0
STANDARD DEVIATION.....GM.....	96	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,362	136	136	363	272
MAXIMUM OVER 3 DAYS.....GM.....	1,632	136	136	363	408
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	171	136	68	125	135
STANDARD DEVIATION.....GM.....	91	0	35	73	76
PERCENTILES: <u>1/</u>					
5TH.....GM.....	61				
25TH.....GM.....	136				
50TH.....GM.....	136				
75TH.....GM.....	272				
90TH.....GM.....	272				
95TH.....GM.....	272				
99TH.....GM.....	529				
MAXIMUM.....GM.....	896	136	136	363	272
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	52.7	0	1.0	0	2.0
PROPORTION OF EATING OCCASIONS.....PCT.....	3.0	0	7.3	0	4.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.2	2.5	2.0	2.5	2.5	2.2	3.9	5.4	5.3	6.2	3.9	3.3
1.9	1.9	1.8	1.8	2.2	1.7	2.8	3.8	3.8	3.6	2.5	2.4
.3	.6	.2	.7	.3	.4	.9	1.1	.9	1.8	1.0	.3
0	.1	0	0	(2)	.1	.2	.5	.6	.8	.3	.5
1.2	1.3	1.1	1.4	1.1	1.3	1.4	1.5	1.5	1.7	1.5	1.5
.4	.7	.3	.6	.4	.6	.7	.9	.8	1.0	.6	.7
2.0	4.0	2.0	3.0	3.0	3.0	4.0	6.0	4.0	5.0	3.0	3.0
63	68	77	89	74	71	87	84	97	98	123	77
46	44	59	68	38	50	72	65	59	60	87	42
	6			27	11	23	21	45	45		
	45			45	45	45	45	45	45		
	57			85	68	91	68	91	91		
	91			91	91	91	91	91	112		
	91			107	107	160	181	181	181		
	107			136	181	181	227	213	181		
	272			208	272	453	272	272	227		
146	166	182	188	201	176	192	171	207	212	242	185
83	87	79	104	83	96	154	95	73	85	224	87
218	150	332	237	175	177	179	179	212	171	272	217
67	85	111	129	88	113	70	81	64	63	0	67
0	181	0	0	160	112	215	199	220	185	272	113
0	0	0	0	0	48	139	89	46	91	0	33
272	544	544	544	544	544	1,362	544	408	1,088	896	272
544	816	816	1,088	624	816	1,362	1,360	816	1,632	896	544
165	152	209	189	193	164	180	165	201	173	252	155
85	77	111	113	94	96	101	84	66	74	143	73
	32			68	32	68	40	136	80		
	136			136	80	136	136	136	136		
	136			160	136	136	136	160	136		
	204			272	272	272	272	272	272		
	272			272	272	272	272	272	272		
	272			320	272	272	272	272	272		
	272			544	544	681	320	272	408		
272	272	544	544	544	544	681	544	272	544	896	272
3.1	1.8	2.2	0	0	10.1	2.9	17.9	1.0	8.1	.7	1.8
5.9	2.5	7.2	0	0	6.6	1.1	3.1	1.2	4.4	2.4	3.6

TABLE 4.10.--FRUIT COCKTAIL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.9	1.4	4.6	4.5	8.3
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	1.4	3.4	3.9	7.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	1.0	.5	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.0	1.4	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.7	.5	.4
MAXIMUM.....DO.....	4.0	1.0	4.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	62	32	58	53	57
STANDARD DEVIATION.....GM.....	41	14	41	30	33
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11			14	21
25TH.....GM.....	43			28	42
50TH.....GM.....	43			43	43
75TH.....GM.....	85			85	85
90TH.....GM.....	85			85	85
95TH.....GM.....	149			99	126
99TH.....GM.....	184			149	170
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	163	96	130	139	157
STANDARD DEVIATION.....GM.....	93	43	69	74	84
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	158	0	149	142	155
STANDARD DEVIATION.....GM.....	75	0	71	34	52
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	180	0	170	127	150
STANDARD DEVIATION.....GM.....	82	0	0	0	21
MAXIMUM IN ANY 1 DAY.....GM.....	701	128	510	319	510
MAXIMUM OVER 3 DAYS.....GM.....	1,403	128	606	446	510
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	159	96	125	135	152
STANDARD DEVIATION.....GM.....	88	43	61	66	76
PERCENTILES: <u>1/</u>					
5TH.....GM.....	32			63	63
25TH.....GM.....	125			84	125
50TH.....GM.....	128			128	128
75TH.....GM.....	255			155	250
90TH.....GM.....	255			255	255
95TH.....GM.....	255			255	255
99TH.....GM.....	510			279	319
MAXIMUM.....GM.....	510	128	255	279	510
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	10.8	0	.9	1.6	3.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	1.4	1.7	2.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.2	7.5	5.2	3.8	1.9	2.6	3.3	3.1	5.5	3.2	2.6	3.9
6.5	6.7	4.9	3.5	1.7	2.4	2.8	2.8	3.7	2.5	1.7	3.4
.6	.7	.3	.3	.2	.2	.4	.3	1.4	.6	.6	.3
(2)	.1	0	0	0	0	.1	(2)	.4	.2	.3	.2
1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.1	1.4	1.3	1.4	1.2
.3	.4	.2	.3	.4	.3	.5	.4	.7	.6	.7	.5
3.0	3.0	2.0	2.0	3.0	2.0	4.0	4.0	3.0	3.0	3.0	3.0
56	60	61	59	76	58	73	55	81	71	86	61
30	35	35	38	57	31	44	33	56	72	83	41
21	21	21	10	21	11	21	10	16	5		
43	43	43	42	43	42	43	42	43	30		
43	43	43	43	64	43	74	43	64	57		
85	85	85	85	85	85	85	85	85	85		
85	85	85	85	170	85	128	85	170	135		
128	128	128	170	170	93	170	85	191	170		
170	170	170	170	276	170	255	170	255	468		
160	161	179	164	203	168	184	153	180	144	91	161
86	79	101	104	131	91	88	90	91	101	56	115
121	169	132	162	233	136	192	120	148	161	255	169
45	79	71	67	142	49	70	46	87	62	0	31
127	133	0	0	0	0	216	141	199	272	255	103
0	5	0	0	0	0	42	65	50	147	0	0
510	510	510	510	701	510	510	510	510	510	255	510
510	893	510	510	829	510	765	551	765	1,403	765	510
152	160	175	163	202	156	182	145	169	162	179	153
76	77	100	101	134	80	87	85	86	108	90	103
64	64	64	31	45	32	64	31	32	16		
125	128	125	123	128	123	128	90	97	90		
128	128	128	128	128	128	180	128	171	128		
250	255	255	255	255	255	255	191	255	250		
255	255	255	255	510	255	255	255	255	255		
255	255	383	255	510	255	255	255	255	383		
383	510	510	510	510	383	510	510	510	510		
510	510	510	510	510	383	510	510	510	510	255	510
.7	2.4	1.0	0	0	0	0	0	0	0	.8	0
.4	1.3	1.3	0	0	0	0	0	0	0	3.8	0

TABLE 4.11.--GRAPEJUICE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.8	2.6	5.2	6.8	5.9
ON ONLY 1 OF 3 DAYS.....PCT.....	1.9	1.8	3.3	4.8	3.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	.7	1.5	1.2	1.4
ON ALL 3 OF 3 DAYS.....PCT.....	.3	.1	.4	.8	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.6	1.7	1.5	1.7
STANDARD DEVIATION.....DO.....	.9	.9	.9	.9	1.2
MAXIMUM.....DO.....	7.0	4.0	4.0	7.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	114	66	93	92	106
STANDARD DEVIATION.....GM.....	95	57	66	68	86
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	32		21	32	32
25TH.....GM.....	63		42	42	63
50TH.....GM.....	84		84	63	84
75TH.....GM.....	148		127	116	127
90TH.....GM.....	211		169	179	190
95TH.....GM.....	253		211	190	253
99TH.....GM.....	464		337	316	569
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	233	181	198	193	221
STANDARD DEVIATION.....GM.....	149	151	125	97	103
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	232	79	211	186	164
STANDARD DEVIATION.....GM.....	126	54	117	81	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	246	211	142	201	264
STANDARD DEVIATION.....GM.....	157	0	56	102	141
MAXIMUM IN ANY 1 DAY.....GM.....	1,265	506	633	759	759
MAXIMUM OVER 3 DAYS.....GM.....	3,036	633	1,012	1,771	1,708
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	212	121	166	180	183
STANDARD DEVIATION.....GM.....	110	69	85	75	78
PERCENTILES: <u>1/</u>					
5TH.....GM.....	83		63	63	63
25TH.....GM.....	127		127	127	127
50TH.....GM.....	190		127	190	190
75TH.....GM.....	253		190	190	253
90TH.....GM.....	316		253	253	253
95TH.....GM.....	380		253	253	253
99TH.....GM.....	569		506	380	506
MAXIMUM.....GM.....	1,012	253	506	569	506
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	4.5	0	0	1.4	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	.8	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.9	4.8	2.2	2.0	2.1	3.5	1.3	1.9	1.9	2.2	.8	1.6
2.1	3.4	1.8	1.5	1.3	2.3	.8	1.2	1.1	1.5	.2	.5
.6	1.0	.2	.3	.3	1.0	.3	.5	.5	.5	.3	.9
.2	.4	.2	.2	.5	.2	.2	.2	.3	.2	.3	.2
1.5	1.6	1.4	1.4	1.8	1.6	1.6	1.6	1.7	1.5	3.6	2.0
.8	.8	.9	.7	1.0	.9	1.0	.8	.8	1.0	2.6	.8
4.0	6.0	5.0	3.0	5.0	6.0	6.0	4.0	3.0	5.0	7.0	3.0
100	106	141	126	170	122	114	100	122	105	322	150
70	79	117	93	104	120	88	66	104	97	304	85
32	42			42	42	21	21				
63	63			84	63	63	42				
84	84			127	84	84	84				
127	127			253	169	158	169				
200	210			337	211	211	190				
211	274			337	253	295	211				
464	506			464	1,012	422	253				
218	238	324	267	370	238	248	188	193	190	190	183
119	130	250	111	241	111	240	99	72	61	0	52
254	205	525	358	335	252	217	241	272	181	158	298
142	118	160	203	161	105	129	101	221	35	0	107
156	243	220	244	266	395	200	175	245	354	717	155
38	118	121	85	77	338	37	17	10	134	0	62
1,012	822	1,012	759	1,012	1,012	1,265	633	759	633	759	506
1,392	1,518	1,392	1,265	1,392	3,036	1,265	949	1,012	1,392	2,151	759
202	205	301	274	285	234	209	193	214	205	268	229
105	96	207	136	135	101	142	86	104	60	102	95
95	95			127	127	83	63				
127	127			253	190	127	127				
190	190			253	253	190	190				
253	253			316	253	253	253				
316	253			506	316	253	253				
380	316			506	506	380	380				
506	506			759	506	1,012	380				
759	822	1,012	759	759	506	1,012	633	506	316	506	380
0	0	0	0	1.3	1.8	0	0	0	0	0	0
0	0	0	0	.9	.6	0	0	0	0	0	0

TABLE 4.12.--RAW PEACHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	AGE (YEARS) AND SEX OF INDIVIDUAL				
	TOTAL	UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.5	1.2	3.5	3.8	4.5
ON ONLY 1 OF 3 DAYS.....PCT.....	3.0	1.0	2.7	2.6	3.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	.2	.6	1.0	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	.1	.2	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.6	1.2	1.4	1.6	1.5
STANDARD DEVIATION.....DO.....	1.0	.4	1.0	1.1	.9
MAXIMUM.....DO.....	7.0	2.0	7.0	7.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	84	46	60	67	74
STANDARD DEVIATION.....GM.....	66	28	58	50	52
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	25			25	30
25TH.....GM.....	51			33	51
50TH.....GM.....	51			51	51
75TH.....GM.....	101			96	101
90TH.....GM.....	152			101	152
95TH.....GM.....	203			135	152
99TH.....GM.....	355			355	304
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	164	104	121	135	160
STANDARD DEVIATION.....GM.....	86	38	55	58	82
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	176	152	152	155	179
STANDARD DEVIATION.....GM.....	74	0	15	33	95
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	224	0	355	197	152
STANDARD DEVIATION.....GM.....	100	0	0	115	0
MAXIMUM IN ANY 1 DAY.....GM.....	760	152	456	456	500
MAXIMUM OVER 3 DAYS.....GM.....	1,520	304	1,064	1,064	912
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	160	118	129	128	145
STANDARD DEVIATION.....GM.....	75	39	48	36	68
PERCENTILES: ^{1/}					
5TH.....GM.....	76			76	76
25TH.....GM.....	152			100	100
50TH.....GM.....	152			152	152
75TH.....GM.....	152			152	152
90TH.....GM.....	304			152	152
95TH.....GM.....	304			152	250
99TH.....GM.....	456			200	456
MAXIMUM.....GM.....	760	152	300	200	500
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	7.8	0	0	0	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	1.2

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.5	4.9	4.0	3.3	2.8	4.0	4.2	5.7	5.8	8.9	4.1	6.9
2.9	3.3	3.1	2.1	1.9	2.7	2.8	3.6	3.8	5.3	2.1	4.3
.5	1.3	.8	1.0	.6	1.0	1.1	1.5	1.6	2.8	2.0	2.3
.2	.4	.1	.2	.3	.3	.3	.7	.4	.8	0	.3
1.5	1.6	1.3	1.5	1.6	1.6	1.6	1.7	1.7	1.6	1.5	1.6
1.1	1.1	.5	.7	.9	1.0	1.0	1.0	1.1	.8	.5	.8
6.0	6.0	3.0	4.0	6.0	7.0	7.0	6.0	5.0	4.0	2.0	4.0
83	84	87	71	86	83	87	92	96	86	69	81
69	65	65	42	64	64	63	77	82	62	25	59
25	25	30		25	25	33	30	30	30		17
51	51	51		51	51	51	51	51	51		51
51	51	51		59	51	51	59	51	59		59
101	101	101		101	101	101	101	109	101		101
203	152	203		193	152	153	152	253	152		203
253	203	203		253	178	203	262	279	203		211
304	355	304		304	402	355	405	304	456		226
167	157	197	156	161	170	183	169	150	159	143	141
92	90	134	88	59	104	88	83	43	58	40	82
279	192	228	132	185	166	166	161	250	181	137	196
97	101	100	28	85	60	52	45	134	61	17	78
235	207	253	181	211	206	255	242	247	182	0	192
64	48	0	21	60	101	85	116	39	117	0	50
608	760	608	608	608	760	608	760	608	608	200	456
912	1,064	912	608	912	1,216	1,260	1,520	912	1,368	304	678
170	153	205	142	165	156	166	165	173	160	139	152
77	68	111	66	58	71	76	84	75	67	33	70
89	76	89		89	76	76	76	76	76		50
152	152	152		152	152	152	152	152	150		89
152	152	152		152	152	152	152	152	152		152
152	152	304		152	152	152	152	178	152		178
304	178	304		304	253	304	304	304	304		304
304	304	456		304	304	304	304	304	304		304
456	456	608		304	356	504	456	304	304		356
608	608	608	608	304	760	608	760	304	456	200	356
0	1.5	0	1.2	0	0	0	1.7	1.8	0	0	0
0	.9	0	1.7	0	0	0	.3	1.6	0	0	0

TABLE 4.13.--COOKED, CANNED, AND FROZEN PEACHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.1	28.3	10.5	6.9	12.5
ON ONLY 1 OF 3 DAYS.....PCT.....	6.4	16.4	7.2	6.1	11.0
ON ONLY 2 OF 3 DAYS.....PCT.....	1.3	8.7	3.0	.7	1.3
ON ALL 3 OF 3 DAYS.....PCT.....	.3	3.2	.3	.2	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER.....	1.3	1.9	1.4	1.2	1.1
STANDARD DEVIATION.....DO.....	.6	1.1	.6	.4	.4
MAXIMUM.....DC.....	7.0	7.0	4.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	68	51	57	58	54
STANDARD DEVIATION.....GM.....	52	35	39	35	35
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	20	9	10	16	15
25TH.....GM.....	42	28	28	32	32
50TH.....GM.....	49	42	43	43	43
75TH.....GM.....	85	70	82	83	71
90TH.....GM.....	128	96	96	85	91
95TH.....GM.....	171	113	128	128	123
99TH.....GM.....	256	162	192	171	171
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	166	107	135	158	143
STANDARD DEVIATION.....GM.....	104	59	84	93	81
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	153	104	119	132	132
STANDARD DEVIATION.....GM.....	76	54	68	61	38
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	178	84	123	106	205
STANDARD DEVIATION.....GM.....	120	45	42	30	86
MAXIMUM IN ANY 1 DAY.....GM.....	1,280	340	425	512	512
MAXIMUM OVER 3 DAYS.....GM.....	2,816	616	638	512	1,024
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	156	82	123	149	141
STANDARD DEVIATION.....GM.....	96	49	74	87	72
PERCENTILES: 1/					
5TH.....GM.....	48	14	31	57	48
25TH.....GM.....	96	56	64	96	96
50TH.....GM.....	128	68	112	128	128
75TH.....GM.....	192	113	128	192	192
90TH.....GM.....	256	135	220	256	256
95TH.....GM.....	288	170	288	256	256
99TH.....GM.....	512	224	384	512	384
MAXIMUM.....GM.....	1,280	288	384	512	384
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER.....	32.3	0	0	1.1	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	0	0	.8	.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.7	10.6	8.4	6.9	4.2	4.0	8.1	6.7	13.1	11.3	12.0	13.2
9.8	9.7	7.9	6.3	3.5	3.4	6.3	5.2	8.8	7.3	9.1	8.7
.9	.9	.5	.6	.6	.6	1.4	1.2	3.1	2.8	2.0	3.7
0	.1	0	0	.1	(2)	.4	.2	1.3	1.1	.8	.9
1.1	1.1	1.1	1.1	1.2	1.2	1.3	1.3	1.5	1.5	1.4	1.6
.3	.3	.2	.3	.4	.5	.6	.6	.8	.8	.7	.9
2.0	3.0	2.0	2.0	3.0	4.0	5.0	4.0	6.0	5.0	4.0	4.0
51	54	58	53	83	67	79	74	93	76	76	77
31	33	35	30	46	41	57	65	88	55	44	59
11	21	16	16	21	21	21	21	21	21	25	21
32	32	42	32	43	42	43	42	43	42	43	42
43	43	43	43	83	64	64	53	83	64	64	57
64	64	85	83	96	85	85	85	113	85	91	85
85	85	96	85	160	125	161	128	171	152	148	169
107	115	128	96	171	171	203	182	229	189	171	192
167	171	192	147	213	213	256	293	341	213	197	296
148	149	164	145	217	173	194	173	215	151	193	147
89	82	93	73	115	94	132	113	142	74	112	84
114	142	183	151	200	167	172	166	155	154	157	160
45	79	63	58	56	58	88	87	81	77	43	69
0	111	0	0	213	213	181	238	214	176	138	227
0	0	0	0	0	0	63	177	199	63	54	53
512	512	576	288	640	512	1,280	1,024	1,024	672	512	531
512	768	576	512	640	768	1,280	2,048	2,816	1,152	592	888
142	145	166	146	210	166	180	168	185	148	169	145
85	81	91	71	106	86	115	109	120	71	81	63
32	63	48	48	64	63	60	63	48	61	74	63
96	96	125	96	128	125	125	96	125	97	128	96
128	128	128	128	192	128	128	128	148	128	148	128
192	192	250	192	256	250	256	222	250	192	218	192
256	256	256	256	375	256	256	256	256	256	256	250
256	256	288	256	384	288	313	375	384	256	273	256
500	384	576	288	513	512	577	513	768	320	500	288
512	512	576	288	640	512	1,280	1,024	768	640	500	339
2.9	2.0	2.1	0	1.1	6.3	4.3	6.3	.9	0	1.3	2.2
1.2	.8	1.7	0	.6	2.4	.8	1.0	.4	0	1.5	1.1

TABLE 4.14.--RAW PEARS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.1	1.2	2.3	2.9	4.0
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	.3	1.7	2.3	3.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	.7	.4	.4	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.2	.2	.2	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.4	2.0	1.4	1.2	1.3
STANDARD DEVIATION.....DO.....	.8	.6	.7	.5	.7
MAXIMUM.....DO.....	6.0	3.0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	75	37	61	62	71
STANDARD DEVIATION.....GM.....	50	35	43	34	40
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	27			18	28
25TH.....GM.....	55			55	55
50TH.....GM.....	55			55	55
75TH.....GM.....	109			55	82
90TH.....GM.....	114			110	109
95TH.....GM.....	164			149	164
99TH.....GM.....	273			164	219
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	168	20	117	147	173
STANDARD DEVIATION.....GM.....	85	0	69	51	58
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	173	38	164	184	167
STANDARD DEVIATION.....GM.....	58	5	0	37	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	184	110	148	103	193
STANDARD DEVIATION.....GM.....	63	0	14	40	27
MAXIMUM IN ANY 1 DAY.....GM.....	2,132	164	328	328	492
MAXIMUM OVER 3 DAYS.....GM.....	2,132	329	492	492	656
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	163	56	131	150	163
STANDARD DEVIATION.....GM.....	69	40	43	57	42
PERCENTILES: ^{1/}					
5TH.....GM.....	82			55	82
25TH.....GM.....	164			141	164
50TH.....GM.....	164			164	164
75TH.....GM.....	164			164	164
90TH.....GM.....	164			164	164
95TH.....GM.....	328			200	164
99TH.....GM.....	328			328	328
MAXIMUM.....GM.....	2,132	164	164	328	328
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	.6	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	(2)	0	0	0	0

^{1/} SEE TABLE NOTES IN APPENDIX.

^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.7	3.3	3.2	1.4	1.8	2.1	4.0	3.9	4.9	4.3	3.5	4.2
2.4	2.6	2.8	1.1	1.1	1.6	2.9	3.0	3.0	3.4	1.7	3.2
.2	.7	.3	.2	.5	.4	.8	.6	1.6	.7	.6	1.0
(2)	.1	.1	0	.2	.2	.3	.3	.3	.2	1.2	0
1.2	1.3	1.3	1.2	1.6	1.4	1.4	1.4	1.5	1.3	2.2	1.3
.5	.7	.7	.5	.9	.7	.7	.9	.6	.6	1.1	.4
4.0	5.0	5.0	3.0	5.0	4.0	5.0	6.0	3.0	3.0	4.0	2.0
63	71	83	69	92	70	80	77	74	69	111	65
34	40	86	27	59	44	49	57	28	38	42	40
27	27			27	14	27	28	27	14		
55	55			55	55	55	55	55	55		
55	55			55	55	55	55	55	55		
55	55			109	91	109	72	109	82		
109	109			164	109	164	164	109	109		
164	164			273	164	164	219	109	164		
219	273			273	219	273	273	110	164		
165	163	230	179	168	155	177	166	172	157	252	157
65	51	268	62	51	71	89	44	57	48	126	36
182	177	164	165	181	159	180	192	153	182	164	160
34	42	0	1	40	13	49	87	27	42	0	96
219	195	168	0	210	178	187	218	82	164	150	0
0	82	5	0	57	48	46	76	0	0	14	0
492	492	2,132	328	328	492	656	656	330	328	492	328
656	820	2,132	332	820	656	820	1,230	330	492	492	656
163	161	195	167	171	156	171	166	149	162	151	153
46	42	219	57	47	56	57	63	39	46	44	64
82	82			82	82	82	82	82	82		
164	164			164	164	164	164	152	164		
164	164			164	164	164	164	164	164		
164	164			164	164	164	164	164	164		
164	164			168	164	165	164	164	164		
200	165			328	164	328	328	165	246		
328	328			328	492	328	410	246	328		
328	492	2,132	328	328	492	328	656	246	328	246	328
0	.6	0	0	0	0	0	0	0	0	0	0
0	.7	0	0	0	0	0	0	0	0	0	0

TABLE 4.15.--COOKED AND CANNED PEARS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.9	23.1	5.4	3.2	4.6
ON ONLY 1 OF 3 DAYS.....PCT.....	3.1	13.8	4.1	2.7	4.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	6.0	1.2	.5	.1
ON ALL 3 OF 3 DAYS.....PCT.....	.1	3.2	.1	(2)	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	1.3	2.1	1.3	1.2	1.1
STANDARD DEVIATION.....DO.....	.7	1.2	.5	.5	.3
MAXIMUM.....DO.....	8.0	8.0	4.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	65	52	47	63	49
STANDARD DEVIATION.....GM.....	46	32	29	50	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	21	9	16	19	11
25TH.....GM.....	31	28	28	31	31
50TH.....GM.....	53	43	43	45	43
75TH.....GM.....	84	72	63	63	63
90TH.....GM.....	125	99	75	101	94
95TH.....GM.....	157	128	104	188	94
99TH.....GM.....	251	135	207	302	128
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	156	111	111	149	141
STANDARD DEVIATION.....GM.....	86	65	47	81	78
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	166	101	112	189	144
STANDARD DEVIATION.....GM.....	90	46	31	119	36
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	148	89	207	188	94
STANDARD DEVIATION.....GM.....	73	31	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	752	424	250	564	383
MAXIMUM OVER 3 DAYS.....GM.....	1,504	424	620	907	383
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	149	75	112	154	138
STANDARD DEVIATION.....GM.....	85	44	47	89	74
PERCENTILES: ^{1/}					
5TH.....GM.....	42	14	47	47	32
25TH.....GM.....	94	42	77	94	94
50TH.....GM.....	128	64	97	128	128
75TH.....GM.....	188	112	133	188	188
90TH.....GM.....	255	128	188	255	255
95TH.....GM.....	282	135	188	282	282
99TH.....GM.....	383	224	250	510	383
MAXIMUM.....GM.....	752	224	250	510	383
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	6.5	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.9	5.5	3.1	2.7	2.0	1.8	3.9	3.7	5.2	6.2	5.8	5.6
4.3	5.1	2.7	2.3	1.6	1.5	2.8	3.1	4.4	3.9	4.2	4.0
.4	.4	.3	.3	.4	.3	1.0	.4	.6	1.6	1.6	1.3
.1	.1	.1	.1	(2)	0	.1	.1	.2	.6	0	.3
1.2	1.1	1.1	1.2	1.2	1.2	1.3	1.2	1.2	1.5	1.3	1.4
.4	.4	.4	.4	.5	.4	.5	.6	.5	.8	.4	.6
3.0	3.0	3.0	3.0	3.0	3.0	3.0	6.0	3.0	4.0	2.0	3.0
52	52	73	63	73	64	79	64	78	86	71	72
39	30	51	37	41	36	51	47	74	63	47	42
21	13			21	21	21	19	31	19		25
31	31			43	31	43	31	43	43		42
43	43			63	63	63	56	63	64		63
63	63			101	83	105	77	85	113		85
85	85			128	125	157	125	163	148		149
128	105			146	137	170	157	179	188		149
251	128			188	170	251	245	501	340		181
129	148	179	166	190	171	179	159	171	160	152	164
67	87	84	92	97	88	102	86	61	79	74	88
162	127	224	152	165	157	191	155	294	198	185	161
82	30	99	69	66	61	79	69	227	92	76	47
208	77	251	126	188	0	167	194	163	173	0	146
41	0	0	0	0	0	15	86	0	75	0	48
510	658	397	383	454	510	510	638	752	510	383	446
792	658	773	564	680	510	752	1,024	1,504	1,020	609	542
135	142	191	160	179	160	179	156	189	169	166	151
68	82	90	85	90	69	84	87	125	82	81	64
63	38			64	64	64	63	77	58		64
94	94			125	94	122	94	128	128		94
128	128			188	152	188	128	188	154		128
188	188			191	213	250	191	255	188		188
191	255			304	255	282	256	255	255		255
255	255			376	255	376	304	282	282		255
383	383			383	282	383	383	752	510		282
510	658	397	383	454	282	448	638	752	510	383	282
0	0	0	0	0	1.0	0	1.8	0	0	1.6	2.2
0	0	0	0	0	.9	0	.6	0	0	4.0	2.7

TABLE 4.16.--COOKED AND CANNED PINEAPPLE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.6	.5	2.2	1.6	2.8
ON ONLY 1 OF 3 DAYS.....PCT.....	2.2	.5	2.0	1.3	2.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	.2	.4	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.2	1.0	1.1	1.3	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.3	.5	.4
MAXIMUM.....DO.....	4.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	41	18	24	40	35
STANDARD DEVIATION.....GM.....	33	20	15	26	19
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	7				8
25TH.....GM.....	21				21
50TH.....GM.....	39				39
75TH.....GM.....	42				42
90TH.....GM.....	83				63
95TH.....GM.....	85				82
99TH.....GM.....	170				83
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	102	53	68	87	99
STANDARD DEVIATION.....GM.....	62	60	34	45	51
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	121	0	59	113	97
STANDARD DEVIATION.....GM.....	80	0	43	35	30
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	134	0	0	0	0
STANDARD DEVIATION.....GM.....	52	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	515	125	125	206	250
MAXIMUM OVER 3 DAYS.....GM.....	824	125	204	312	250
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	105	53	66	94	96
STANDARD DEVIATION.....GM.....	64	60	37	44	50
PERCENTILES: ^{1/}					
5TH.....GM.....	23				23
25TH.....GM.....	58				58
50TH.....GM.....	106				103
75TH.....GM.....	125				125
90TH.....GM.....	206				128
95TH.....GM.....	250				188
99TH.....GM.....	308				250
MAXIMUM.....GM.....	375	125	125	206	250
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	35.0	0	2.8	2.2	1.2
PROPORTION OF EATING OCCASIONS.....PCT.....	3.0	0	11.2	6.2	2.1

^{1/} SEE TABLE NOTES IN APPENDIX.

^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.6	3.2	2.0	1.9	1.7	2.6	2.2	3.4	2.8	3.7	2.0	4.6
2.5	2.9	1.8	1.9	1.4	2.4	1.8	2.7	2.4	3.3	1.7	3.5
.1	.3	.2	(2)	.2	.1	.4	.7	.2	.3	.3	1.0
0	0	0	0	(2)	.1	0	(2)	.2	.1	0	.1
1.1	1.1	1.1	1.0	1.2	1.1	1.2	1.2	1.3	1.2	1.3	1.3
.2	.4	.3	.2	.4	.4	.4	.5	.6	.5	.7	.6
2.0	3.0	2.0	2.0	3.0	3.0	2.0	3.0	3.0	3.0	3.0	4.0
36	51	40	35	47	38	43	42	47	36	46	49
24	51	28	20	32	30	32	38	29	23	21	39
10	12			15	8	6	5		10		
19	34			28	19	21	19		20		
39	42			41	34	39	37		34		
42	42			63	42	55	43		42		
61	82			83	83	83	82		83		
83	123			109	85	83	104		85		
167	275			168	170	167	250		96		
99	118	104	102	126	96	108	95	120	94	135	105
52	51	59	55	70	65	73	64	63	53	67	62
150	261	130	125	119	152	115	113	70	93	76	118
77	156	69	0	84	61	60	76	0	47	0	60
0	0	0	0	78	125	0	255	126	85	0	180
0	0	0	0	0	0	0	0	0	0	0	0
250	515	250	250	328	375	412	500	250	254	250	255
500	824	412	250	505	510	500	765	378	287	250	540
102	135	109	104	122	103	110	101	111	91	109	112
57	72	61	54	73	65	68	65	49	52	68	67
20	37			23	23	20	24		23		
58	103			63	58	58	58		58		
116	123			116	103	103	95		103		
125	125			142	125	125	125		125		
183	250			246	206	246	206		128		
250	309			255	250	250	250		159		
250	309			328	255	255	255		254		
250	309	250	250	328	375	309	313	250	254	250	255
1.0	0	1.1	0	6.3	11.5	0	3.9	0	1.0	4.2	0
1.8	0	3.6	0	8.3	7.5	0	1.3	0	1.3	31.0	0

TABLE 4.17.--RAISINS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.2	0	7.0	4.8	3.8
ON ONLY 1 OF 3 DAYS.....PCT.....	1.6	0	4.2	3.5	2.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	0	1.9	.8	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	.9	.5	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	1.4	0	1.6	1.4	1.5
STANDARD DEVIATION.....DO.....	.7	0	.8	.7	.7
MAXIMUM.....DO.....	5.0	0	4.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	16	0	15	14	15
STANDARD DEVIATION.....GM.....	14	0	12	11	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	2		1	1	1
25TH.....GM.....	6		6	6	6
50TH.....GM.....	12		12	12	12
75TH.....GM.....	24		21	22	21
90TH.....GM.....	30		35	28	28
95TH.....GM.....	48		36	29	29
99TH.....GM.....	73		48	37	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	37	0	29	32	31
STANDARD DEVIATION.....GM.....	33	0	23	20	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	35	0	28	31	36
STANDARD DEVIATION.....GM.....	23	0	18	12	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	28	0	29	27	22
STANDARD DEVIATION.....GM.....	21	0	17	20	10
MAXIMUM IN ANY 1 DAY.....GM.....	290	0	86	109	129
MAXIMUM OVER 3 DAYS.....GM.....	290	0	218	254	172
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	33	0	28	30	30
STANDARD DEVIATION.....GM.....	28	0	20	19	16
PERCENTILES: ^{1/}					
5TH.....GM.....	3		3	3	3
25TH.....GM.....	14		14	17	17
50TH.....GM.....	28		18	28	28
75TH.....GM.....	43		36	43	43
90TH.....GM.....	73		73	55	43
95TH.....GM.....	73		73	73	55
99TH.....GM.....	145		86	73	73
MAXIMUM.....GM.....	290	0	86	109	73
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	35.8	0	2.3	2.6	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	3.0	0	2.0	2.2	1.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.8	1.5	1.7	1.2	1.7	2.8	1.9	1.6	2.1	2.0	2.3	1.8
1.2	1.2	1.2	1.0	1.2	1.8	1.4	1.2	1.8	1.5	1.7	1.3
.5	.2	.5	.2	.2	.7	.3	.3	.2	.3	0	.6
.1	.1	0	0	.2	.3	.1	.1	.1	.2	.6	0
1.5	1.2	1.3	1.2	1.4	1.5	1.3	1.5	1.2	1.4	1.5	1.3
.7	.5	.6	.4	.8	.8	.7	.7	.5	.7	.9	.5
4.0	3.0	3.0	2.0	4.0	5.0	4.0	4.0	3.0	3.0	3.0	2.0
20	19	18	19	18	18	15	17	9	11	8	9
18	17	12	17	16	18	13	13	6	8	7	12
				1	2	2	2				
				6	6	6	6				
				14	12	12	15				
				19	24	18	24				
				48	30	36	29				
				48	64	48	36				
				73	97	48	63				
43	52	47	40	43	40	37	43	23	27	22	19
47	43	37	23	36	43	34	33	13	12	16	15
48	49	35	73	28	34	35	37	35	19	0	22
26	37	11	28	28	23	22	18	8	14	0	29
31	12	0	0	35	37	17	25	9	26	12	0
8	0	0	0	20	26	14	24	0	13	9	0
181	145	145	145	145	290	145	188	43	43	43	73
181	231	145	218	218	290	181	243	85	109	71	145
41	46	40	49	37	35	33	35	23	25	16	21
41	41	29	31	33	34	30	25	14	13	14	23
				4	5	2	3				
				17	17	14	14				
				28	28	28	36				
				43	36	43	48				
				73	73	73	73				
				109	96	73	73				
				145	145	145	145				
181	145	145	145	145	290	145	145	43	43	43	73
0	.9	.9	.9	2.4	0	5.2	7.5	3.4	2.0	3.2	3.0
0	2.2	2.9	4.3	2.5	0	4.2	4.6	12.1	4.1	16.3	12.8

TABLE 4.18.--RAW STRAWBERRIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.1	.2	1.5	1.2	1.6
ON ONLY 1 OF 3 DAYS.....PCT.....	1.6	0	.9	1.0	1.5
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	.2	.6	.1	.1
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	2.0	1.5	1.2	1.2
STANDARD DEVIATION.....DO.....	.8	.0	.8	.6	.4
MAXIMUM.....DO.....	6.0	2.0	4.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	46	80	45	28	34
STANDARD DEVIATION.....GM.....	44	0	45	17	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	12				
25TH.....GM.....	25				
50TH.....GM.....	37				
75TH.....GM.....	50				
90TH.....GM.....	99				
95TH.....GM.....	120				
99TH.....GM.....	225				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	101	0	75	70	88
STANDARD DEVIATION.....GM.....	60	0	34	38	42
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	115	120	110	87	134
STANDARD DEVIATION.....GM.....	76	0	88	18	96
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	139	0	0	0	0
STANDARD DEVIATION.....GM.....	93	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	600	150	298	149	373
MAXIMUM OVER 3 DAYS.....GM.....	1,273	240	596	225	522
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	100	120	87	69	87
STANDARD DEVIATION.....GM.....	58	30	41	34	44
PERCENTILES: <u>1/</u>					
5TH.....GM.....	37				
25TH.....GM.....	75				
50TH.....GM.....	75				
75TH.....GM.....	149				
90TH.....GM.....	149				
95TH.....GM.....	180				
99TH.....GM.....	298				
MAXIMUM.....GM.....	447	150	149	149	224
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	16.4	0	0	1.3	1.3
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	0	5.2	3.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.2	2.2	1.6	1.9	1.6	2.2	2.2	2.2	4.9	3.1	2.8	2.7
1.0	2.0	1.2	1.6	1.5	1.8	1.8	1.7	2.8	1.9	2.1	1.6
.1	.3	.4	.2	.1	.4	.3	.4	1.5	.8	.5	.8
.1	0	0	.1	(2)	(2)	.1	.1	.6	.4	.2	.3
1.5	1.2	1.4	1.2	1.2	1.3	1.4	1.4	1.8	1.6	1.4	1.6
1.2	.4	.7	.6	.7	.8	.8	.7	1.1	.8	.9	.7
5.0	3.0	3.0	3.0	6.0	6.0	6.0	4.0	6.0	4.0	4.0	3.0
49	36	57	34	45	43	48	47	62	53	51	53
47	24	39	28	49	49	49	41	44	44	25	55
				11	9	12	15	6	9		
				25	19	25	25	25	25		
				40	30	37	37	50	38		
				50	50	50	50	99	63		
				50	90	87	75	137	113		
				100	99	99	114	149	149		
				348	225	302	273	150	186		
94	96	130	74	105	98	106	103	117	107	131	91
60	55	69	43	54	61	61	48	101	66	63	40
137	94	141	127	190	97	141	139	114	79	111	69
31	55	73	50	134	56	63	106	40	50	40	31
139	0	0	74	221	320	142	74	137	136	62	194
79	0	0	0	111	99	132	39	13	47	0	54
298	315	298	199	447	600	596	447	447	338	298	298
671	447	522	348	1,043	1,273	1,192	820	450	559	298	745
95	91	121	82	115	97	106	101	103	98	105	102
53	50	63	45	66	60	65	59	46	55	60	69
				37	15	37	37	28	28		
				75	57	75	75	75	75		
				120	75	75	75	90	75		
				149	149	149	149	149	149		
				149	150	149	149	149	149		
				150	180	298	224	162	224		
				447	300	298	298	224	338		
298	315	298	149	447	300	298	298	224	338	298	298
0	0	0	0	2.5	6.2	2.2	3.0	0	0	0	0
0	0	0	0	3.2	4.0	1.5	1.4	0	0	0	0

TABLE 5.01.--YEAST BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	93.7	17.6	88.0	95.1	97.2
CN ONLY 1 OF 3 DAYS.....PCT.....	11.2	6.5	14.7	10.1	9.3
ON ONLY 2 OF 3 DAYS.....PCT.....	24.5	5.7	29.1	28.3	27.7
ON ALL 3 OF 3 DAYS.....PCT.....	58.0	5.5	44.1	56.7	60.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	4.1	2.5	3.5	4.0	4.1
STANDARD DEVIATION.....DO.....	2.0	1.6	1.8	1.8	1.8
MAXIMUM.....DO.....	14.0	8.0	9.0	10.0	11.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	62	17	33	47	55
STANDARD DEVIATION.....GM.....	40	12	22	26	29
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15	2	7	13	15
25TH.....GM.....	33	7	17	27	33
50TH.....GM.....	55	13	29	43	50
75TH.....GM.....	82	25	46	62	72
90TH.....GM.....	113	32	63	81	92
95TH.....GM.....	137	45	75	92	106
99TH.....GM.....	194	53	97	123	144
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	57	21	35	45	53
STANDARD DEVIATION.....GM.....	38	16	23	23	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	64	24	39	52	59
STANDARD DEVIATION.....GM.....	34	14	22	25	25
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	79	29	46	59	67
STANDARD DEVIATION.....GM.....	39	11	21	24	26
MAXIMUM IN ANY 1 DAY.....GM.....	838	100	262	277	352
MAXIMUM OVER 3 DAYS.....GM.....	1,588	160	427	520	578
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	46	20	28	36	40
STANDARD DEVIATION.....GM.....	26	11	16	17	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	21	7	11	13	21
25TH.....GM.....	25	11	21	25	25
50TH.....GM.....	44	20	25	28	40
75TH.....GM.....	50	25	40	50	50
90TH.....GM.....	75	28	50	50	56
95TH.....GM.....	100	50	50	57	75
99TH.....GM.....	140	50	80	100	100
MAXIMUM.....GM.....	700	75	193	242	228
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	638.4	0	9.5	29.2	36.0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	.3	.5	.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
96.9	96.4	96.2	93.7	96.1	92.9	95.5	92.9	97.2	95.3	96.7	97.4
9.9	13.4	9.2	14.5	11.1	15.8	8.9	11.8	7.0	7.4	7.3	8.2
27.3	28.4	23.6	28.8	24.8	29.5	19.5	23.5	18.4	22.9	13.9	16.5
59.6	54.5	63.4	50.3	60.2	47.6	67.2	57.7	71.7	64.9	75.5	72.7
4.2	3.8	4.4	3.8	4.1	3.5	4.5	3.9	4.7	4.3	5.0	4.5
1.9	1.8	2.0	1.9	1.9	1.8	2.1	1.9	2.1	2.0	2.2	2.1
11.0	10.0	14.0	10.0	12.0	10.0	14.0	14.0	12.0	10.0	12.0	12.0
69	57	87	56	83	52	80	54	71	54	68	54
41	32	52	32	50	32	47	33	40	31	36	29
17	13	17	13	17	12	17	13	17	15	17	14
40	33	50	31	47	29	47	31	43	32	42	32
63	52	79	50	75	47	73	49	65	50	65	50
90	75	115	75	111	70	105	72	92	71	92	72
121	98	155	98	150	93	138	95	121	94	119	91
143	113	178	114	177	111	162	111	147	110	136	107
193	152	250	148	237	150	232	149	188	161	168	144
64	60	80	58	78	55	66	50	69	43	57	48
39	42	59	34	47	33	40	29	35	22	31	35
70	64	87	65	86	59	75	58	63	53	60	52
34	29	47	30	44	26	37	30	33	26	29	26
87	73	106	73	105	71	96	68	83	66	78	63
40	30	49	29	48	30	44	31	38	30	33	27
767	535	838	390	740	526	549	554	340	400	380	300
1,588	755	1,363	644	1,269	757	1,012	1,099	776	658	575	650
49	44	59	44	61	44	53	41	45	38	41	36
28	23	35	21	35	22	29	21	23	19	22	17
22	22	23	22	25	22	22	21	22	20	22	18
28	25	40	28	44	28	40	25	25	25	25	23
50	44	50	44	50	44	50	42	44	34	44	28
50	50	70	50	75	50	56	50	50	50	50	50
80	57	100	57	100	57	100	56	70	56	56	50
100	84	114	80	114	80	100	75	88	60	75	57
142	125	170	113	170	114	150	113	124	105	113	100
617	510	638	340	700	476	454	454	234	340	280	236
30.5	36.1	22.2	20.7	91.2	97.9	90.9	100.5	22.3	22.5	11.0	17.9
.4	.5	.4	.4	.6	.6	.4	.4	.4	.3	.4	.4

TABLE 5.02.--WHITE BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	75.7	12.8	76.1	83.7	86.9
ON ONLY 1 OF 3 DAYS.....PCT.....	20.3	4.4	20.5	16.9	20.7
ON ONLY 2 OF 3 DAYS.....PCT.....	23.4	4.9	25.6	29.8	30.0
ON ALL 3 OF 3 DAYS.....PCT.....	32.0	3.4	30.0	37.0	36.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	3.1	2.4	3.0	3.2	3.1
STANDARD DEVIATION.....DO.....	1.8	1.3	1.7	1.7	1.7
MAXIMUM.....DO.....	14.0	7.0	9.0	9.0	10.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	45	17	27	37	40
STANDARD DEVIATION.....GM.....	33	12	19	23	25
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	8	4	4	8	8
25TH.....GM.....	19	7	13	19	20
50TH.....GM.....	37	16	24	33	33
75TH.....GM.....	61	25	37	50	53
90TH.....GM.....	87	32	53	67	75
95TH.....GM.....	108	39	65	81	87
99TH.....GM.....	158	53	93	112	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	51	22	31	40	46
STANDARD DEVIATION.....GM.....	31	17	20	20	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	56	28	36	48	51
STANDARD DEVIATION.....GM.....	29	14	20	23	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	68	28	41	52	59
STANDARD DEVIATION.....GM.....	34	10	19	21	23
MAXIMUM IN ANY 1 DAY.....GM.....	838	100	227	242	249
MAXIMUM OVER 3 DAYS.....GM.....	1,363	160	366	421	485
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	43	21	27	34	39
STANDARD DEVIATION.....GM.....	23	12	15	16	17
PERCENTILES: <u>1</u> /					
5TH.....GM.....	22	11	11	13	20
25TH.....GM.....	25	13	22	22	25
50TH.....GM.....	44	22	25	25	44
75TH.....GM.....	50	25	25	50	50
90TH.....GM.....	57	27	50	50	50
95TH.....GM.....	100	50	50	50	57
99TH.....GM.....	113	75	57	100	100
MAXIMUM.....GM.....	638	75	160	242	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	265.5	0	5.5	12.6	27.0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	.2	.3	.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
85.6	84.4	84.8	80.8	76.5	70.7	76.8	71.5	74.1	69.9	74.1	72.3
21.8	23.7	19.0	24.5	21.9	23.3	20.0	20.1	15.7	17.5	14.0	13.4
28.4	28.4	27.6	28.0	23.0	23.8	19.7	21.7	19.2	21.4	13.5	15.9
35.5	32.2	38.1	28.3	31.6	23.6	37.2	29.7	39.2	31.1	46.5	43.1
3.1	3.0	3.3	2.9	3.1	2.7	3.4	3.0	3.7	3.2	4.2	3.8
1.7	1.7	1.8	1.7	1.8	1.6	2.0	1.8	2.1	1.9	2.3	2.1
9.0	9.0	10.0	9.0	10.0	9.0	14.0	14.0	12.0	9.0	9.0	12.0
48	41	61	40	59	38	56	39	51	38	52	43
33	27	43	27	43	26	40	27	35	25	33	27
8	8	15	8	15	8	11	7	8	7	9	8
24	17	30	17	29	17	25	17	24	17	25	22
42	33	50	33	49	33	48	33	46	33	50	37
65	56	81	54	81	50	75	50	73	50	69	58
91	75	120	77	115	73	110	73	96	73	94	77
109	92	144	94	142	86	132	89	115	83	110	94
161	127	193	121	206	123	190	129	163	121	151	129
55	50	65	49	66	49	58	45	51	42	46	41
32	28	39	24	39	27	37	23	31	23	25	20
62	55	71	55	77	52	66	49	55	47	54	48
30	23	35	25	38	23	32	24	27	22	26	24
72	63	90	64	90	62	81	59	72	56	68	56
33	27	43	27	44	25	40	28	33	25	30	26
667	350	838	340	498	300	454	554	300	245	350	300
789	725	1,363	594	1,140	680	1,012	908	625	483	575	650
47	42	56	42	58	42	50	39	42	35	38	34
24	18	30	18	31	17	25	18	19	15	16	14
22	22	22	22	22	22	22	20	21	20	22	17
25	25	44	25	44	25	40	25	25	22	25	22
50	44	50	44	50	44	50	44	44	25	40	25
50	50	57	50	66	50	50	50	50	50	50	46
75	50	100	50	100	50	88	50	50	50	50	50
100	66	100	60	100	60	100	50	75	50	60	50
113	100	150	100	150	100	150	100	100	85	100	66
567	227	638	340	454	227	454	454	227	170	150	113
16.3	15.8	9.3	9.6	29.5	23.9	43.3	34.3	12.2	11.5	6.8	8.0
.3	.3	.2	.3	.3	.2	.3	.2	.4	.3	.4	.3

TABLE 5.03.--WHITE ROLLS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	36.8	1.9	24.5	35.4	46.3
ON ONLY 1 OF 3 DAYS.....PCT.....	25.4	1.8	19.7	26.8	30.5
ON ONLY 2 OF 3 DAYS.....PCT.....	8.8	.2	4.1	7.4	12.2
ON ALL 3 OF 3 DAYS.....PCT.....	2.6	0	.7	1.3	3.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.1	1.3	1.4	1.5
STANDARD DEVIATION.....DO.....	.8	.3	.6	.6	.8
MAXIMUM.....DO.....	8.0	2.0	4.0	4.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	25	7	14	18	21
STANDARD DEVIATION.....GM.....	18	4	9	10	13
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		5	7	9
25TH.....GM.....	13		9	13	13
50TH.....GM.....	19		13	13	15
75TH.....GM.....	28		17	23	27
90TH.....GM.....	47		27	31	40
95TH.....GM.....	60		28	38	48
99TH.....GM.....	93		40	50	69
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	52	19	35	41	46
STANDARD DEVIATION.....GM.....	30	11	20	18	23
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	54	16	35	42	46
STANDARD DEVIATION.....GM.....	26	0	10	11	17
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	58	0	33	40	43
STANDARD DEVIATION.....GM.....	26	0	6	12	14
MAXIMUM IN ANY 1 DAY.....GM.....	454	40	190	170	227
MAXIMUM OVER 3 DAYS.....GM.....	628	40	190	185	254
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	49	19	33	39	42
STANDARD DEVIATION.....GM.....	26	10	15	15	18
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28		14	19	28
25TH.....GM.....	35		20	28	28
50TH.....GM.....	40		30	40	40
75TH.....GM.....	56		40	40	40
90TH.....GM.....	80		40	56	64
95TH.....GM.....	100		56	70	80
99TH.....GM.....	140		80	85	100
MAXIMUM.....GM.....	354	40	160	170	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	37.6	0	0	2.2	2.3
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	.3	.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
52.9	46.8	49.7	42.7	43.0	36.7	38.7	31.5	26.1	23.8	16.8	18.3
34.1	31.6	30.6	27.9	28.0	27.1	26.6	23.1	17.9	16.1	12.0	11.8
14.2	11.7	13.7	12.1	11.8	8.0	8.4	6.5	6.2	6.0	3.6	5.0
4.6	3.5	5.3	2.7	3.3	1.5	3.7	1.8	1.9	1.7	1.3	1.5
1.6	1.5	1.7	1.5	1.6	1.4	1.6	1.4	1.6	1.5	1.5	1.6
.8	.8	1.0	.8	.9	.7	.9	.8	.9	.9	.8	.9
5.0	7.0	8.0	6.0	8.0	5.0	6.0	6.0	6.0	6.0	5.0	6.0
26	22	33	24	33	21	28	21	25	21	24	21
18	15	24	17	24	14	20	16	19	15	20	16
9	9	9	9	9	9	9	9	9	9	9	7
13	13	13	13	13	13	13	13	12	13	13	9
19	15	27	19	27	13	24	13	21	13	19	13
32	27	40	28	43	27	37	27	31	27	31	27
53	42	64	47	67	38	53	40	45	40	40	40
59	53	80	56	80	47	69	49	56	49	53	55
81	72	120	80	107	75	98	76	105	80	113	82
52	47	64	50	68	48	60	46	54	41	52	41
28	22	40	27	42	23	32	23	28	15	28	29
52	48	68	55	69	50	62	47	55	45	46	45
23	21	37	29	31	19	29	19	25	20	16	16
58	50	66	49	76	55	60	56	57	54	72	51
20	23	28	14	27	28	26	29	31	24	35	19
300	315	420	284	361	454	340	284	235	205	160	236
447	542	628	425	480	556	528	567	430	324	340	245
49	45	59	47	63	46	55	44	48	41	49	40
24	20	32	23	34	22	29	21	24	17	27	22
28	28	28	28	28	28	28	25	25	25	21	14
37	35	40	37	40	37	37	28	28	28	28	28
40	40	40	40	56	40	40	40	40	40	40	35
56	40	80	56	80	56	74	50	57	40	64	40
80	80	100	80	112	80	84	74	80	57	85	57
84	80	120	100	120	84	113	84	90	80	112	80
120	113	168	125	170	120	160	113	135	113	120	105
300	280	280	227	354	236	255	224	170	135	120	236
7.3	3.1	0	1.5	8.8	2.0	3.9	4.6	1.0	.8	0	0
.4	.2	0	.2	.3	.1	.1	.1	.2	.1	0	0

TABLE 5.04.--FRENCH, VIENNA, AND ITALIAN BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.2	.2	2.8	4.2	6.5
ON ONLY 1 OF 3 DAYS.....PCT.....	4.7	.2	2.1	3.5	5.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	.6	.4	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.5	0	.1	.2	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.0	1.3	1.2	1.2
STANDARD DEVIATION.....DO.....	1.0	.0	.5	.5	.5
MAXIMUM.....DO.....	12.0	1.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	26	9	15	15	16
STANDARD DEVIATION.....GM.....	31	0	18	17	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7			5	6
25TH.....GM.....	8			7	8
50TH.....GM.....	17			8	10
75TH.....GM.....	30			17	19
90TH.....GM.....	53			33	33
95TH.....GM.....	80			38	38
99TH.....GM.....	151			113	95
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	51	28	30	34	36
STANDARD DEVIATION.....GM.....	42	0	26	32	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	66	0	36	48	52
STANDARD DEVIATION.....GM.....	56	0	22	28	38
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	80	0	94	47	25
STANDARD DEVIATION.....GM.....	54	0	0	34	0
MAXIMUM IN ANY 1 DAY.....GM.....	567	28	113	227	227
MAXIMUM OVER 3 DAYS.....GM.....	908	28	284	340	340
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	54	28	37	38	39
STANDARD DEVIATION.....GM.....	45	0	30	32	30
PERCENTILES: <u>1/</u>					
5TH.....GM.....	18			15	18
25TH.....GM.....	25			20	25
50TH.....GM.....	40			25	25
75TH.....GM.....	60			40	50
90TH.....GM.....	113			61	80
95TH.....GM.....	113			113	113
99TH.....GM.....	227			227	198
MAXIMUM.....GM.....	567	28	113	227	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	19.5	0	0	1.1	1.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	1.2	.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.2	7.6	8.1	6.8	6.9	6.0	8.1	6.3	5.8	2.9	3.9	2.6
6.0	6.5	6.5	5.4	5.1	4.6	6.0	4.5	3.1	2.3	2.4	1.3
1.1	1.1	1.1	1.0	1.2	1.0	1.4	1.0	1.7	.3	.2	.9
.2	(2)	.5	.4	.6	.4	.7	.7	1.1	.3	1.3	.4
1.3	1.2	1.4	1.3	1.5	1.4	1.6	1.6	2.2	1.6	2.8	2.3
.6	.5	.7	.7	1.0	.7	1.1	1.1	1.8	1.4	2.5	1.8
5.0	3.0	4.0	5.0	7.0	4.0	7.0	8.0	12.0	7.0	8.0	7.0
21	17	27	22	40	23	32	26	28	23	42	28
22	14	26	25	48	26	32	33	25	27	45	23
7	6	7	7	8	7	8	5	7	4		
8	8	13	8	13	9	13	10	9	8		
15	13	17	13	23	17	20	17	20	13		
25	17	28	25	42	30	38	27	33	24		
40	38	66	57	95	44	76	43	63	57		
64	45	79	85	151	58	85	108	76	104		
95	76	140	113	265	151	173	151	148	111		
52	42	56	41	64	46	66	48	37	43	40	27
56	31	39	28	38	28	53	49	16	36	15	10
55	44	76	69	111	51	80	54	65	32	113	65
39	26	52	47	86	23	52	67	29	9	0	26
59	85	80	79	129	88	73	72	53	80	89	56
21	0	34	35	69	57	56	44	34	37	47	18
567	227	250	340	454	227	454	454	148	170	170	113
567	255	425	390	908	680	709	908	444	332	454	240
51	42	60	51	80	50	61	50	38	43	45	37
49	31	37	41	59	33	46	49	22	30	22	14
20	15	20	15	20	20	20	15	20	15		
25	20	36	25	40	25	30	25	25	23		
40	30	50	40	60	46	50	40	36	36		
57	50	75	57	113	57	75	57	40	50		
113	85	113	113	142	113	113	113	60	69		
120	113	142	113	184	113	142	113	80	113		
227	151	170	227	227	227	227	227	120	170		
567	227	227	340	454	227	454	454	120	170	113	70
0	1.2	0	0	4.1	4.5	1.7	3.8	2.0	0	0	0
0	.6	0	0	1.0	1.0	.3	.5	1.4	0	0	0

TABLE 5.05.--WHOLE AND CRACKED WHEAT BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.8	2.4	13.7	13.5	12.7
ON ONLY 1 OF 3 DAYS.....PCT.....	7.0	1.5	5.3	5.3	5.8
ON ONLY 2 OF 3 DAYS.....PCT.....	5.1	.6	4.9	4.2	3.9
ON ALL 3 OF 3 DAYS.....PCT.....	5.7	.4	3.4	4.0	3.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	2.6	1.6	2.3	2.6	2.2
STANDARD DEVIATION.....DO.....	1.7	.9	1.3	1.6	1.3
MAXIMUM.....DO.....	12.0	4.0	6.0	8.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	38	12	21	29	29
STANDARD DEVIATION.....GM.....	30	8	14	20	21
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	8		4	5	8
25TH.....GM.....	18		9	17	15
50TH.....GM.....	29		19	25	19
75TH.....GM.....	50		28	36	44
90TH.....GM.....	75		37	61	56
95TH.....GM.....	94		52	74	65
99TH.....GM.....	149		63	87	109
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	51	25	27	40	41
STANDARD DEVIATION.....GM.....	30	14	13	22	17
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	55	23	37	44	48
STANDARD DEVIATION.....GM.....	26	7	14	17	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	65	25	33	50	55
STANDARD DEVIATION.....GM.....	34	10	15	21	20
MAXIMUM IN ANY 1 DAY.....GM.....	362	53	128	140	150
MAXIMUM OVER 3 DAYS.....GM.....	972	103	196	268	373
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	44	23	27	34	40
STANDARD DEVIATION.....GM.....	22	10	14	16	16
PERCENTILES: <u>1/</u>					
5TH.....GM.....	21		11	13	21
25TH.....GM.....	25		14	25	25
50TH.....GM.....	50		25	28	35
75TH.....GM.....	56		28	50	56
90TH.....GM.....	56		56	56	56
95TH.....GM.....	84		56	56	56
99TH.....GM.....	112		56	57	84
MAXIMUM.....GM.....	227	50	100	112	100
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	13.2	0	.9	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	.3	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.7	10.4	12.5	11.0	19.0	18.8	19.9	20.5	24.5	28.4	25.7	26.3
4.7	5.9	5.5	6.0	8.5	7.9	7.0	8.4	7.3	7.6	5.3	6.8
3.4	2.4	3.7	3.0	5.4	6.4	6.0	5.7	5.2	7.7	5.8	5.0
2.5	2.1	3.3	2.0	5.2	4.4	6.9	6.4	12.0	13.1	14.6	14.4
2.4	2.2	2.4	2.1	2.4	2.3	2.7	2.5	3.3	3.2	3.7	3.4
1.6	1.5	1.7	1.4	1.6	1.3	1.8	1.6	1.9	1.9	2.1	2.2
8.0	9.0	9.0	8.0	10.0	8.0	9.0	10.0	9.0	9.0	9.0	12.0
40	31	50	29	49	32	46	31	49	38	54	38
32	23	47	20	39	21	36	22	31	26	36	26
8	8	9	8	12	8	9	8	8	8	9	8
17	9	19	17	19	17	19	17	25	19	25	19
28	19	35	19	37	28	37	25	44	33	50	33
56	44	69	37	67	45	58	44	65	52	73	48
83	64	112	57	108	60	93	59	92	70	105	75
96	81	140	66	120	75	117	74	113	86	113	85
141	112	231	99	200	105	170	101	131	125	168	140
55	50	61	49	71	48	55	46	66	48	42	44
42	28	30	27	43	25	26	21	50	25	13	27
67	53	70	54	75	53	61	46	63	46	54	45
32	22	37	18	36	18	30	19	26	17	16	16
74	65	101	58	91	57	80	53	69	55	75	52
36	21	57	17	42	23	38	23	29	27	32	26
346	165	324	170	362	224	282	196	255	217	224	154
636	336	972	302	724	411	728	492	501	584	504	420
49	43	62	43	62	42	52	38	45	36	44	34
23	18	33	16	31	16	24	15	21	14	18	15
25	23	25	23	25	23	24	21	24	21	25	20
28	28	50	28	50	26	28	25	28	25	28	25
50	50	56	50	56	50	50	28	50	28	50	28
56	56	57	56	56	56	56	50	56	50	56	50
73	56	112	56	112	56	84	56	69	56	56	56
100	57	112	56	112	56	112	56	84	56	75	56
112	112	168	75	168	100	112	75	112	63	112	56
224	138	224	170	227	138	225	140	170	100	125	125
0	1.5	0	0	2.3	2.6	2.3	1.0	0	0	0	2.6
0	.3	0	0	.1	.1	.1	(2)	0	0	0	.3

TABLE 5.06.--RYE AND PUMPERNICKEL BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.0	.4	1.9	1.7	2.0
ON ONLY 1 OF 3 DAYS.....PCT.....	4.3	.4	1.4	1.2	1.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.6	0	.5	.3	.6
ON ALL 3 OF 3 DAYS.....PCT.....	1.2	0	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.9	1.0	1.6	1.5	1.7
STANDARD DEVIATION.....DO.....	1.3	.0	1.1	1.0	1.0
MAXIMUM.....DO.....	9.0	1.0	5.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	28	3	17	17	21
STANDARD DEVIATION.....GM.....	24	1	17	14	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	8				
25TH.....GM.....	17				
50TH.....GM.....	17				
75TH.....GM.....	33				
90TH.....GM.....	55				
95TH.....GM.....	73				
99TH.....GM.....	112				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	9	25	29	38
STANDARD DEVIATION.....GM.....	24	2	13	13	22
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	54	0	54	55	47
STANDARD DEVIATION.....GM.....	26	0	34	26	24
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	61	0	37	38	43
STANDARD DEVIATION.....GM.....	34	0	0	4	7
MAXIMUM IN ANY 1 DAY.....GM.....	567	11	123	123	122
MAXIMUM OVER 3 DAYS.....GM.....	794	11	223	198	200
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	45	9	31	33	37
STANDARD DEVIATION.....GM.....	22	2	17	14	16
PERCENTILES: <u>1/</u>					
5TH.....GM.....	22				
25TH.....GM.....	25				
50TH.....GM.....	50				
75TH.....GM.....	50				
90TH.....GM.....	57				
95TH.....GM.....	100				
99TH.....GM.....	113				
MAXIMUM.....GM.....	340	11	73	57	112
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	1.1	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	(2)	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.9	2.7	4.4	3.8	7.9	5.9	12.6	9.6	11.9	9.3	8.6	7.6
1.9	1.9	2.9	3.1	5.2	4.4	7.4	5.6	5.8	4.9	3.8	2.7
.7	.7	.8	.2	1.4	1.1	2.5	2.5	3.4	2.1	2.2	2.6
.2	.1	.6	.4	1.3	.4	2.6	1.5	2.8	2.3	2.6	2.4
1.5	1.4	1.8	1.4	1.7	1.5	2.0	1.9	2.1	2.1	2.5	2.4
.9	.7	1.3	1.1	1.1	1.0	1.5	1.2	1.4	1.4	1.8	1.7
6.0	5.0	6.0	6.0	7.0	7.0	8.0	7.0	7.0	7.0	6.0	9.0
26	21	34	20	31	21	36	24	32	26	35	29
33	14	28	15	25	15	32	18	25	20	25	22
7	7	8	7	8	7	8	8	8	8		8
15	8	17	9	17	15	17	15	17	8		17
17	17	17	17	17	17	23	17	25	17		25
33	31	50	17	36	25	48	33	38	33		33
41	33	79	48	67	36	67	46	50	53		67
57	50	93	50	83	50	100	55	81	67		83
265	92	108	83	112	95	183	88	144	77		100
43	45	56	44	53	45	56	45	47	41	46	39
20	22	38	16	25	19	26	21	20	31	22	17
58	46	83	43	69	54	61	46	54	41	70	43
39	13	39	20	32	28	25	18	19	16	46	11
79	71	74	55	69	46	78	49	61	49	56	48
77	21	19	14	25	15	43	24	33	20	12	29
567	100	227	100	225	170	300	340	255	200	163	150
794	275	325	250	468	296	700	590	432	294	300	300
51	44	58	42	54	41	52	39	47	36	42	36
41	17	30	16	21	16	24	18	21	14	23	16
22	22	25	22	22	22	24	22	22	22		22
25	25	46	25	50	25	44	25	25	25		25
50	50	50	50	50	50	50	44	50	28		25
50	50	50	50	50	50	50	50	50	50		50
57	56	100	50	100	50	100	50	57	50		57
113	57	100	64	100	56	100	56	100	50		75
227	100	150	100	113	113	128	88	113	50		75
340	100	227	100	142	113	200	227	113	113	150	75
0	0	0	0	0	0	0	1.1	0	0	0	0
0	0	0	0	0	0	0	.1	0	0	0	0

TABLE 5.07.--HIGH FIBER, WHEAT GERM, BRAN, OATMEAL, AND MULTIGRAIN BREADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.6	.4	.9	.8	1.0
ON ONLY 1 OF 3 DAYS.....PCT.....	.7	.4	.4	.5	.5
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	0	.4	.1	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	.1	.1	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	2.4	1.0	1.8	1.7	1.8
STANDARD DEVIATION.....DO.....	1.6	.0	.6	1.1	.9
MAXIMUM.....DO.....	9.0	1.0	3.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	38	7	15	20	25
STANDARD DEVIATION.....GM.....	31	4	8	20	13
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	8				
25TH.....GM.....	17				
50TH.....GM.....	31				
75TH.....GM.....	49				
90TH.....GM.....	79				
95TH.....GM.....	102				
99TH.....GM.....	139				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	53	22	22	29	47
STANDARD DEVIATION.....GM.....	31	11	5	18	16
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	61	0	34	40	44
STANDARD DEVIATION.....GM.....	29	0	6	7	18
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	72	0	15	62	41
STANDARD DEVIATION.....GM.....	36	0	0	11	7
MAXIMUM IN ANY 1 DAY.....GM.....	248	34	56	84	81
MAXIMUM OVER 3 DAYS.....GM.....	627	34	81	221	156
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	48	22	25	36	41
STANDARD DEVIATION.....GM.....	24	11	13	14	15
PERCENTILES: <u>1</u> /					
5TH.....GM.....	23				
25TH.....GM.....	28				
50TH.....GM.....	50				
75TH.....GM.....	57				
90TH.....GM.....	68				
95TH.....GM.....	100				
99TH.....GM.....	136				
MAXIMUM.....GM.....	200	34	56	56	75
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	0	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	0	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
.7	1.0	.8	.7	1.4	1.4	1.9	2.4	2.6	3.4	2.1	2.4
.4	.6	.4	.3	.5	.6	.7	1.2	1.2	1.3	1.2	1.0
.3	.2	.1	.2	.5	.6	.7	.7	.4	.7	.2	1.1
(2)	.2	.3	.2	.4	.2	.6	.5	1.0	1.4	.8	.3
1.7	1.6	3.0	2.4	2.6	2.2	2.8	2.2	2.8	2.6	3.0	2.6
1.0	.8	2.5	1.4	1.7	1.4	1.9	1.4	2.0	1.7	2.3	1.5
5.0	3.0	8.0	5.0	9.0	7.0	9.0	6.0	8.0	8.0	7.0	7.0
29	23	55	40	59	31	49	31	59	41	41	32
19	12	38	26	42	21	30	22	51	34	43	17
				17	8	9	8		8		
				25	15	25	15		15		
				52	24	45	23		37		
				89	45	65	39		57		
				102	65	88	59		66		
				139	79	121	79		99		
				209	83	136	95		170		
49	54	72	51	77	39	79	49	47	42	48	60
30	25	47	12	33	15	46	23	16	25	33	27
60	36	68	54	79	61	73	54	68	60	72	52
24	11	0	8	34	29	37	20	41	18	0	16
64	42	99	80	105	54	75	59	112	65	76	55
0	9	19	2	41	13	26	24	33	38	53	23
130	112	146	112	227	173	204	136	248	204	168	126
200	169	357	248	627	250	408	316	496	510	392	270
50	44	54	51	69	42	52	42	62	47	40	36
23	16	21	16	32	22	26	17	28	17	15	13
				25	23	23	23		24		
				50	25	31	25		31		
				68	34	50	46		46		
				92	50	68	56		66		
				100	68	75	62		68		
				113	68	100	68		68		
				200	136	136	102		68		
112	69	100	68	200	136	170	136	124	84	56	62
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0

TABLE 5.08.--ENGLISH MUFFINS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.0	0	2.7	4.2	4.2
ON ONLY 1 OF 3 DAYS.....PCT.....	3.2	0	2.4	2.9	3.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.3	0	.3	1.0	1.0
ON ALL 3 OF 3 DAYS.....PCT.....	.5	0	.1	.4	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	0	1.2	1.4	1.3
STANDARD DEVIATION.....CO.....	.7	0	.5	.6	.5
MAXIMUM.....DO.....	5.0	0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	28	0	15	20	25
STANDARD DEVIATION.....GM.....	18	0	11	13	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9			8	9
25TH.....GM.....	19			9	17
50TH.....GM.....	19			19	19
75TH.....GM.....	38			19	33
90TH.....GM.....	57			38	38
95TH.....GM.....	57			50	57
99TH.....GM.....	95			76	114
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	60	0	38	42	62
STANDARD DEVIATION.....GM.....	27	0	22	15	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	60	0	38	43	60
STANDARD DEVIATION.....GM.....	24	0	12	17	35
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	57	0	52	43	30
STANDARD DEVIATION.....GM.....	23	0	0	19	14
MAXIMUM IN ANY 1 DAY.....GM.....	228	0	113	113	228
MAXIMUM OVER 3 DAYS.....GM.....	528	0	157	227	341
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	57	0	39	43	58
STANDARD DEVIATION.....GM.....	25	0	20	18	36
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28			14	25
25TH.....GM.....	50			28	28
50TH.....GM.....	57			50	57
75TH.....GM.....	57			57	57
90TH.....GM.....	100			57	114
95TH.....GM.....	114			57	114
99TH.....GM.....	142			113	227
MAXIMUM.....GM.....	228	0	113	113	228
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	0	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	0	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.7	4.6	4.9	4.4	5.9	5.5	5.2	5.9	4.7	4.8	1.3	3.3
3.2	3.4	2.9	2.7	3.8	3.9	3.1	3.7	2.8	2.9	.9	1.9
1.3	.9	1.4	1.6	1.4	1.4	1.4	1.4	1.5	1.4	0	.7
.3	.3	.5	.1	.6	.2	.7	.8	.4	.5	.4	.7
1.4	1.4	1.5	1.4	1.5	1.4	1.6	1.6	1.5	1.5	1.6	1.7
.7	.6	.7	.6	.7	.6	.8	.8	.6	.7	.9	.8
4.0	3.0	4.0	3.0	4.0	4.0	5.0	5.0	3.0	3.0	3.0	3.0
31	27	39	29	32	26	31	28	30	27	24	29
24	15	26	19	20	14	16	18	15	19	11	16
9	9	17	9	9	10	17	9	17	9		
19	19	19	19	19	19	19	17	19	17		
19	19	33	19	19	19	25	19	19	19		
38	38	55	38	38	36	38	38	36	36		
57	50	71	38	57	48	57	55	57	50		
76	57	105	67	76	57	57	57	57	57		
85	76	114	133	114	75	76	95	67	113		
60	64	77	60	69	59	65	55	58	49	52	57
25	28	43	22	34	21	25	23	16	18	3	20
70	61	76	68	61	60	59	56	63	51	0	59
23	30	30	32	24	19	24	22	18	14	0	18
88	49	84	36	67	57	54	53	54	67	42	47
38	9	27	0	25	7	10	22	3	26	0	13
228	171	228	228	228	171	171	171	114	114	57	100
528	228	397	399	342	227	342	342	200	340	125	171
66	59	78	61	64	58	58	52	58	53	46	52
30	28	38	23	29	19	23	21	15	20	10	19
28	28	40	28	28	28	28	25	50	28		
51	50	57	50	50	50	50	50	50	50		
57	57	57	57	57	57	57	57	57	50		
57	57	112	57	57	57	57	57	57	57		
114	114	114	100	114	57	100	57	57	57		
114	114	114	113	114	112	114	100	100	113		
150	171	228	171	171	114	114	142	114	114		
228	171	228	171	228	171	171	171	114	114	57	100
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0

TABLE 5.09.--BISCUITS OTHER THAN REFRIGERATOR TYPE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.5	.8	6.6	7.3	9.3
ON ONLY 1 OF 3 DAYS.....PCT.....	5.1	.6	4.9	5.6	7.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.5	.1	1.3	1.3	1.7
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	.5	.4	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.4	1.5	1.4	1.4
STANDARD DEVIATION.....DO.....	1.0	.8	1.0	.9	.8
MAXIMUM.....DO.....	8.0	3.0	6.0	6.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	35	8	17	21	25
STANDARD DEVIATION.....GM.....	31	7	14	14	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		5	5	9
25TH.....GM.....	19		9	9	13
50TH.....GM.....	25		13	19	19
75TH.....GM.....	42		25	25	28
90TH.....GM.....	75		38	38	49
95TH.....GM.....	93		47	51	63
99TH.....GM.....	168		75	78	89
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	16	42	50	56
STANDARD DEVIATION.....GM.....	53	7	29	27	27
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	73	35	29	42	66
STANDARD DEVIATION.....GM.....	42	0	18	20	29
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	81	0	47	55	48
STANDARD DEVIATION.....GM.....	46	0	14	20	19
MAXIMUM IN ANY 1 DAY.....GM.....	748	56	156	168	192
MAXIMUM OVER 3 DAYS.....GM.....	888	70	224	280	268
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	67	19	35	44	53
STANDARD DEVIATION.....GM.....	43	8	24	25	28
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28		11	14	28
25TH.....GM.....	38		21	28	38
50TH.....GM.....	56		28	38	56
75TH.....GM.....	84		38	56	60
90TH.....GM.....	114		57	84	84
95TH.....GM.....	140		76	84	112
99TH.....GM.....	228		113	114	170
MAXIMUM.....GM.....	448	28	156	152	192
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	18.2	0	0	.8	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	.5	.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
9.3	7.1	8.3	6.9	6.8	6.9	7.9	7.7	10.0	6.8	8.5	5.3
6.3	5.3	5.6	5.1	4.7	5.1	4.8	5.0	6.3	4.5	4.3	2.7
2.3	1.3	1.4	1.0	1.4	1.3	1.9	1.7	1.6	1.2	2.7	1.2
.7	.5	1.3	.7	.7	.6	1.2	1.0	2.1	1.1	1.5	1.4
1.5	1.4	1.6	1.5	1.5	1.4	1.8	1.6	1.9	1.8	2.2	2.0
.7	.7	1.0	.8	.9	.8	1.1	1.0	1.4	1.4	1.3	1.5
6.0	4.0	5.0	5.0	6.0	5.0	6.0	7.0	7.0	8.0	7.0	8.0
33	28	47	29	47	31	49	32	48	32	55	34
23	17	43	24	38	23	42	26	43	28	40	28
9	9	9	9	13	9	9	9	9	9		9
19	19	19	13	25	14	24	14	19	13		14
28	25	37	19	37	25	37	24	28	25		25
42	37	63	40	56	38	63	38	67	35		45
56	56	89	57	93	56	93	65	103	80		56
75	65	131	84	112	79	126	76	114	103		122
129	89	240	114	189	127	196	139	224	149		126
74	67	85	55	104	72	87	62	83	58	84	53
39	35	58	34	95	55	51	40	52	27	32	27
68	59	108	77	96	63	92	60	92	57	122	52
21	23	57	32	43	33	47	33	48	27	65	17
75	61	98	72	94	63	106	72	98	75	81	67
47	13	62	24	40	25	59	36	57	41	32	35
240	224	520	210	748	380	468	380	336	168	308	169
660	266	820	354	748	420	888	504	672	448	532	378
68	61	87	60	93	64	83	58	74	53	75	51
37	29	54	36	58	43	46	34	43	26	38	25
28	28	28	28	28	28	28	28	28	28		28
42	38	56	38	56	38	56	28	56	28		28
56	56	75	56	84	56	76	56	60	56		56
84	76	112	76	112	76	112	76	84	76		60
112	84	168	112	152	113	128	84	120	84		76
140	114	224	120	170	114	168	120	152	114		84
228	168	280	196	336	240	240	168	227	120		168
240	190	284	210	448	380	360	380	240	120	224	168
2.5	.7	0	0	3.8	0	1.5	4.9	0	2.2	0	0
.9	.3	0	0	.9	0	.2	.6	0	1.0	0	0

TABLE 5.10.--REFRIGERATOR-TYPE BISCUITS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.5	.1	3.1	4.6	5.1
ON ONLY 1 OF 3 DAYS.....PCT.....	2.6	.1	2.5	3.5	4.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.6	.9	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	0	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.0	1.2	1.3	1.3
STANDARD DEVIATION.....DO.....	.8	.0	.5	.7	.7
MAXIMUM.....DO.....	7.0	1.0	3.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	24	7	10	14	16
STANDARD DEVIATION.....GM.....	18	0	5	9	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7			7	7
25TH.....GM.....	13			7	7
50TH.....GM.....	20			13	13
75TH.....GM.....	28			19	20
90TH.....GM.....	47			27	27
95TH.....GM.....	60			27	40
99TH.....GM.....	82			40	60
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	52	20	28	33	38
STANDARD DEVIATION.....GM.....	30	0	14	15	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	53	0	20	35	36
STANDARD DEVIATION.....GM.....	26	0	7	13	12
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	59	0	0	35	56
STANDARD DEVIATION.....GM.....	29	0	0	12	6
MAXIMUM IN ANY 1 DAY.....GM.....	340	20	93	80	120
MAXIMUM OVER 3 DAYS.....GM.....	553	20	93	160	180
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	50	20	25	32	36
STANDARD DEVIATION.....GM.....	28	0	13	14	18
PERCENTILES: <u>1/</u>					
5TH.....GM.....	20			20	20
25TH.....GM.....	40			20	20
50TH.....GM.....	40			30	40
75TH.....GM.....	60			40	40
90TH.....GM.....	80			50	60
95TH.....GM.....	100			60	80
99TH.....GM.....	125			80	100
MAXIMUM.....GM.....	340	20	93	80	100
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	4.3	0	0	0	1.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.1	3.8	3.9	3.1	4.3	3.6	3.4	2.9	3.7	3.0	3.3	2.9
3.1	2.9	3.1	2.2	2.9	3.0	2.5	2.0	2.2	1.4	1.0	.9
.9	.7	.6	.8	1.1	.4	.6	.6	.9	1.3	.7	1.4
.1	.1	.3	.1	.2	.1	.3	.3	.6	.3	1.5	.5
1.3	1.3	1.4	1.3	1.4	1.3	1.4	1.5	1.7	1.8	3.2	2.1
.7	.7	.9	.6	.7	.6	.7	.9	1.1	.8	2.0	1.0
4.0	5.0	6.0	4.0	4.0	4.0	4.0	6.0	7.0	3.0	7.0	5.0
24	20	29	20	34	19	29	21	42	26	54	27
18	13	22	13	18	11	17	15	38	19	34	14
7	7	7		13	7	13	7		7		
13	13	13		20	13	13	13		13		
19	19	20		33	13	27	13		19		
27	27	33		47	20	40	27		40		
40	40	67		60	36	57	40		40		
53	42	76		67	47	67	53		53		
80	73	107		80	58	80	80		112		
53	47	63	45	79	47	67	42	65	43	48	65
32	26	38	20	36	23	32	19	27	13	18	32
61	51	74	47	74	46	60	46	75	45	70	39
36	23	38	20	22	19	24	19	31	16	7	17
49	43	75	71	63	48	59	44	111	59	83	40
9	3	18	0	21	8	16	24	44	28	24	17
340	120	227	113	200	140	170	180	220	112	142	128
440	220	320	213	325	173	300	380	553	336	369	200
53	46	63	46	72	45	62	41	74	45	50	40
37	24	37	20	30	22	28	18	33	21	15	25
20	20	20		40	20	20	20		20		
40	20	40		50	20	40	25		40		
40	40	60		60	40	60	40		40		
60	60	80		80	60	80	50		50		
80	80	100		120	80	100	60		60		
100	100	120		120	80	120	80		100		
240	120	227		160	113	170	85		112		
340	120	227	113	200	120	170	113	180	112	85	128
0	0	0	0	0	0	1.5	1.5	0	0	0	0
0	0	0	0	0	0	.7	.5	0	0	0	0

TABLE 5.11.--CORNBREAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.1	2.0	7.9	9.9	10.2
ON ONLY 1 OF 3 DAYS.....PCT.....	6.9	1.5	4.9	6.6	7.5
ON ONLY 2 OF 3 DAYS.....PCT.....	2.4	.2	2.1	2.5	2.3
ON ALL 3 OF 3 DAYS.....PCT.....	.8	.3	.9	.7	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.7	1.6	1.5	1.4
STANDARD DEVIATION.....DO.....	.9	1.4	.8	.8	.7
MAXIMUM.....DO.....	9.0	5.0	4.0	4.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	49	15	28	33	33
STANDARD DEVIATION.....GM.....	51	13	25	29	28
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		3	6	6
25TH.....GM.....	22		11	13	15
50TH.....GM.....	33		22	26	26
75TH.....GM.....	57		37	47	40
90TH.....GM.....	100		63	66	69
95TH.....GM.....	138		83	91	84
99TH.....GM.....	266		102	133	121
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	103	26	56	71	77
STANDARD DEVIATION.....GM.....	87	13	47	63	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	106	61	54	72	67
STANDARD DEVIATION.....GM.....	80	0	25	45	32
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	116	30	63	72	106
STANDARD DEVIATION.....GM.....	94	16	39	33	47
MAXIMUM IN ANY 1 DAY.....GM.....	1,062	78	266	398	531
MAXIMUM OVER 3 DAYS.....GM.....	2,188	144	448	515	681
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	98	25	53	67	73
STANDARD DEVIATION.....GM.....	78	16	39	45	53
PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		8	9	17
25TH.....GM.....	46		31	33	40
50TH.....GM.....	78		40	60	75
75TH.....GM.....	133		78	78	78
90TH.....GM.....	180		100	128	133
95TH.....GM.....	266		120	149	156
99TH.....GM.....	398		199	199	224
MAXIMUM.....GM.....	797	78	266	266	531
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	142.0	0	3.1	6.6	5.9
PROPORTION OF EATING OCCASIONS.....PCT.....	2.4	0	2.3	2.6	2.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.5	10.4	12.4	9.1	7.2	8.7	10.6	11.1	14.8	12.9	14.1	11.2
7.0	7.4	8.8	6.9	5.6	6.0	7.3	7.6	8.8	7.9	8.7	6.3
2.9	2.1	2.7	1.7	1.3	2.4	2.4	2.6	4.0	3.0	3.4	3.0
.6	.9	.8	.4	.3	.3	.9	.9	2.0	2.0	2.0	2.0
1.5	1.5	1.4	1.3	1.3	1.4	1.5	1.5	1.8	1.8	1.8	2.0
.8	.8	.7	.6	.6	.7	.9	.9	1.1	1.0	1.3	1.3
6.0	4.0	4.0	4.0	4.0	4.0	6.0	6.0	6.0	6.0	9.0	6.0
49	43	57	39	56	44	61	48	53	52	80	67
47	39	55	31	45	38	63	53	50	47	109	64
11	7	11	6	13	11	11	11	11	9	11	10
22	19	25	17	26	22	26	19	24	22	27	25
36	27	39	27	44	30	44	28	39	41	52	50
61	52	66	50	77	52	71	52	61	69	103	80
104	89	105	78	104	92	138	94	100	103	160	177
127	104	177	104	133	117	177	133	156	151	177	199
211	216	299	174	183	185	302	266	252	266	730	291
101	89	121	88	133	93	125	93	95	108	174	133
73	68	99	69	82	69	111	79	61	90	162	125
108	106	123	96	140	106	131	101	109	103	114	110
83	76	82	53	106	72	95	80	78	75	37	101
117	93	146	94	112	97	142	128	106	89	183	126
79	45	99	15	42	29	112	100	75	62	240	73
598	531	664	531	598	598	797	1,062	623	531	844	623
1,195	789	930	531	1,195	797	1,793	1,793	936	890	2,188	872
97	90	118	89	127	93	120	93	89	87	137	101
74	66	89	61	86	64	100	74	60	71	112	86
29	21	33	28	33	28	23	25	17	17	39	17
45	45	57	52	78	50	66	45	50	45	72	40
78	78	85	78	117	78	85	75	75	75	90	78
133	112	149	113	156	120	156	120	133	100	160	117
156	149	266	149	234	156	227	156	156	156	299	224
240	224	284	199	266	224	266	266	234	249	398	266
448	312	415	299	448	299	532	300	266	332	531	531
598	531	664	531	598	531	797	797	398	531	531	531
15.7	6.8	5.0	5.4	13.6	12.2	26.4	26.3	3.8	1.4	2.6	7.3
4.7	2.1	2.0	3.0	3.6	1.9	3.3	2.2	1.3	.4	1.9	3.3

TABLE 5.12.--PANCAKES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.3	3.0	12.2	12.7	11.9
ON ONLY 1 OF 3 DAYS.....PCT.....	7.3	2.7	11.0	11.4	10.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	.3	1.1	1.2	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.1	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.3	.3	.3	.4
MAXIMUM.....DO.....	3.0	2.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	43	14	22	28	36
STANDARD DEVIATION.....GM.....	36	12	21	21	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		5	9	9
25TH.....GM.....	18		9	18	18
50TH.....GM.....	33		18	24	27
75TH.....GM.....	50		24	36	49
90TH.....GM.....	74		49	54	73
95TH.....GM.....	108		54	73	74
99TH.....GM.....	195		110	98	146
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	113	36	57	77	96
STANDARD DEVIATION.....GM.....	88	29	47	53	60
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	114	54	67	74	99
STANDARD DEVIATION.....GM.....	76	0	60	44	62
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	119	0	66	74	109
STANDARD DEVIATION.....GM.....	66	0	43	37	25
MAXIMUM IN ANY 1 DAY.....GM.....	940	85	329	294	365
MAXIMUM OVER 3 DAYS.....GM.....	940	108	504	329	504
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	113	39	59	76	96
STANDARD DEVIATION.....GM.....	85	27	50	52	59
PERCENTILES: <u>1/</u>					
5TH.....GM.....	27		14	27	27
25TH.....GM.....	54		27	41	54
50TH.....GM.....	81		50	54	81
75TH.....GM.....	146		73	95	113
90TH.....GM.....	219		146	146	189
95TH.....GM.....	282		146	198	219
99TH.....GM.....	438		284	229	252
MAXIMUM.....GM.....	940	85	329	294	365
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	9.2	0	.7	1.4	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	.5	.6	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
13.5	10.7	9.8	9.8	8.5	7.6	8.0	5.8	5.6	4.9	6.0	4.3
12.0	9.6	8.5	8.4	7.6	6.9	6.4	5.2	4.6	4.7	5.4	4.2
1.2	.9	1.0	1.3	.9	.7	1.3	.5	.8	0	.5	.1
.3	.2	.3	0	(2)	(2)	.3	.1	.2	.2	.1	0
1.2	1.1	1.2	1.1	1.1	1.1	1.2	1.1	1.2	1.1	1.2	1.0
.4	.4	.5	.3	.3	.3	.5	.4	.5	.4	.4	.1
3.0	3.0	3.0	2.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0
45	38	63	46	59	36	55	38	52	33	47	31
31	35	48	40	41	24	45	36	42	20	46	14
18	9	18	18	18	9	18	9	9	12		
25	18	33	18	36	18	27	18	24	18		
36	27	50	33	49	27	37	27	36	27		
54	45	73	49	73	49	72	45	66	42		
91	72	122	97	97	72	108	73	122	57		
108	97	173	146	146	73	153	97	157	73		
146	188	243	210	195	146	219	197	189	97		
123	103	165	114	158	101	139	100	132	98	115	94
80	94	120	96	95	62	105	89	103	60	54	41
103	99	134	144	163	92	125	104	119	0	227	40
38	76	53	78	101	53	77	64	77	0	155	0
127	85	192	0	189	73	139	132	116	39	27	0
46	15	66	0	0	0	65	77	54	3	0	0
438	940	730	752	584	564	584	940	470	292	378	185
616	940	886	752	882	564	752	940	567	292	756	185
118	101	161	121	159	98	134	102	125	92	121	92
72	89	110	93	97	61	96	87	83	60	89	41
54	27	54	49	54	27	37	27	27	27		
73	54	85	54	81	54	81	54	54	54		
100	81	146	81	144	81	108	81	98	81		
146	108	170	146	219	128	162	113	162	108		
200	188	292	219	292	185	219	219	219	162		
282	252	370	292	365	219	296	219	252	219		
438	514	588	564	441	292	565	441	438	292		
438	940	730	752	584	564	584	940	438	292	378	185
0	.9	0	2.5	0	0	2.2	0	0	1.5	0	0
0	.3	0	1.5	0	0	.5	0	0	1.6	0	0

TABLE 5.13.--WAFFLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.9	.6	3.4	5.7	5.9
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	.6	2.4	4.4	4.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.9	1.3	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.1	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.0	1.3	1.3	1.3
STANDARD DEVIATION.....DO.....	.5	.0	.5	.5	.5
MAXIMUM.....DO.....	4.0	1.0	3.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	35	10	24	30	29
STANDARD DEVIATION.....GM.....	30	4	23	26	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7			7	7
25TH.....GM.....	13			13	13
50TH.....GM.....	26			23	26
75TH.....GM.....	47			33	33
90TH.....GM.....	67			67	53
95TH.....GM.....	95			82	83
99TH.....GM.....	158			153	121
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	92	30	53	62	68
STANDARD DEVIATION.....GM.....	82	13	44	36	42
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	80	0	59	80	71
STANDARD DEVIATION.....GM.....	48	0	45	45	46
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	0	66	153	80
STANDARD DEVIATION.....GM.....	36	0	0	0	33
MAXIMUM IN ANY 1 DAY.....GM.....	711	43	200	258	283
MAXIMUM OVER 3 DAYS.....GM.....	711	43	350	458	400
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	87	30	56	69	69
STANDARD DEVIATION.....GM.....	74	13	45	41	45
PERCENTILES: <u>1/</u>					
5TH.....GM.....	20			20	20
25TH.....GM.....	40			39	40
50TH.....GM.....	78			50	50
75TH.....GM.....	100			100	85
90TH.....GM.....	158			113	100
95TH.....GM.....	200			158	158
99TH.....GM.....	400			200	250
MAXIMUM.....GM.....	711	43	200	200	283
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	31.5	0	0	4.2	3.7
PROPORTION OF EATING OCCASIONS.....PCT.....	2.4	0	0	3.3	2.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.2	4.1	3.5	2.4	2.6	2.5	2.2	1.8	2.3	1.6	2.6	1.3
4.2	3.4	2.9	2.3	2.1	2.2	2.0	1.6	2.1	1.0	1.5	1.0
.8	.7	.4	.1	.4	.2	.2	.1	.1	.5	.3	.3
.2	(2)	.1	0	.2	(2)	.1	.1	.1	.1	.8	0
1.2	1.2	1.2	1.0	1.3	1.1	1.2	1.1	1.2	1.4	1.7	1.3
.5	.4	.5	.2	.6	.4	.4	.4	.5	.6	.9	.4
3.0	3.0	3.0	2.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0
36	32	50	27	48	35	43	29	39	28	41	42
28	29	29	18	38	36	33	22	30	16	21	37
10	7			13	7	13	7				
17	13			17	17	20	13				
27	23			33	26	33	26				
50	33			67	33	52	33				
75	75			100	67	80	57				
94	95			117	94	105	76				
133	133			237	189	189	100				
88	75	134	83	129	105	122	81	99	59	98	108
68	74	77	56	121	111	99	56	56	22	49	55
81	92	115	39	98	49	83	80	200	67	35	91
36	44	48	0	36	32	49	48	0	23	0	97
100	59	84	0	65	79	59	52	20	40	64	0
44	0	18	0	21	0	32	42	0	0	14	0
400	494	340	284	711	566	566	267	234	100	158	316
524	494	400	284	711	566	566	316	400	200	234	474
87	80	125	79	112	95	110	78	101	61	71	101
62	68	70	55	100	102	91	55	65	23	37	80
20	20			40	20	39	20				
50	40			50	40	50	40				
78	57			80	78	85	78				
100	100			150	100	150	100				
150	158			200	200	200	158				
200	200			300	235	300	170				
400	400			711	566	566	267				
400	494	340	284	711	566	566	267	234	100	158	316
4.3	2.1	2.0	0	11.6	0	3.6	0	0	0	0	0
3.2	2.0	3.5	0	8.8	0	2.8	0	0	0	0	0

TABLE 5.14.--TORTILLAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.9	.8	3.9	5.1	4.7
ON ONLY 1 OF 3 DAYS.....PCT.....	1.6	.8	2.2	2.4	2.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	0	.3	.8	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.8	0	1.4	1.9	1.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.4	1.0	2.5	2.9	2.3
STANDARD DEVIATION.....DO.....	2.0	.0	1.8	2.3	2.0
MAXIMUM.....DO.....	9.0	1.0	7.0	9.0	9.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	55	5	21	35	43
STANDARD DEVIATION.....GM.....	62	2	17	31	54
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	10			6	10
25TH.....GM.....	20			10	10
50TH.....GM.....	30			25	23
75TH.....GM.....	63			53	47
90TH.....GM.....	130			76	100
95TH.....GM.....	180			95	110
99TH.....GM.....	284			180	280
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	68	16	30	42	50
STANDARD DEVIATION.....GM.....	46	7	13	21	26
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	76	0	41	50	50
STANDARD DEVIATION.....GM.....	40	0	17	22	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	121	0	37	62	108
STANDARD DEVIATION.....GM.....	79	0	16	33	72
MAXIMUM IN ANY 1 DAY.....GM.....	630	28	70	180	350
MAXIMUM OVER 3 DAYS.....GM.....	1,680	28	175	540	840
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	69	16	26	36	55
STANDARD DEVIATION.....GM.....	39	7	11	16	29
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28			15	28
25TH.....GM.....	30			30	30
50TH.....GM.....	60			30	57
75TH.....GM.....	90			35	70
90TH.....GM.....	120			60	90
95TH.....GM.....	140			60	105
99TH.....GM.....	210			90	140
MAXIMUM.....GM.....	227	28	60	90	140
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	10.8	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.0	4.3	3.4	4.0	3.2	3.4	2.2	2.3	.4	.7	1.7	.3
2.2	2.2	2.1	2.3	1.6	1.8	1.2	1.4	.3	.2	.7	.1
.7	.8	.6	.9	.7	.6	.3	.3	0	.4	.3	.2
1.1	1.3	.6	.9	.9	1.0	.7	.6	.2	.1	.8	0
2.2	2.3	1.9	2.1	2.5	2.4	2.3	2.5	2.9	2.3	2.5	2.0
1.6	1.9	1.6	1.5	2.1	1.9	1.8	2.2	2.4	1.5	1.6	.0
6.0	9.0	9.0	9.0	9.0	9.0	8.0	9.0	6.0	6.0	5.0	2.0
54	50	63	48	81	54	76	52	53	56	53	47
42	52	60	42	82	55	92	58	29	53	32	4
10	10		10	10	10	10	10				
20	20		20	23	20	20	19				
40	30		38	47	30	47	30				
80	60		60	133	60	93	67				
117	110		82	240	120	150	132				
140	170		120	260	180	268	150				
190	292		240	330	280	560	315				
82	67	91	82	83	67	82	67	90	41	60	120
46	44	43	63	50	38	57	43	0	14	0	0
87	67	123	77	96	64	88	72	0	77	85	75
39	29	61	28	44	31	38	43	0	24	0	0
107	103	158	99	193	119	165	128	90	195	82	0
38	65	67	56	75	63	118	66	0	0	20	0
280	315	420	245	420	360	630	385	150	195	120	120
570	875	840	720	990	840	1,680	945	270	585	300	150
74	66	100	69	97	67	97	64	54	72	63	71
31	33	48	33	42	31	51	31	35	23	8	14
30	30		30	30	30	30	30				
60	35		35	60	35	60	35				
60	60		60	105	60	90	60				
90	75		90	120	90	120	90				
120	105		120	142	105	180	105				
128	140		120	180	120	210	113				
140	210		140	180	180	210	175				
180	227	210	150	227	227	210	210	120	105	85	90
0	.6	2.2	.7	2.4	.9	1.7	2.3	0	0	0	0
0	.3	2.5	.5	.8	.2	.7	.6	0	0	0	0

TABLE 5.15.--CAKES AND CUPCAKES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	25.5	1.6	17.4	25.3	34.4
ON ONLY 1 OF 3 DAYS.....PCT.....	17.3	1.1	14.7	19.1	22.3
ON ONLY 2 OF 3 DAYS.....PCT.....	6.2	.5	2.0	4.8	9.6
ON ALL 3 OF 3 DAYS.....PCT.....	1.9	0	.6	1.5	2.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.3	1.3	1.5	1.6
STANDARD DEVIATION.....DO.....	.9	.5	.6	.9	.9
MAXIMUM.....DO.....	8.0	2.0	4.0	6.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	41	23	22	30	35
STANDARD DEVIATION.....GM.....	38	16	20	26	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		3	7	9
25TH.....GM.....	16		9	13	16
50TH.....GM.....	31		15	21	28
75TH.....GM.....	52		30	38	42
90TH.....GM.....	83		48	60	65
95TH.....GM.....	112		57	77	88
99TH.....GM.....	192		114	137	128
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	84	55	55	66	73
STANDARD DEVIATION.....GM.....	64	44	42	51	50
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	90	51	57	74	74
STANDARD DEVIATION.....GM.....	63	22	38	53	38
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	95	0	65	63	69
STANDARD DEVIATION.....GM.....	57	0	38	33	40
MAXIMUM IN ANY 1 DAY.....GM.....	1,382	142	259	411	460
MAXIMUM OVER 3 DAYS.....GM.....	1,460	142	405	570	610
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	79	53	51	61	66
STANDARD DEVIATION.....GM.....	59	37	38	45	42
PERCENTILES: <u>1/</u>					
5TH.....GM.....	23		9	19	25
25TH.....GM.....	41		28	35	38
50TH.....GM.....	63		41	46	52
75TH.....GM.....	99		65	78	88
90TH.....GM.....	144		100	113	113
95TH.....GM.....	184		138	147	139
99TH.....GM.....	284		179	228	227
MAXIMUM.....GM.....	1,382	142	259	358	358
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2357.2	2.6	28.0	104.8	121.9
PROPORTION OF EATING OCCASIONS.....PCT.....	15.8	24.7	11.9	16.6	12.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
36.4	35.2	31.0	26.5	24.5	20.2	27.2	22.8	24.8	22.7	22.4	25.8
23.5	24.5	20.7	19.5	15.5	15.0	17.2	16.2	16.3	14.5	14.5	16.1
9.9	8.8	8.7	5.7	6.4	4.2	6.9	5.2	5.8	6.0	5.5	8.1
3.1	1.9	1.7	1.3	2.5	1.0	3.1	1.4	2.7	2.2	2.4	1.7
1.6	1.5	1.6	1.5	1.7	1.4	1.7	1.5	1.7	1.6	1.8	1.5
.9	.8	.8	.8	1.0	.7	1.0	.8	1.0	.9	1.1	.8
6.0	7.0	6.0	8.0	6.0	5.0	6.0	7.0	6.0	7.0	6.0	5.0
43	39	48	39	53	37	49	35	43	39	53	35
36	33	45	37	48	30	44	31	39	31	55	47
9	9	11	9	12	8	10	7	8	7	11	6
19	16	19	18	23	15	19	14	19	16	19	15
32	30	33	30	38	30	36	27	34	31	35	27
55	50	62	49	65	48	62	44	53	55	76	42
86	76	95	76	108	76	108	71	91	79	118	61
116	104	118	96	143	93	140	90	113	95	147	77
189	157	222	155	244	139	217	170	203	160	339	131
87	81	103	88	101	83	94	75	86	81	94	67
62	58	93	61	73	62	76	54	53	68	69	45
90	89	105	94	121	83	101	77	80	77	96	84
58	48	61	84	87	44	63	50	45	42	43	111
93	95	103	81	101	93	110	91	107	82	164	61
53	61	97	55	56	43	59	46	69	37	89	28
546	438	875	677	743	372	796	560	737	496	498	1,382
823	944	1,460	1,340	1,080	710	1,040	960	907	709	1,017	1,427
80	77	93	80	96	78	89	71	77	73	92	68
56	55	71	59	67	52	62	50	53	52	57	105
25	26	28	24	29	22	28	20	22	22	29	19
41	39	43	42	48	41	44	37	40	38	45	38
66	57	72	64	85	66	80	57	58	57	84	51
96	96	113	101	113	100	113	92	105	95	114	79
158	140	179	143	174	139	170	137	147	125	172	106
186	179	230	185	227	179	200	173	179	173	180	132
284	302	346	302	340	269	306	257	240	227	319	232
410	410	614	648	626	355	614	480	369	496	319	1,382
157.4	155.1	77.0	92.8	231.2	301.6	392.1	409.8	83.1	123.1	24.0	52.7
12.8	13.5	11.5	16.2	14.5	19.8	17.6	17.3	18.0	19.5	11.4	13.3

TABLE 5.16.--CHOCOLATE CAKES AND CUPCAKES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.8	.3	5.8	10.5	16.0
ON ONLY 1 OF 3 DAYS.....PCT.....	6.7	0	5.2	8.8	12.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.7	.3	.5	1.5	3.4
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.2	.2	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.4	2.0	1.3	1.3	1.4
STANDARD DEVIATION.....DO.....	.7	.0	.6	.8	.7
MAXIMUM.....DO.....	6.0	2.0	3.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	36	46	22	26	29
STANDARD DEVIATION.....GM.....	31	0	16	22	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	9		8	8	9
25TH.....GM.....	15		12	13	15
50TH.....GM.....	30		15	19	24
75TH.....GM.....	46		29	31	35
90TH.....GM.....	73		46	53	57
95TH.....GM.....	92		57	59	62
99TH.....GM.....	170		85	92	94
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	84	0	58	62	67
STANDARD DEVIATION.....GM.....	61	0	40	37	39
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	89	69	46	82	70
STANDARD DEVIATION.....GM.....	59	0	17	60	38
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	85	0	67	34	55
STANDARD DEVIATION.....GM.....	52	0	17	17	29
MAXIMUM IN ANY 1 DAY.....GM.....	737	92	184	284	460
MAXIMUM OVER 3 DAYS.....GM.....	884	138	255	567	495
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	79	69	52	59	62
STANDARD DEVIATION.....GM.....	53	23	31	41	35
PERCENTILES: ^{1/}					
5TH.....GM.....	28		23	19	28
25TH.....GM.....	41		35	35	38
50TH.....GM.....	70		41	46	46
75TH.....GM.....	92		57	70	85
90TH.....GM.....	142		92	104	92
95TH.....GM.....	177		118	113	133
99TH.....GM.....	278		171	284	184
MAXIMUM.....GM.....	626	92	171	284	244
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	543.8	1.4	5.1	37.2	41.5
PROPORTION OF EATING OCCASIONS.....PCT.....	11.8	50.0	6.6	15.6	10.2

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.6	13.9	12.4	9.6	9.2	7.3	9.2	6.5	5.1	4.1	6.0	4.9
11.0	11.1	9.3	7.8	6.4	5.8	6.5	5.2	3.8	2.6	5.2	4.0
2.9	2.3	2.9	1.6	2.2	1.4	2.0	1.1	1.2	1.4	.6	.9
.7	.5	.2	.2	.6	.1	.7	.2	.1	.1	.2	.1
1.4	1.4	1.4	1.3	1.5	1.3	1.5	1.3	1.4	1.5	1.4	1.3
.8	.7	.7	.7	.8	.6	.8	.7	.7	.7	.7	.5
6.0	5.0	4.0	4.0	5.0	4.0	4.0	6.0	3.0	4.0	3.0	3.0
36	32	46	37	51	32	44	31	49	42	41	27
26	25	38	33	46	26	35	22	39	30	33	19
11	9	12	11	13	9	12	7	19	6		
16	15	23	19	24	15	19	15	31	17		
30	27	31	28	36	28	31	27	38	35		
47	44	61	46	61	38	58	38	52	61		
67	61	90	62	93	61	87	61	92	78		
88	76	113	89	139	92	96	77	123	94		
150	124	220	141	244	139	207	104	246	189		
90	77	108	86	106	77	96	76	134	88	106	68
59	51	97	51	75	48	64	50	120	55	93	51
72	82	112	109	121	86	95	73	96	87	109	60
38	45	61	87	87	60	51	38	52	34	31	18
76	70	70	67	110	72	98	76	46	141	81	81
49	48	16	28	60	29	56	28	0	47	0	0
371	371	726	663	696	355	454	348	737	227	354	232
547	650	726	884	812	710	650	487	737	567	354	244
75	71	95	83	103	76	90	70	103	85	87	64
47	47	64	49	68	50	54	45	66	47	64	41
28	25	28	30	35	23	35	20	41	23		
39	38	46	46	57	41	46	37	57	46		
62	57	84	70	91	64	85	57	92	91		
92	92	114	92	133	92	112	92	122	108		
138	133	184	162	184	138	162	128	162	157		
168	162	227	186	227	157	184	157	183	177		
266	278	340	221	278	278	278	227	369	186		
371	371	369	313	626	355	454	348	369	227	313	232
37.7	35.0	29.3	26.4	67.6	98.9	77.9	45.0	7.2	20.9	2.0	10.7
8.6	8.6	11.8	14.0	12.6	19.8	11.8	7.4	8.8	19.7	4.5	17.3

TABLE 5.17.--COOKIES

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	30.8	11.9	46.3	48.1	53.2
ON ONLY 1 OF 3 DAYS.....PCT.....	17.4	4.7	22.5	23.7	27.1
ON ONLY 2 OF 3 DAYS.....PCT.....	9.0	5.3	13.9	16.0	17.5
ON ALL 3 OF 3 DAYS.....PCT.....	4.4	1.9	9.9	8.4	8.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.9	2.0	2.1	2.0	2.0
STANDARD DEVIATION.....DO.....	1.2	1.1	1.3	1.2	1.3
MAXIMUM.....DO.....	12.0	6.0	8.0	8.0	9.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	20	10	14	17	19
STANDARD DEVIATION.....GM.....	21	9	13	16	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3	2	2	3	3
25TH.....GM.....	7	4	7	7	7
50TH.....GM.....	14	7	11	12	13
75TH.....GM.....	25	13	18	23	25
90TH.....GM.....	42	26	28	33	38
95TH.....GM.....	57	30	38	47	53
99TH.....GM.....	103	36	66	76	82
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	36	15	23	32	31
STANDARD DEVIATION.....GM.....	37	13	17	33	26
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	38	21	26	32	33
STANDARD DEVIATION.....GM.....	31	15	15	24	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	40	12	26	29	39
STANDARD DEVIATION.....GM.....	29	7	18	17	26
MAXIMUM IN ANY 1 DAY.....GM.....	648	71	165	432	369
MAXIMUM OVER 3 DAYS.....GM.....	1,358	108	312	474	759
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	32	15	21	25	28
STANDARD DEVIATION.....GM.....	30	13	15	22	21
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7	7	5	7	7
25TH.....GM.....	14	7	10	14	14
50TH.....GM.....	26	10	16	20	24
75TH.....GM.....	40	14	26	30	36
90TH.....GM.....	60	35	39	48	56
95TH.....GM.....	84	42	57	57	60
99TH.....GM.....	144	57	64	106	112
MAXIMUM.....GM.....	648	71	165	432	216
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	79.1	1.5	3.6	10.0	5.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	1.2	.4	.6	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
44.4	43.1	37.9	34.9	24.5	24.4	27.3	23.8	27.6	22.6	29.6	27.4
23.7	23.8	20.9	21.4	14.5	15.5	14.8	15.4	13.3	14.0	15.8	12.3
14.5	13.2	11.8	9.2	6.8	6.3	7.7	6.4	8.1	5.5	7.6	8.4
6.1	6.1	5.2	4.3	3.2	2.5	4.7	2.0	6.2	3.1	6.2	6.7
1.9	1.9	1.9	1.8	1.8	1.7	1.9	1.6	2.1	1.7	2.0	2.1
1.2	1.2	1.2	1.1	1.1	1.1	1.2	1.0	1.3	1.1	1.5	1.3
8.0	8.0	7.0	6.0	9.0	8.0	11.0	8.0	8.0	7.0	12.0	6.0
23	20	29	19	27	18	24	15	21	16	21	15
27	19	29	18	26	21	22	14	18	19	26	13
3	3	4	3	3	2	3	2	3	3	3	3
8	8	9	7	9	7	9	6	9	7	7	6
15	15	20	14	19	12	17	11	16	11	14	11
30	27	37	24	36	22	32	19	27	19	27	19
47	43	59	39	57	38	52	30	45	32	37	30
63	57	81	54	82	56	67	40	58	45	64	47
113	88	156	80	113	104	115	69	86	73	196	59
37	36	58	37	53	36	41	30	34	32	35	23
36	32	74	32	52	33	41	25	27	28	30	17
44	40	49	35	53	40	45	31	36	25	39	25
34	33	36	28	46	43	28	19	25	11	27	18
49	39	57	40	53	37	47	33	37	41	41	27
48	21	36	24	26	26	28	23	21	36	44	14
624	336	648	317	540	621	395	255	228	364	227	113
1,358	672	693	439	756	684	427	454	318	510	588	227
36	32	45	31	46	33	38	27	30	28	32	21
36	29	50	26	41	33	29	21	20	28	27	14
7	7	7	7	9	7	8	7	9	7	7	7
16	14	21	14	21	14	18	14	16	14	16	13
27	26	32	25	36	25	30	21	27	22	26	16
42	40	56	41	56	40	48	32	39	32	36	28
64	60	84	60	85	64	72	53	50	48	54	36
85	81	113	81	113	85	94	63	57	57	81	48
168	165	227	113	227	170	142	113	136	144	170	78
624	336	648	240	384	432	284	192	170	338	192	113
5.9	9.8	3.0	4.7	11.5	0	11.8	6.8	0	3.5	0	2.1
.3	.6	.3	.5	.7	0	.5	.2	0	.5	0	.4

TABLE 5.18.--CHOCOLATE CHIP COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.2	3.1	16.1	17.6	16.3
ON ONLY 1 OF 3 DAYS.....PCT.....	5.7	2.3	10.8	11.1	11.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.8	.5	3.7	5.3	3.7
ON ALL 3 OF 3 DAYS.....PCT.....	.6	.3	1.5	1.2	1.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.5	1.7	1.7	1.6
STANDARD DEVIATION.....DO.....	1.0	1.1	1.2	1.0	.9
MAXIMUM.....DO.....	12.0	5.0	8.0	8.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	12	11	9	12	12
STANDARD DEVIATION.....GM.....	13	6	8	11	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		2	2	2
25TH.....GM.....	5		5	5	5
50TH.....GM.....	9		7	9	9
75TH.....GM.....	14		12	14	14
90TH.....GM.....	28		19	23	26
95TH.....GM.....	37		24	28	33
99TH.....GM.....	61		45	57	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	26	30	17	23	24
STANDARD DEVIATION.....GM.....	25	17	11	19	22
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	15	22	26	26
STANDARD DEVIATION.....GM.....	24	1	11	20	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	26	17	23	23	25
STANDARD DEVIATION.....GM.....	18	9	13	13	10
MAXIMUM IN ANY 1 DAY.....GM.....	340	57	113	180	170
MAXIMUM OVER 3 DAYS.....GM.....	567	77	198	294	227
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	24	21	17	21	22
STANDARD DEVIATION.....GM.....	20	15	11	14	16
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		7	7	7
25TH.....GM.....	14		10	14	14
50TH.....GM.....	21		14	14	15
75TH.....GM.....	28		21	28	28
90TH.....GM.....	42		28	35	35
95TH.....GM.....	57		30	44	48
99TH.....GM.....	113		57	85	105
MAXIMUM.....GM.....	227	57	85	113	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	10.4	.7	0	1.2	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	2.7	0	.2	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.0	12.3	12.7	11.6	6.4	6.4	6.2	5.0	3.6	3.0	3.5	3.7
9.5	8.5	8.8	8.3	4.1	4.7	4.4	3.9	2.1	2.1	2.4	1.9
3.6	2.7	3.0	2.6	1.6	1.2	1.1	1.0	.6	.9	.5	.9
.9	1.2	.9	.6	.7	.4	.7	.1	.9	0	.6	.9
1.6	1.6	1.6	1.4	1.7	1.5	1.5	1.3	1.7	1.4	2.4	1.9
.9	1.0	1.0	.7	.9	1.0	.9	.6	1.0	.7	2.8	1.0
7.0	7.0	6.0	5.0	6.0	7.0	6.0	4.0	4.0	4.0	12.0	4.0
15	13	16	11	18	13	13	9	13	7	16	8
16	12	16	11	15	16	11	9	12	4	25	4
2	2	2	2	3	2	3	2		2		
5	5	7	5	7	5	5	5		5		
9	9	12	9	14	8	9	7		7		
19	14	19	12	28	14	15	9		9		
28	28	35	19	37	28	28	16		12		
47	38	43	23	44	38	38	22		14		
76	70	57	57	66	76	57	38		16		
30	26	38	23	35	29	26	21	22	17	26	14
36	23	35	14	27	31	20	17	9	8	24	10
35	30	30	27	40	32	34	21	25	16	25	12
24	24	15	28	19	41	18	22	8	5	10	2
28	25	38	18	37	24	25	22	24	0	48	14
24	15	34	8	17	13	14	12	19	0	47	(2)
340	170	255	198	227	340	170	255	98	28	126	49
397	227	468	340	290	567	175	354	182	49	308	49
27	24	30	22	33	26	26	20	23	15	21	12
24	19	25	21	20	28	16	16	11	6	12	4
7	7	7	7	10	7	7	7		7		
14	14	14	14	21	14	14	14		14		
21	21	22	21	28	21	21	14		14		
28	28	35	28	42	28	33	21		14		
42	42	57	35	57	44	42	35		28		
71	56	84	42	84	66	56	44		28		
125	113	168	170	105	170	85	85		28		
170	170	170	198	113	227	126	170	49	28	42	28
0	4.0	1.9	1.0	0	0	1.6	0	0	0	0	0
0	.9	.7	.4	0	0	.3	0	0	0	0	0

TABLE 5.19.--OATMEAL COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.5	.9	4.6	4.3	6.7
ON ONLY 1 OF 3 DAYS.....PCT.....	2.5	.6	3.4	3.5	4.9
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	.3	1.0	.5	1.3
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.2	.3	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	1.7	1.5	1.4	1.5
STANDARD DEVIATION.....DO.....	1.0	.9	1.1	.9	.9
MAXIMUM.....DO.....	7.0	3.0	7.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	18	9	14	13	18
STANDARD DEVIATION.....GM.....	18	4	14	8	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4			4	4
25TH.....GM.....	9			9	9
50TH.....GM.....	13			9	13
75TH.....GM.....	22			17	22
90TH.....GM.....	39			26	39
95TH.....GM.....	52			30	52
99TH.....GM.....	91			43	82
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	38	19	26	31	36
STANDARD DEVIATION.....GM.....	29	6	12	15	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	19	40	35	44
STANDARD DEVIATION.....GM.....	31	0	25	11	26
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	0	54	32	44
STANDARD DEVIATION.....GM.....	28	0	15	7	25
MAXIMUM IN ANY 1 DAY.....GM.....	364	26	165	91	170
MAXIMUM OVER 3 DAYS.....GM.....	429	39	196	130	281
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	36	15	28	27	34
STANDARD DEVIATION.....GM.....	26	5	18	12	20
PERCENTILES: <u>1/</u>					
5TH.....GM.....	13			13	13
25TH.....GM.....	23			13	26
50TH.....GM.....	26			26	26
75TH.....GM.....	39			39	39
90TH.....GM.....	65			39	57
95TH.....GM.....	78			52	65
99TH.....GM.....	130			57	113
MAXIMUM.....GM.....	338	26	165	65	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	7.2	0	.9	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	1.2	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.0	4.1	4.3	3.8	3.0	2.7	3.4	2.5	4.4	2.5	5.1	3.9
3.8	3.2	3.6	3.4	2.3	1.9	2.1	1.8	2.7	1.9	3.0	2.2
1.1	.6	.7	.3	.6	.7	.9	.7	1.3	.3	1.6	.9
.2	.3	.1	.2	.1	.2	.5	.1	.5	.2	.5	.7
1.5	1.4	1.4	1.2	1.5	1.6	1.8	1.5	1.7	1.5	1.9	2.0
.9	.8	.8	.6	1.1	1.2	1.0	.7	.9	1.0	1.3	1.4
5.0	4.0	4.0	3.0	7.0	7.0	5.0	5.0	4.0	5.0	5.0	6.0
19	16	22	14	23	20	25	14	16	17	18	19
19	11	20	11	18	21	23	10	9	29	11	17
4	4	4	4	6	4	4	4				
9	9	9	4	9	9	9	9				
13	13	17	13	17	13	17	9				
22	22	26	17	30	27	35	17				
35	28	43	35	48	48	48	28				
52	35	82	43	61	56	78	38				
112	52	86	48	91	113	113	52				
38	37	52	37	51	37	48	31	32	26	38	35
30	27	44	30	35	29	40	18	18	13	15	20
58	37	59	53	53	52	48	33	35	38	43	35
43	18	39	20	24	43	33	19	8	2	20	37
28	30	61	15	63	50	53	32	27	80	13	36
4	6	10	2	31	11	31	7	1	65	0	15
260	195	247	143	161	170	234	113	71	364	78	113
368	195	257	143	273	340	397	156	91	429	130	227
38	33	47	34	45	37	43	30	28	35	29	28
29	23	28	23	28	30	28	17	10	56	13	21
13	13	13	13	13	13	13	13				
26	23	26	13	26	13	26	13				
26	26	39	26	39	26	39	26				
46	39	65	52	52	46	52	39				
78	65	78	65	78	78	78	52				
85	78	113	78	92	113	104	69				
184	104	127	130	156	170	156	78				
260	195	130	130	161	170	227	113	65	338	52	113
.8	0	0	0	0	0	0	3.0	0	2.6	0	0
.5	0	0	0	0	0	0	1.1	0	4.0	0	0

TABLE 5.20.--PEANUT BUTTER COOKIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.4	.7	2.4	2.9	5.7
ON ONLY 1 OF 3 DAYS.....PCT.....	1.9	.7	1.6	2.2	4.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.6	.6	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.1	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.0	1.6	1.5	1.4
STANDARD DEVIATION.....DO.....	.8	.0	.9	.9	1.0
MAXIMUM.....DO.....	7.0	1.0	5.0	6.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	15	6	13	14	12
STANDARD DEVIATION.....GM.....	15	2	7	9	13
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4			4	4
25TH.....GM.....	6			8	4
50TH.....GM.....	9			12	8
75TH.....GM.....	17			19	15
90TH.....GM.....	28			24	20
95TH.....GM.....	40			28	46
99TH.....GM.....	72			47	53
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	33	19	34	35	24
STANDARD DEVIATION.....GM.....	28	6	21	19	16
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	35	0	23	30	35
STANDARD DEVIATION.....GM.....	31	0	7	18	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	39	0	22	35	41
STANDARD DEVIATION.....GM.....	20	0	0	8	18
MAXIMUM IN ANY 1 DAY.....GM.....	624	24	81	85	113
MAXIMUM OVER 3 DAYS.....GM.....	636	24	81	142	216
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	31	19	24	29	26
STANDARD DEVIATION.....GM.....	29	6	11	16	17
PERCENTILES: <u>1/</u>					
5TH.....GM.....	12			12	12
25TH.....GM.....	12			12	12
50TH.....GM.....	24			24	24
75TH.....GM.....	36			36	36
90TH.....GM.....	57			52	48
95TH.....GM.....	72			60	52
99TH.....GM.....	113			85	113
MAXIMUM.....GM.....	624	24	60	85	113
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	0	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	0	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.4	5.0	3.7	3.8	1.9	1.8	1.4	1.4	2.1	1.4	.3	1.3
4.5	4.5	2.8	3.2	1.5	1.3	1.0	1.1	1.7	.7	.3	.5
.8	.5	.6	.5	.3	.4	.2	.3	.4	.5	0	.7
.1	0	.3	(2)	.1	.1	.1	.1	0	.3	0	.2
1.2	1.2	1.5	1.2	1.4	1.6	1.4	1.5	1.4	2.2	1.0	2.0
.6	.4	.9	.5	.6	1.0	.7	.9	.7	1.8	.0	.7
4.0	3.0	5.0	3.0	3.0	5.0	4.0	6.0	3.0	7.0	1.0	3.0
13	11	19	12	22	16	20	13	15	17	8	12
21	9	18	9	18	14	17	9	11	21	0	7
4	4	4	4	4	4	4	4				
4	4	8	4	8	4	8	8				
8	8	12	8	16	12	12	8				
15	14	24	16	29	24	20	16				
24	20	51	24	36	38	52	24				
24	33	63	33	64	44	64	38				
48	48	83	44	89	80	66	56				
27	29	42	31	55	37	40	32	32	17	24	30
17	23	38	24	53	32	29	22	15	6	0	10
49	35	44	32	50	37	37	26	51	21	0	13
71	18	27	15	19	28	20	14	15	7	0	2
35	0	54	24	41	26	62	24	0	52	0	28
7	0	22	0	10	12	6	15	0	24	0	0
624	170	227	106	268	168	156	113	72	120	24	36
636	170	248	132	268	240	197	168	120	216	24	85
32	28	40	30	48	30	41	26	32	24	24	18
53	20	32	21	42	23	28	16	14	11	0	9
12	12	12	12	12	12	12	12				
12	12	24	12	24	12	24	12				
24	24	28	24	36	24	36	24				
36	36	48	36	50	36	48	36				
48	56	72	60	72	72	72	57				
72	72	104	72	96	84	104	57				
96	85	156	106	268	113	156	72				
624	144	227	106	268	113	156	85	57	48	24	36
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0

TABLE 5.21.--SUGAR COOKIES

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.2	.9	3.6	4.4	5.3
ON ONLY 1 OF 3 DAYS.....PCT.....	2.3	.7	2.6	3.5	4.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.6	.7	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.2	.2	.4	.3	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	1.5	1.4	1.6	1.4	1.4
STANDARD DEVIATION.....DO.....	.8	.8	1.1	.8	.9
MAXIMUM.....DO.....	6.0	3.0	5.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	18	12	15	17	16
STANDARD DEVIATION.....GM.....	16	8	16	17	13
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	5			5	5
25TH.....GM.....	8			9	5
50TH.....GM.....	11			11	11
75TH.....GM.....	21			21	21
90TH.....GM.....	38			43	32
95TH.....GM.....	53			57	48
99TH.....GM.....	75			87	64
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	40	27	28	43	34
STANDARD DEVIATION.....GM.....	34	13	13	42	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	41	0	27	28	41
STANDARD DEVIATION.....GM.....	24	0	15	14	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	42	27	49	52	48
STANDARD DEVIATION.....GM.....	22	0	25	20	9
MAXIMUM IN ANY 1 DAY.....GM.....	320	48	113	224	170
MAXIMUM OVER 3 DAYS.....GM.....	340	80	227	260	192
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	37	27	28	37	34
STANDARD DEVIATION.....GM.....	27	10	14	26	21
PERCENTILES: ^{1/}					
5TH.....GM.....	8			8	16
25TH.....GM.....	16			16	16
50TH.....GM.....	32			32	32
75TH.....GM.....	48			48	48
90TH.....GM.....	64			64	57
95TH.....GM.....	81			96	64
99TH.....GM.....	142			128	85
MAXIMUM.....GM.....	320	48	64	142	170
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	9.1	0	1.1	0	1.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	1.8	0	.8

^{1/} SEE TABLE NOTES IN APPENDIX.^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.9	5.0	3.1	3.1	1.3	2.0	2.9	3.0	4.1	3.3	6.4	5.6
3.3	3.9	2.3	2.2	.9	1.6	1.9	2.3	2.3	2.3	4.0	3.0
.6	.9	.6	.7	.4	.3	.8	.6	.9	.8	1.4	1.5
.1	.2	.2	.1	(2)	.1	.1	.1	.9	.2	1.0	1.2
1.2	1.4	1.4	1.4	1.5	1.4	1.5	1.4	1.9	1.4	1.7	1.9
.5	.8	.8	.7	.7	.7	.7	.7	1.3	.6	1.0	1.2
4.0	5.0	5.0	3.0	4.0	4.0	4.0	4.0	5.0	3.0	4.0	6.0
15	18	25	18	27	16	24	15	20	13	22	16
13	16	21	17	19	12	18	13	18	8	20	13
3	3			5	5	4	5		5		5
5	5			11	9	11	5		9		11
11	11			21	11	21	11		11		11
19	21			43	21	32	16		18		18
32	43			53	36	47	27		27		32
47	54			64	48	54	38		37		59
59	64			95	48	113	69		37		59
39	36	64	34	71	40	53	35	27	32	35	32
34	26	61	23	62	28	36	27	14	22	22	15
36	58	41	53	51	38	46	38	39	28	64	20
21	26	19	39	16	18	21	27	11	9	39	9
37	50	60	29	45	38	73	29	40	23	34	33
0	11	7	3	0	13	26	7	24	10	13	17
162	144	320	240	284	144	142	162	112	112	135	96
176	255	320	272	284	144	340	227	256	112	216	176
37	39	53	37	55	36	48	32	31	29	38	25
29	25	47	32	44	23	29	24	15	13	21	12
8	8			16	13	8	8		16		8
16	16			27	16	28	16		16		16
32	32			48	32	48	28		28		24
48	54			64	48	64	32		32		32
64	64			128	64	96	64		48		32
81	81			128	85	108	80		54		48
162	142			284	113	135	144		64		64
162	144	320	240	284	113	142	162	64	64	81	64
0	.8	0	0	0	0	1.7	2.1	0	0	0	2.1
0	.5	0	0	0	0	.8	.7	0	0	0	2.0

TABLE 5.22.--PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	11.9	.5	4.7	7.1	8.1
ON ONLY 1 OF 3 DAYS.....PCT.....	9.3	.5	4.0	5.9	7.1
ON ONLY 2 OF 3 DAYS.....PCT.....	2.1	0	.7	1.2	1.0
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	0	(2)	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.2	1.3	1.3	1.2
STANDARD DEVIATION.....DO.....	.6	.4	.6	.5	.4
MAXIMUM.....DO.....	5.0	2.0	4.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	57	22	37	44	45
STANDARD DEVIATION.....GM.....	38	15	24	24	31
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	19			15	11
25TH.....GM.....	34			28	25
50TH.....GM.....	45			39	39
75TH.....GM.....	70			56	50
90TH.....GM.....	102			79	77
95TH.....GM.....	135			80	110
99TH.....GM.....	197			133	180
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	136	65	103	117	113
STANDARD DEVIATION.....GM.....	70	46	56	57	58
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	144	0	84	102	134
STANDARD DEVIATION.....GM.....	68	0	57	40	73
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	134	0	0	133	133
STANDARD DEVIATION.....GM.....	51	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	960	140	228	275	450
MAXIMUM OVER 3 DAYS.....GM.....	1,283	140	405	480	675
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	129	53	88	106	116
STANDARD DEVIATION.....GM.....	60	30	50	48	58
PERCENTILES: <u>1/</u>					
5TH.....GM.....	57			43	32
25TH.....GM.....	97			73	69
50TH.....GM.....	120			101	114
75TH.....GM.....	150			130	135
90TH.....GM.....	195			158	187
95TH.....GM.....	236			228	228
99TH.....GM.....	360			264	270
MAXIMUM.....GM.....	896	102	228	264	342
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	1054.2	0	11.4	26.6	24.9
PROPORTION OF EATING OCCASIONS.....PCT.....	17.6	0	18.2	17.3	14.4

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.2	10.6	13.6	9.2	13.6	10.0	15.7	11.9	19.5	14.6	17.2	15.6
8.3	9.1	10.9	8.2	10.6	8.5	11.7	9.3	13.8	11.1	10.7	11.3
1.8	1.3	2.4	.8	2.7	1.2	3.3	2.1	4.9	3.1	4.7	3.1
.1	.2	.3	.2	.3	.2	.7	.5	.8	.3	1.8	1.2
1.3	1.2	1.3	1.2	1.3	1.2	1.4	1.4	1.4	1.4	1.7	1.4
.6	.6	.6	.5	.7	.6	.6	.6	.7	.7	.7	.7
4.0	5.0	4.0	3.0	5.0	5.0	4.0	4.0	5.0	4.0	3.0	4.0
58	53	62	50	65	51	60	54	67	61	76	58
34	34	41	28	47	31	37	33	49	47	46	35
22	19	23	20	28	19	20	19	22	20	20	19
38	32	39	35	38	33	38	34	38	34	43	34
45	43	45	43	45	43	45	45	51	45	70	45
75	65	80	51	76	61	76	65	86	71	94	68
101	88	110	79	120	88	119	95	132	126	135	126
135	150	135	88	166	108	137	113	151	152	160	132
161	168	225	169	253	163	160	190	280	251	246	158
150	135	146	131	151	128	137	129	146	136	172	134
83	71	67	57	82	56	64	63	76	96	102	68
138	147	179	134	167	128	151	127	160	155	141	116
49	69	79	68	93	46	59	49	76	80	44	50
118	133	100	118	178	162	119	118	152	156	140	131
44	30	15	23	48	30	25	59	96	29	63	6
680	454	466	506	900	342	472	672	506	960	506	384
680	724	802	675	1,152	619	870	830	1,283	960	737	474
133	129	144	126	147	123	131	119	140	130	137	121
55	62	66	47	77	53	51	54	66	63	74	50
59	57	65	60	72	56	57	49	64	60	60	57
101	93	103	101	105	93	101	85	105	90	98	85
128	118	130	128	130	118	128	118	130	125	126	120
152	140	158	135	158	135	152	140	158	158	152	135
227	225	249	189	236	192	195	170	203	192	227	189
252	236	280	225	284	234	228	225	240	210	270	203
284	384	360	270	450	283	312	284	420	448	454	288
360	450	405	350	896	342	450	672	480	480	506	384
50.3	47.3	50.3	34.2	154.8	127.7	197.1	186.9	27.1	52.1	12.1	51.5
18.1	16.8	20.4	21.3	21.8	19.3	18.4	16.4	8.6	14.7	7.9	23.3

TABLE 5.23.--FRUIT PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.1	0	2.1	4.0	4.9
ON ONLY 1 OF 3 DAYS.....PCT.....	5.8	0	1.8	3.4	4.6
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	.3	.6	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	0	1.3	1.2	1.1
STANDARD DEVIATION.....DO.....	.6	0	.7	.5	.3
MAXIMUM.....DO.....	5.0	0	4.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	57	0	35	43	42
STANDARD DEVIATION.....GM.....	39	0	27	25	29
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	21			16	15
25TH.....GM.....	35			25	25
50TH.....GM.....	45			35	39
75TH.....GM.....	67			56	45
90TH.....GM.....	104			79	76
95TH.....GM.....	135			80	90
99TH.....GM.....	208			160	180
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	138	0	81	111	114
STANDARD DEVIATION.....GM.....	72	0	40	56	53
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	152	0	125	116	201
STANDARD DEVIATION.....GM.....	70	0	56	46	82
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	150	0	0	0	0
STANDARD DEVIATION.....GM.....	55	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	960	0	203	270	450
MAXIMUM OVER 3 DAYS.....GM.....	1,283	0	405	480	675
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	133	0	79	106	120
STANDARD DEVIATION.....GM.....	60	0	31	48	56
PERCENTILES: <u>1/</u>					
5TH.....GM.....	57			48	50
25TH.....GM.....	101			75	85
50TH.....GM.....	128			101	118
75TH.....GM.....	158			128	135
90TH.....GM.....	200			158	225
95TH.....GM.....	236			236	236
99TH.....GM.....	384			240	270
MAXIMUM.....GM.....	540	0	158	240	270
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	531.4	0	4.2	5.9	16.2
PROPORTION OF EATING OCCASIONS.....PCT.....	15.4	0	14.5	7.1	17.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.7	6.0	8.2	4.8	8.1	5.4	10.2	6.7	11.9	9.5	11.7	10.7
5.6	5.6	7.0	4.4	6.6	4.6	8.3	5.7	8.5	6.7	7.6	8.1
1.0	.4	1.0	.3	1.3	.7	1.8	.8	3.2	2.4	3.2	2.0
.1	.1	.2	.1	.2	.1	.2	.2	.2	.3	.9	.6
1.3	1.2	1.2	1.1	1.3	1.2	1.3	1.3	1.4	1.5	1.6	1.4
.6	.5	.6	.4	.6	.5	.6	.6	.7	.7	.7	.7
4.0	4.0	4.0	3.0	5.0	4.0	4.0	4.0	5.0	4.0	3.0	4.0
58	51	60	51	65	52	57	51	72	65	77	60
36	38	41	30	47	31	34	32	55	49	47	35
23	19	23	24	28	23	23	19	23	23	23	23
36	28	39	38	39	34	38	34	39	35	45	38
45	42	45	43	45	45	45	45	53	45	70	51
68	53	65	53	76	63	68	58	90	87	90	70
101	96	113	79	120	90	105	90	140	126	135	128
143	151	143	117	180	105	128	105	157	147	169	132
165	177	225	225	292	163	160	190	280	251	246	158
150	130	144	134	155	132	138	128	156	141	175	147
90	76	60	47	88	54	65	56	89	104	88	74
147	204	216	165	170	133	154	131	169	152	134	125
54	84	81	92	91	49	55	52	84	76	40	54
103	157	101	135	177	172	118	138	225	156	192	135
42	0	18	0	41	27	10	63	132	29	47	6
680	454	450	450	500	316	472	450	506	960	506	384
680	724	754	675	900	619	675	830	1,283	960	737	474
133	132	148	135	152	129	135	120	149	133	141	131
54	70	65	47	73	50	53	48	75	60	73	54
57	57	68	71	75	64	68	53	68	68	60	68
101	85	113	105	113	96	105	85	112	96	105	96
128	118	135	128	135	118	128	118	135	126	126	128
151	135	170	135	158	158	158	135	158	158	158	158
227	225	237	192	256	192	196	180	225	189	227	192
256	270	300	227	284	236	236	210	300	196	270	210
284	384	360	270	450	270	360	236	420	480	506	288
360	450	405	350	540	283	450	384	480	480	506	384
26.6	17.5	22.2	19.5	71.4	55.0	117.5	89.6	21.9	27.0	8.2	28.6
14.5	11.5	15.9	24.3	17.4	15.7	18.3	14.9	11.4	11.1	7.9	19.4

TABLE 5.24.--CREAM AND CUSTARD PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.1	.1	2.5	2.7	3.1
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	.1	2.3	2.3	2.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	0	.1	.4	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	(2)	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	2.0	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.0	.4	.5	.5
MAXIMUM.....DO.....	4.0	2.0	2.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	52	47	38	45	45
STANDARD DEVIATION.....GM.....	31	0	20	24	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	19				9
25TH.....GM.....	33				30
50TH.....GM.....	43				41
75TH.....GM.....	63				51
90TH.....GM.....	87				76
95TH.....GM.....	113				95
99TH.....GM.....	163				133
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	140	117	124	119
STANDARD DEVIATION.....GM.....	69	0	61	60	63
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	127	0	27	86	99
STANDARD DEVIATION.....GM.....	57	0	6	32	39
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	117	0	0	133	133
STANDARD DEVIATION.....GM.....	43	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	896	140	228	275	342
MAXIMUM OVER 3 DAYS.....GM.....	896	140	228	400	400
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	124	70	99	109	112
STANDARD DEVIATION.....GM.....	62	31	61	52	59
PERCENTILES: <u>1/</u>					
5TH.....GM.....	49				28
25TH.....GM.....	95				65
50TH.....GM.....	120				114
75TH.....GM.....	138				130
90TH.....GM.....	188				171
95TH.....GM.....	228				228
99TH.....GM.....	340				342
MAXIMUM.....GM.....	896	102	228	264	342
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	379.7	0	6.2	14.0	8.7
PROPORTION OF EATING OCCASIONS.....PCT.....	19.5	0	20.9	24.5	12.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.8	4.0	4.3	3.6	4.4	3.7	4.7	4.5	7.3	4.7	4.2	5.2
2.4	3.4	3.4	3.3	3.6	3.3	3.8	3.7	5.7	4.2	2.3	4.1
.5	.5	.9	.1	.8	.3	.7	.8	1.6	.5	1.3	.7
0	.1	0	.1	(2)	.1	.2	.1	0	0	.5	.4
1.2	1.2	1.3	1.1	1.3	1.2	1.3	1.3	1.3	1.2	1.7	1.3
.5	.6	.5	.5	.6	.5	.6	.5	.5	.4	.8	.6
3.0	4.0	3.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0	3.0	3.0
56	52	57	46	60	47	57	51	53	49	69	45
26	27	37	22	43	25	32	30	27	36	40	28
22	17	11	19	22	15	22	19	21	14		19
38	34	38	33	38	31	38	33	34	27		33
51	43	43	42	43	43	43	43	43	43		38
76	75	76	51	70	51	67	59	60	56		43
87	87	101	75	102	87	101	87	93	77		87
114	88	114	85	131	108	137	101	120	112		130
130	150	267	132	253	114	152	183	135	193		130
152	135	138	127	151	127	136	131	132	130	145	108
66	60	69	47	101	64	55	68	49	88	93	32
126	123	146	123	141	123	146	120	126	144	144	75
35	34	76	58	62	39	51	58	51	66	60	33
0	120	0	105	253	102	124	57	0	0	91	130
0	30	0	23	0	13	22	0	0	0	6	(2)
342	304	466	260	896	342	420	672	304	578	396	203
390	503	802	397	896	378	473	672	405	578	479	391
138	125	135	120	141	120	129	119	124	122	121	100
55	52	65	45	87	60	47	64	46	69	44	37
65	54	49	42	65	43	60	42	48	43		33
112	95	100	98	100	88	100	85	100	65		65
130	114	130	119	130	114	130	114	130	112		101
152	152	152	130	152	130	152	138	152	152		130
227	222	227	182	224	195	195	176	177	210		130
252	228	284	224	260	252	228	228	203	228		152
342	260	342	228	454	342	228	325	280	448		203
342	264	342	228	896	342	420	672	280	448	240	203
18.0	23.1	20.9	8.6	55.8	53.9	49.7	72.6	4.2	19.3	3.9	20.5
24.8	21.5	27.8	14.4	25.2	23.6	16.2	17.8	4.0	19.6	10.2	29.8

TABLE 5.25.--DOUGHNUTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.9	.8	6.6	8.6	10.9
ON ONLY 1 OF 3 DAYS.....PCT.....	7.3	.8	5.6	7.0	9.2
ON ONLY 2 OF 3 DAYS.....PCT.....	2.0	0	.6	1.4	1.6
ON ALL 3 OF 3 DAYS.....PCT.....	.6	0	.4	.3	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.2	1.3	1.3	1.3
STANDARD DEVIATION.....DO.....	.7	.4	.8	.6	.5
MAXIMUM.....DO.....	7.0	2.0	5.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	30	15	20	24	25
STANDARD DEVIATION.....GM.....	23	8	13	15	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		4	8	9
25TH.....GM.....	14		14	14	14
50TH.....GM.....	28		14	17	19
75TH.....GM.....	38		28	28	29
90TH.....GM.....	56		38	42	42
95TH.....GM.....	75		47	56	56
99TH.....GM.....	112		57	84	84
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	69	44	50	61	65
STANDARD DEVIATION.....GM.....	49	25	30	40	37
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	69	0	50	53	66
STANDARD DEVIATION.....GM.....	38	0	9	23	27
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	61	0	48	48	49
STANDARD DEVIATION.....GM.....	27	0	9	10	7
MAXIMUM IN ANY 1 DAY.....GM.....	914	84	142	294	336
MAXIMUM OVER 3 DAYS.....GM.....	1,276	84	171	294	336
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	64	36	47	54	60
STANDARD DEVIATION.....GM.....	40	22	26	28	30
PERCENTILES: <u>1/</u>					
5TH.....GM.....	26		13	25	26
25TH.....GM.....	42		28	42	42
50TH.....GM.....	43		42	42	43
75TH.....GM.....	84		52	57	84
90TH.....GM.....	106		84	85	88
95TH.....GM.....	126		112	126	126
99TH.....GM.....	208		142	168	168
MAXIMUM.....GM.....	851	84	142	171	171
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	20.8	0	0	2.9	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	1.5	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
12.0	12.9	13.2	12.9	12.5	9.1	11.7	8.6	6.8	5.2	7.1	4.4
9.0	10.6	9.2	10.2	8.4	6.9	7.4	6.9	4.4	3.6	4.0	2.7
2.6	2.0	3.5	2.1	3.2	1.6	3.2	1.3	1.6	1.2	1.8	.6
.4	.4	.5	.5	.9	.6	1.1	.5	.8	.5	1.3	1.1
1.4	1.3	1.4	1.3	1.5	1.3	1.6	1.3	1.6	1.5	1.7	1.7
.7	.6	.7	.6	.8	.7	.8	.6	.8	.9	.8	1.0
4.0	5.0	5.0	4.0	5.0	6.0	7.0	5.0	4.0	6.0	4.0	4.0
32	27	43	27	39	27	31	26	29	26	27	33
23	20	42	18	29	18	20	19	15	21	15	18
14	9	14	9	14	9	14	11	14	7		
14	14	19	14	19	14	14	14	14	14		
28	21	28	19	28	19	28	19	28	17		
42	29	56	31	56	33	42	28	42	28		
56	50	84	52	76	56	56	52	56	42		
76	71	113	69	86	58	76	63	56	73		
141	89	196	84	151	84	98	113	56	126		
77	66	89	64	89	65	65	63	62	56	53	73
56	42	66	36	81	37	37	40	34	33	18	36
72	67	108	69	80	60	65	60	63	62	47	82
38	30	87	31	30	24	25	33	17	45	15	31
60	68	95	60	77	61	61	61	45	45	48	41
28	38	55	21	31	19	22	22	4	14	15	12
448	258	851	284	914	252	252	340	168	252	84	171
448	486	1,276	284	914	344	336	420	168	378	225	227
67	62	91	63	77	60	60	58	55	51	49	56
39	36	74	34	52	32	30	32	27	30	17	28
28	26	42	26	42	28	28	25	25	21		
42	42	43	42	42	42	42	42	42	42		
50	43	84	52	84	43	43	42	42	42		
84	84	113	84	84	84	84	84	66	56		
126	99	168	104	126	94	84	85	84	112		
142	126	208	126	168	126	114	114	84	126		
168	227	336	170	252	171	171	168	168	126		
340	252	851	284	638	252	252	340	168	227	84	126
1.0	0	0	1.3	3.0	2.0	4.4	3.4	2.8	0	0	0
.3	0	0	.5	.4	.3	.5	.4	2.4	0	0	0

TABLE 5.26.--SWEET ROLLS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE	MALE	MALE	MALE
		AND FEMALE	AND FEMALE	AND FEMALE	AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.5	0	2.0	2.7	4.3
ON ONLY 1 OF 3 DAYS.....PCT.....	4.3	0	1.8	2.2	4.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	.2	.3	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	0	.1	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER....	1.3	0	1.2	1.3	1.1
STANDARD DEVIATION.....DO.....	.7	0	.4	.8	.2
MAXIMUM.....DO.....	7.0	0	2.0	5.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	32	0	19	28	23
STANDARD DEVIATION.....GM.....	23	0	10	25	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	12				12
25TH.....GM.....	17				17
50TH.....GM.....	24				17
75TH.....GM.....	38				28
90TH.....GM.....	57				38
95TH.....GM.....	71				50
99TH.....GM.....	118				57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	76	0	57	65	68
STANDARD DEVIATION.....GM.....	46	0	30	39	33
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	73	0	35	57	48
STANDARD DEVIATION.....GM.....	32	0	14	15	7
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	72	0	0	133	0
STANDARD DEVIATION.....GM.....	39	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	510	0	113	200	170
MAXIMUM OVER 3 DAYS.....GM.....	780	0	113	400	170
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	72	0	50	62	65
STANDARD DEVIATION.....GM.....	39	0	29	32	31
PERCENTILES: <u>1</u> /					
5TH.....GM.....	35				36
25TH.....GM.....	50				50
50TH.....GM.....	56				50
75TH.....GM.....	85				71
90TH.....GM.....	113				102
95TH.....GM.....	142				142
99TH.....GM.....	227				170
MAXIMUM.....GM.....	510	0	113	200	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER....	12.1	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.8	5.7	7.4	5.6	6.4	4.8	7.4	6.0	6.0	4.0	5.8	6.4
4.5	4.9	5.7	4.9	4.7	3.8	5.3	4.5	5.0	2.7	3.0	4.6
1.1	.7	1.4	.6	1.2	.9	1.4	1.1	.6	.5	1.9	1.6
.2	.1	.2	.2	.5	.2	.6	.4	.4	.8	.9	.3
1.3	1.2	1.3	1.2	1.5	1.3	1.4	1.4	1.3	1.6	1.6	1.3
.7	.4	.6	.5	.8	.5	.7	.7	.6	.9	.7	.6
6.0	3.0	3.0	3.0	7.0	3.0	5.0	4.0	3.0	4.0	3.0	3.0
38	29	36	26	40	30	35	30	32	33	32	25
31	20	24	14	31	20	24	19	20	23	15	15
17	12	17	12	17	12	17	12	17	17		12
17	17	17	17	19	17	17	17	17	17		17
33	21	28	19	33	24	33	24	24	24		17
47	33	50	33	47	33	47	33	38	38		33
67	50	67	47	76	50	67	57	71	67		47
113	57	76	52	113	67	76	68	76	83		67
200	127	113	67	130	113	142	95	95	115		67
91	74	90	68	91	69	81	68	85	72	64	51
61	50	60	29	59	33	43	40	53	46	25	12
83	77	89	63	83	74	72	68	75	59	54	63
38	34	34	13	41	27	33	23	35	18	13	23
117	55	54	63	84	85	74	68	55	57	57	51
76	28	34	13	54	33	37	22	13	28	7	5
425	510	340	200	425	213	250	400	227	340	100	100
695	510	484	263	780	400	542	400	284	344	200	200
85	72	83	64	82	71	74	65	75	62	58	56
51	47	46	23	52	32	37	30	45	36	19	20
35	35	35	35	50	35	35	28	50	18		25
50	50	50	50	50	50	50	50	50	50		50
71	50	71	50	71	57	50	50	50	50		50
100	85	100	71	100	100	100	85	100	71		57
142	107	150	100	113	113	113	100	113	100		100
170	150	170	100	150	140	150	113	211	100		100
284	210	284	142	340	170	227	200	227	150		100
400	510	284	150	340	213	227	284	227	340	100	100
0	6.0	2.2	0	2.1	1.8	0	0	0	0	0	0
0	4.2	1.6	0	.6	.6	0	0	0	0	0	0

TABLE 5.27.--CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	26.2	13.8	38.1	32.8	26.2
ON ONLY 1 OF 3 DAYS.....PCT.....	17.9	4.9	20.2	21.0	19.0
ON ONLY 2 OF 3 DAYS.....PCT.....	6.0	4.6	11.7	9.5	6.0
ON ALL 3 OF 3 DAYS.....PCT.....	2.4	4.3	6.3	2.3	1.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	2.6	1.9	1.6	1.4
STANDARD DEVIATION.....DO.....	.9	1.9	1.3	1.0	.7
MAXIMUM.....DO.....	9.0	9.0	9.0	9.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	11	9	9	10	10
STANDARD DEVIATION.....GM.....	13	10	10	10	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2	2	1	2	2
25TH.....GM.....	4	3	3	4	4
50TH.....GM.....	8	6	6	6	6
75TH.....GM.....	13	10	11	12	12
90TH.....GM.....	23	19	18	21	20
95TH.....GM.....	35	34	28	33	28
99TH.....GM.....	65	56	56	42	56
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	23	10	16	23	20
STANDARD DEVIATION.....GM.....	23	5	18	28	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	24	11	15	19	23
STANDARD DEVIATION.....GM.....	21	5	14	14	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	26	17	19	21	29
STANDARD DEVIATION.....GM.....	25	14	13	11	28
MAXIMUM IN ANY 1 DAY.....GM.....	364	57	170	239	255
MAXIMUM OVER 3 DAYS.....GM.....	1,036	168	227	239	454
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	22	10	14	18	20
STANDARD DEVIATION.....GM.....	21	9	14	20	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	6	3	3	6	6
25TH.....GM.....	12	6	6	9	9
50TH.....GM.....	15	7	12	12	14
75TH.....GM.....	24	14	14	21	24
90TH.....GM.....	42	15	28	30	42
95TH.....GM.....	57	28	28	57	57
99TH.....GM.....	113	56	85	113	113
MAXIMUM.....GM.....	340	57	113	227	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	123.1	3.3	4.8	2.2	8.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	1.8	.6	.2	1.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
22.1	22.1	18.0	19.6	18.5	23.1	26.5	29.9	37.4	34.9	31.6	36.0
16.7	17.3	14.1	15.0	13.7	16.5	18.3	20.0	24.3	21.1	18.3	20.2
4.7	3.8	2.7	3.5	3.9	5.4	5.7	7.2	7.9	8.7	5.0	8.6
.7	1.0	1.3	1.0	.9	1.3	2.4	2.7	5.2	5.1	8.3	7.3
1.4	1.3	1.4	1.4	1.4	1.4	1.6	1.6	1.7	1.6	2.0	1.8
.7	.6	.8	.7	.8	.8	1.0	.9	1.2	.9	1.4	1.2
6.0	4.0	6.0	5.0	6.0	9.0	9.0	6.0	9.0	6.0	7.0	8.0
11	9	14	11	15	10	13	11	12	9	15	11
11	8	15	11	19	11	14	14	13	9	16	13
2	2	2	2	3	2	2	2	2	2	2	2
4	4	5	4	5	4	5	4	5	4	5	4
8	6	9	7	8	7	8	7	8	6	8	6
14	10	19	12	19	12	16	12	12	11	18	12
22	19	29	22	37	19	28	20	28	18	37	22
33	26	50	30	47	30	41	28	36	23	48	30
57	38	76	56	85	57	76	59	70	57	67	66
26	21	32	23	33	22	27	20	22	17	21	16
27	18	33	21	34	20	25	19	18	14	20	16
28	21	39	28	34	24	30	24	21	17	33	19
20	15	28	26	29	19	25	21	11	14	28	16
24	20	32	23	51	25	32	25	31	19	27	24
14	12	24	14	43	25	21	32	22	14	18	20
340	175	227	227	340	224	245	364	143	113	113	132
340	204	336	340	579	421	361	1,036	255	232	227	340
24	20	32	23	33	21	26	20	21	17	22	18
24	16	29	21	35	20	23	20	15	14	19	16
6	6	6	6	6	6	6	6	6	6	6	6
12	12	12	12	12	12	12	12	12	12	12	9
18	14	24	18	24	15	18	14	18	12	18	12
30	24	41	28	40	28	30	24	28	18	24	18
45	39	57	42	57	42	56	32	36	28	44	30
57	56	85	57	113	56	60	56	48	44	57	56
120	85	170	113	170	110	113	113	84	85	113	85
340	113	227	170	340	224	245	196	135	113	113	113
8.1	3.5	8.5	4.7	10.7	22.0	15.4	12.4	5.0	8.0	1.4	4.8
1.3	.6	2.5	1.2	1.0	1.2	.8	.4	.7	.8	.4	.7

TABLE 5.28.--GRAHAM CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,8
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.1	5.2	8.2	6.2	3
ON ONLY 1 OF 3 DAYS.....PCT.....	1.4	2.9	4.7	4.4	2
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	1.1	2.0	1.0	
ON ALL 3 OF 3 DAYS.....PCT.....	.3	1.2	1.5	.8	
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	2.5	1.8	1.5	1
STANDARD DEVIATION.....DO.....	1.0	2.3	.9	.8	
MAXIMUM.....DO.....	9.0	9.0	5.0	4.0	4
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	16	12	11	14	
STANDARD DEVIATION.....GM.....	15	13	9	13	
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		4	5	
25TH.....GM.....	5		5	5	
50TH.....GM.....	9		9	9	
75TH.....GM.....	19		14	19	
90TH.....GM.....	35		19	33	
95TH.....GM.....	47		24	38	
99TH.....GM.....	75		56	76	
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	34	16	23	34	
STANDARD DEVIATION.....GM.....	30	8	14	38	
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	32	11	20	25	
STANDARD DEVIATION.....GM.....	22	5	14	13	
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	32	33	19	23	
STANDARD DEVIATION.....GM.....	21	12	12	11	
MAXIMUM IN ANY 1 DAY.....GM.....	227	57	84	227	1
MAXIMUM OVER 3 DAYS.....GM.....	336	168	168	227	1
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	30	14	19	27	
STANDARD DEVIATION.....GM.....	24	12	12	27	
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		7	7	
25TH.....GM.....	14		14	14	
50TH.....GM.....	28		14	21	
75TH.....GM.....	42		28	28	
90TH.....GM.....	56		28	42	
95TH.....GM.....	84		28	64	
99TH.....GM.....	140		71	151	1
MAXIMUM.....GM.....	227	57	84	227	1
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	4.1	1.0	0	0	
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	1.5	0	0	

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.2	2.3	1.1	1.4	.6	1.4	1.3	1.7	2.8	2.5	2.1	2.4
1.6	1.9	1.0	1.2	.4	1.0	.9	1.1	1.4	1.5	1.6	1.6
.4	.3	.1	.2	.1	.2	.2	.4	.7	.5	.5	.6
.1	.1	(2)	.1	(2)	.2	.2	.2	.6	.6	0	.2
1.5	1.4	1.3	1.3	1.5	1.7	1.7	1.6	1.8	1.8	1.3	1.4
1.0	.7	1.0	.6	.7	1.1	1.3	1.0	.8	1.0	.4	.6
6.0	4.0	6.0	3.0	3.0	6.0	8.0	6.0	3.0	5.0	2.0	3.0
12	13	22	16	27	19	22	16	23	16	19	14
10	9	26	13	20	20	16	14	19	14	11	13
	2				5	5	5				
	5				7	9	9				
	9				12	19	14				
	19				19	28	19				
	28				47	42	28				
	28				56	47	47				
	38				112	75	79				
30	34	53	45	57	35	43	35	27	32	42	25
21	25	55	42	48	29	30	25	21	22	27	9
22	32	56	26	63	45	42	35	44	23	49	28
15	9	0	10	31	27	24	24	16	10	0	13
32	20	112	23	47	37	44	31	50	29	0	56
17	7	0	5	0	34	15	18	14	19	0	0
140	113	227	170	154	154	140	154	112	84	85	56
168	113	336	170	252	336	224	238	196	196	98	168
25	28	54	38	55	35	37	31	40	27	45	31
18	19	47	35	39	23	23	24	23	19	21	15
	14				14	14	14				
	14				14	28	14				
	28				28	28	28				
	28				56	42	42				
	56				70	57	56				
	56				84	84	84				
	113				85	140	140				
140	113	227	170	154	85	140	154	112	84	85	56
0	0	1.0	.6	0	0	.8	0	0	0	0	.8
0	0	4.9	2.3	0	0	.7	0	0	0	0	2.4

TABLE 5.29.--SALTINES AND SODA CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	AGE (YEARS) AND SEX OF INDIVIDUAL				
	TOTAL	UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	14.5	6.3	20.1	17.9	14.1
ON ONLY 1 OF 3 DAYS.....PCT.....	10.9	3.1	12.6	13.8	11.7
ON ONLY 2 OF 3 DAYS.....PCT.....	2.8	1.9	5.1	3.6	2.3
ON ALL 3 OF 3 DAYS.....PCT.....	.8	1.3	2.5	.4	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	2.0	1.6	1.4	1.2
STANDARD DEVIATION.....DO.....	.7	1.2	1.0	.6	.5
MAXIMUM.....DO.....	9.0	5.0	6.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	8	5	6	6	7
STANDARD DEVIATION.....GM.....	9	4	8	6	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	4		2	3	3
50TH.....GM.....	6		4	4	4
75TH.....GM.....	9		6	7	8
90TH.....GM.....	16		11	12	14
95TH.....GM.....	22		17	14	20
99TH.....GM.....	48		40	38	40
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	18	7	10	15	16
STANDARD DEVIATION.....GM.....	17	4	12	15	16
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	11	13	13	20
STANDARD DEVIATION.....GM.....	19	5	16	9	15
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	9	13	10	20
STANDARD DEVIATION.....GM.....	16	5	12	3	15
MAXIMUM IN ANY 1 DAY.....GM.....	227	30	170	113	170
MAXIMUM OVER 3 DAYS.....GM.....	355	60	227	113	182
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	18	8	11	13	17
STANDARD DEVIATION.....GM.....	17	5	14	12	17
PERCENTILES: <u>1/</u>					
5TH.....GM.....	6		3	6	6
25TH.....GM.....	9		6	6	6
50TH.....GM.....	12		6	12	12
75TH.....GM.....	18		12	12	18
90TH.....GM.....	30		18	21	30
95TH.....GM.....	42		24	28	36
99TH.....GM.....	113		113	57	60
MAXIMUM.....GM.....	227	30	113	113	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	31.7	1.0	2.6	1.3	.9
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	1.5	.8	.3	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
12.7	12.4	10.2	10.1	9.2	11.8	14.7	17.2	23.9	21.0	16.9	21.2
10.1	10.8	8.6	8.0	7.1	9.2	10.5	12.9	17.8	14.2	9.9	12.9
2.4	1.3	1.1	1.8	1.9	2.0	3.1	3.5	3.9	5.1	3.2	5.6
.2	.2	.4	.4	.2	.6	1.1	.8	2.3	1.8	3.8	2.7
1.3	1.2	1.2	1.3	1.3	1.3	1.5	1.4	1.5	1.5	1.9	1.6
.6	.5	.6	.7	.6	.6	.9	.7	.9	.7	1.4	.9
5.0	3.0	4.0	5.0	5.0	6.0	9.0	5.0	6.0	4.0	7.0	6.0
9	6	12	9	11	7	11	8	9	8	12	8
8	5	13	10	13	6	12	9	8	8	13	8
2	2	2	2	2	2	2	2	2	2	4	2
4	3	4	4	4	4	4	4	4	4	5	4
6	4	8	6	6	6	8	6	7	6	6	6
10	8	12	10	10	8	12	8	10	9	12	10
18	12	20	19	21	12	22	14	16	15	31	16
26	16	40	24	38	16	32	19	28	18	47	19
41	26	76	76	76	28	57	40	48	47	76	30
20	16	26	22	25	16	23	16	20	15	18	14
17	13	23	25	26	11	22	13	17	11	14	7
25	18	42	22	23	16	26	20	16	19	25	18
18	10	38	19	18	9	24	20	6	17	26	18
23	12	27	19	51	14	24	21	24	15	23	14
15	3	19	13	44	13	16	16	13	6	16	4
132	131	227	227	227	113	227	227	143	113	113	125
227	131	247	227	355	251	340	312	174	170	227	239
21	15	28	20	24	15	22	17	18	16	18	14
18	12	26	19	25	11	20	15	12	12	16	11
6	6	6	6	6	6	6	6	6	6	6	3
12	9	12	12	12	9	12	12	12	12	12	9
15	12	20	15	18	12	18	12	15	12	12	12
27	18	30	24	24	18	24	18	24	18	24	18
36	24	57	36	48	24	36	28	30	24	28	21
48	30	113	48	85	30	57	30	36	30	28	28
113	60	113	113	113	57	113	113	60	57	113	57
120	113	140	140	227	113	227	170	135	113	113	113
3.8	1.6	.9	1.3	1.6	4.6	7.0	2.0	2.1	0	0	1.0
1.1	.5	.5	.7	.3	.6	.7	.1	.5	0	0	.3

TABLE 5.30.--BUTTER CRACKERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.4	2.6	7.6	5.5	5.3
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	1.4	4.8	4.5	4.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	1.0	2.2	.9	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.3	.2	.6	.1	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.5	1.6	1.3	1.3
STANDARD DEVIATION.....DO.....	.7	.6	.8	.7	.6
MAXIMUM.....DO.....	7.0	3.0	6.0	6.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	9	4	6	6	9
STANDARD DEVIATION.....GM.....	10	2	4	5	14
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	4		3	3	4
50TH.....GM.....	6		6	5	6
75TH.....GM.....	10		8	8	9
90TH.....GM.....	18		12	12	13
95TH.....GM.....	24		15	16	18
99TH.....GM.....	50		23	28	30
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	7	14	17	19
STANDARD DEVIATION.....GM.....	19	3	11	13	11
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	8	12	12	19
STANDARD DEVIATION.....GM.....	14	2	4	4	11
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	24	6	13	24	151
STANDARD DEVIATION.....GM.....	22	0	5	(2)	0
MAXIMUM IN ANY 1 DAY.....GM.....	340	15	60	85	255
MAXIMUM OVER 3 DAYS.....GM.....	454	21	69	85	454
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	20	7	12	14	21
STANDARD DEVIATION.....GM.....	18	3	7	11	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	6		3	4	4
25TH.....GM.....	12		6	8	12
50TH.....GM.....	18		12	12	18
75TH.....GM.....	24		15	16	24
90TH.....GM.....	36		18	27	36
95TH.....GM.....	50		28	30	57
99TH.....GM.....	113		30	57	113
MAXIMUM.....GM.....	340	15	30	85	142
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	7.2	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.4	3.4	3.0	3.7	3.6	4.2	4.9	4.5	4.0	4.8	4.6	7.3
2.9	2.8	2.4	2.8	2.9	3.6	3.7	3.4	2.4	3.6	2.9	3.9
.5	.5	.4	.7	.5	.4	.9	.9	1.4	.3	.2	1.7
(2)	.1	.2	.3	.1	.1	.3	.2	.2	.9	1.5	1.7
1.2	1.2	1.3	1.4	1.3	1.2	1.3	1.3	1.6	1.5	1.8	2.0
.5	.5	.7	.7	.6	.4	.6	.6	1.0	.9	1.1	1.4
3.0	3.0	4.0	4.0	4.0	3.0	5.0	4.0	7.0	4.0	4.0	6.0
11	7	13	10	14	8	10	8	11	7	14	12
12	4	10	7	14	6	11	8	10	4	10	17
3	2		3	3	2	2	2		2		1
6	4		4	6	4	4	4		4		4
8	6		8	8	6	6	6		5		8
14	8		14	16	10	12	10		10		12
20	12		21	38	16	20	15		12		21
20	15		27	47	19	24	20		16		50
113	25		30	76	33	54	47		22		113
29	19	29	21	32	21	22	18	25	16	33	14
38	12	18	15	31	13	16	18	18	9	32	9
25	16	37	24	35	23	25	20	17	16	54	17
12	7	25	11	28	10	15	11	8	2	0	9
20	12	23	22	47	27	33	20	34	13	17	28
0	0	8	7	9	14	26	14	28	4	5	29
340	75	100	71	170	60	140	142	99	57	113	113
340	75	180	90	227	135	294	165	210	66	113	340
26	18	29	21	32	21	23	18	21	14	23	18
32	11	19	13	29	13	17	14	13	7	21	19
9	6		6	5	6	6	6		6		3
15	12		12	14	12	12	12		12		9
21	15		18	24	18	18	15		12		12
30	18		30	36	28	28	24		18		18
45	30		36	57	42	45	30		18		30
50	45		48	113	50	56	45		24		40
85	57		71	142	60	85	60		57		113
340	75	100	71	170	60	113	142	84	57	113	113
0	0	0	0	0	2.9	1.3	.8	0	0	0	2.2
0	0	0	0	0	1.1	.4	.2	0	0	0	1.5

TABLE 5.31.--POPCORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.6	.1	5.7	8.5	9.5
ON ONLY 1 OF 3 DAYS.....PCT.....	4.9	.1	4.9	7.6	7.9
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	0	.8	.8	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	.1	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.0	1.2	1.1	1.2
STANDARD DEVIATION.....DO.....	.5	.0	.4	.4	.5
MAXIMUM.....DO.....	4.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	7	24	3	5	6
STANDARD DEVIATION.....GM.....	9	0	4	4	4
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	1	2
25TH.....GM.....	3		2	2	3
50TH.....GM.....	6		3	3	6
75TH.....GM.....	9		3	6	6
90TH.....GM.....	12		6	9	9
95TH.....GM.....	18		8	12	12
99TH.....GM.....	43		38	13	28
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	19	72	10	12	16
STANDARD DEVIATION.....GM.....	23	0	14	8	10
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	19	0	7	15	10
STANDARD DEVIATION.....GM.....	15	0	3	15	4
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	0	0	10	16
STANDARD DEVIATION.....GM.....	20	0	0	0	9
MAXIMUM IN ANY 1 DAY.....GM.....	340	72	113	130	98
MAXIMUM OVER 3 DAYS.....GM.....	340	72	113	139	98
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	19	72	9	12	14
STANDARD DEVIATION.....GM.....	22	0	12	11	9
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		2	3	5
25TH.....GM.....	9		5	6	9
50TH.....GM.....	15		6	9	12
75TH.....GM.....	18		9	18	18
90TH.....GM.....	36		14	18	27
95TH.....GM.....	45		24	27	28
99TH.....GM.....	108		113	36	36
MAXIMUM.....GM.....	340	72	113	130	98
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	66.2	0	0	6.2	12.3
PROPORTION OF EATING OCCASIONS.....PCT.....	2.7	0	0	3.7	5.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
9.6	9.1	6.1	7.8	6.5	7.1	4.1	3.8	1.5	.9	.3	.8
8.7	7.7	4.9	6.6	5.8	6.2	3.5	3.4	1.4	.9	.2	.6
.8	1.1	1.1	.9	.5	.7	.5	.3	.1	0	.1	.2
.1	.3	.1	.3	.1	.2	.1	(2)	0	0	0	0
1.1	1.2	1.3	1.2	1.1	1.2	1.2	1.1	1.5	1.0	1.5	1.2
.4	.6	.5	.5	.4	.5	.5	.4	.9	.0	.5	.4
3.0	4.0	3.0	3.0	4.0	3.0	4.0	3.0	4.0	1.0	2.0	2.0
7	7	8	7	9	7	11	6	9	4	12	5
6	6	10	9	12	9	15	6	9	1	6	1
2	2	2	2	1	1	1	2				
3	3	4	3	3	3	4	3				
6	6	6	6	6	6	8	6				
9	9	9	6	12	9	12	7				
12	12	15	12	19	12	18	12				
15	18	19	24	24	15	32	18				
28	28	84	72	90	38	95	43				
19	18	20	16	25	20	28	16	18	11	18	14
18	14	15	12	34	26	41	11	6	4	0	3
15	17	23	21	26	19	22	23	54	0	27	9
7	15	29	17	11	12	13	15	0	0	0	0
6	14	9	32	22	10	56	38	0	0	0	0
(2)	5	0	27	30	2	12	23	0	0	0	0
170	112	144	180	320	340	284	112	54	18	36	18
170	130	252	216	320	340	284	166	108	18	54	18
18	17	20	18	24	19	27	18	18	11	24	12
17	15	20	20	29	24	36	13	11	4	8	3
5	5	5	5	3	5	5	5				
9	9	9	9	9	9	9	9				
18	14	18	18	18	18	18	18				
18	18	18	18	27	24	28	18				
36	27	36	27	48	36	45	36				
36	36	48	45	68	36	72	36				
85	85	108	112	113	85	284	56				
170	112	144	180	284	340	284	112	54	18	36	18
12.3	9.0	5.8	8.6	3.4	4.5	0	4.1	0	0	0	0
5.5	3.7	5.4	6.2	1.2	1.0	0	1.4	0	0	0	0

TABLE 5.32.--PRETZELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.2	.7	3.2	3.1	3.3
ON ONLY 1 OF 3 DAYS.....PCT.....	1.7	.4	2.3	2.5	2.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	.3	.9	.6	.3
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	(2)	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	1.9	1.5	1.3	1.2
STANDARD DEVIATION.....DO.....	.6	1.1	1.0	.5	.5
MAXIMUM.....DO.....	5.0	4.0	5.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	13	3	9	9	10
STANDARD DEVIATION.....GM.....	15	1	9	9	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1			(2)	1
25TH.....GM.....	5			2	4
50TH.....GM.....	9			6	8
75TH.....GM.....	16			11	11
90TH.....GM.....	28			19	19
95TH.....GM.....	42			30	30
99TH.....GM.....	64			41	75
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	10	20	22	25
STANDARD DEVIATION.....GM.....	28	5	24	22	28
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	3	24	20	26
STANDARD DEVIATION.....GM.....	23	1	14	16	17
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	38	0	0	41	24
STANDARD DEVIATION.....GM.....	37	0	0	0	9
MAXIMUM IN ANY 1 DAY.....GM.....	224	15	113	90	224
MAXIMUM OVER 3 DAYS.....GM.....	480	15	113	123	224
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	29	4	18	21	25
STANDARD DEVIATION.....GM.....	28	4	18	20	21
PERCENTILES: <u>1/</u>					
5TH.....GM.....	3			1	4
25TH.....GM.....	12			5	12
50TH.....GM.....	21			16	23
75TH.....GM.....	36			32	28
90TH.....GM.....	57			56	57
95TH.....GM.....	85			57	68
99TH.....GM.....	160			90	112
MAXIMUM.....GM.....	170	15	113	90	112
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	21.6	.8	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.9	12.7	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.1	3.5	2.9	3.1	2.4	2.3	1.9	1.2	1.2	.5	2.1	.6
2.9	2.5	2.1	2.4	1.8	1.8	1.5	1.0	.4	.3	1.5	.6
.8	.7	.4	.5	.6	.4	.3	.2	.5	.2	.6	0
.4	.2	.3	.2	.1	.1	0	.1	.3	0	0	0
1.5	1.4	1.4	1.3	1.4	1.3	1.2	1.3	1.9	1.4	1.5	1.0
.8	.6	.7	.6	.6	.6	.4	.7	.7	.5	.5	.0
4.0	3.0	3.0	3.0	3.0	4.0	2.0	4.0	3.0	2.0	2.0	1.0
15	14	25	11	19	9	13	10	27	12	5	12
12	12	38	8	15	9	13	12	22	4	3	4
2	2			1	1	1	2				
5	5			8	4	5	3				
11	9			15	6	9	5				
19	16			23	13	17	10				
33	30			47	19	21	19				
38	43			51	28	43	47				
47	57			67	50	76	54				
34	32	38	26	51	20	27	24	70	30	17	36
29	26	24	16	42	17	18	26	66	14	11	13
33	27	34	20	35	29	43	23	19	22	6	0
17	21	21	10	27	20	30	29	8	1	(2)	0
23	34	102	32	35	17	0	28	56	0	0	0
18	12	68	3	8	2	0	18	10	0	0	0
142	135	160	57	172	113	170	142	142	48	30	57
225	170	480	105	202	149	227	163	192	48	30	57
29	30	52	25	42	21	32	23	41	26	10	36
25	26	50	16	38	19	26	28	36	11	6	13
6	2			3	3	6	3				
14	12			17	9	15	9				
23	23			30	15	24	15				
42	43			50	28	45	28				
57	64			113	45	57	48				
85	90			142	57	72	57				
113	113			170	113	170	142				
142	135	160	57	170	113	170	142	142	48	23	57
3.4	1.9	.9	1.1	4.2	0	2.3	7.0	0	0	0	0
2.7	1.9	1.5	1.9	3.3	0	2.1	6.2	0	0	0	0

TABLE 5.33.--POTATO CHIPS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.0	.8	19.3	21.7	26.7
ON ONLY 1 OF 3 DAYS.....PCT.....	12.1	.6	13.7	15.7	17.3
ON ONLY 2 OF 3 DAYS.....PCT.....	3.7	.2	4.7	4.9	7.1
ON ALL 3 OF 3 DAYS.....PCT.....	1.2	0	1.0	1.1	2.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.3	1.4	1.4	1.5
STANDARD DEVIATION.....DO.....	.7	.4	.7	.7	.8
MAXIMUM.....DO.....	6.0	2.0	5.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	11	4	7	8	11
STANDARD DEVIATION.....GM.....	11	1	8	7	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		2	2	2
25TH.....GM.....	5		3	3	4
50TH.....GM.....	7		5	6	7
75TH.....GM.....	13		9	9	13
90TH.....GM.....	24		15	15	21
95TH.....GM.....	33		21	19	31
99TH.....GM.....	57		49	38	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	24	14	14	18	20
STANDARD DEVIATION.....GM.....	27	3	13	17	18
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	24	3	19	16	24
STANDARD DEVIATION.....GM.....	18	0	16	9	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	25	0	20	20	25
STANDARD DEVIATION.....GM.....	15	0	11	11	14
MAXIMUM IN ANY 1 DAY.....GM.....	340	18	142	170	170
MAXIMUM OVER 3 DAYS.....GM.....	340	18	156	170	255
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	23	10	16	17	21
STANDARD DEVIATION.....GM.....	23	6	17	13	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		5	5	5
25TH.....GM.....	9		9	9	9
50TH.....GM.....	18		10	14	18
75TH.....GM.....	28		18	18	23
90TH.....GM.....	43		28	28	36
95TH.....GM.....	57		43	40	57
99TH.....GM.....	120		113	57	128
MAXIMUM.....GM.....	340	18	142	170	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	240.0	0	6.9	6.7	21.7
PROPORTION OF EATING OCCASIONS.....PCT.....	2.6	0	2.4	1.3	2.9

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
29.0	29.0	28.7	25.0	23.5	17.7	12.4	10.2	3.8	3.5	2.7	4.2
20.0	20.0	19.4	18.8	15.1	13.5	8.8	8.4	2.8	2.8	1.6	3.6
6.9	6.5	6.9	5.0	6.0	3.4	2.7	1.4	1.0	.5	1.1	.2
2.1	2.5	2.4	1.2	2.4	.9	.9	.4	0	.1	0	.4
1.5	1.5	1.5	1.4	1.6	1.4	1.4	1.3	1.3	1.2	1.4	1.3
.7	.8	.8	.7	.8	.7	.7	.6	.4	.5	.5	.8
4.0	6.0	6.0	6.0	6.0	5.0	4.0	5.0	2.0	3.0	2.0	4.0
12	12	14	11	14	10	11	9	8	6	5	8
12	11	14	12	14	9	9	10	8	5	3	6
2	2	2	2	2	2	2	2		2		
6	6	6	5	6	5	5	3		3		
9	9	9	7	9	7	7	6		6		
14	14	18	12	18	12	12	9		8		
24	24	28	24	30	19	22	19		9		
38	33	46	38	42	27	31	24		9		
57	57	76	66	66	38	43	57		30		
28	27	33	26	29	24	24	22	22	17	12	19
29	27	38	26	35	25	21	27	26	8	5	12
27	26	27	26	27	22	23	21	18	8	10	13
21	19	20	20	19	12	15	16	9	3	5	0
21	20	30	31	31	21	23	24	0	30	0	20
9	10	18	24	18	8	11	12	0	0	0	8
284	227	227	227	340	340	128	284	142	36	18	73
312	255	245	263	340	340	170	284	142	90	27	76
25	24	28	25	27	21	23	21	20	16	11	18
25	22	29	24	28	20	18	24	22	9	6	16
5	6	5	5	5	5	5	5		3		
11	11	14	11	10	9	14	9		8		
18	18	18	18	18	18	18	18		18		
28	28	32	28	30	27	28	24		20		
54	43	57	53	57	36	43	36		28		
57	57	85	60	57	57	57	57		36		
128	128	170	113	142	85	90	128		36		
284	227	227	227	340	340	128	284	142	36	18	70
24.1	24.7	14.4	17.4	32.1	18.8	31.5	33.4	5.2	3.2	0	0
2.7	2.7	2.4	3.5	2.2	1.4	3.6	3.7	9.7	4.4	0	0

TABLE 5.34.--CORN-BASED SALTY SNACKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.9	.6	6.6	8.6	10.3
ON ONLY 1 OF 3 DAYS.....PCT.....	4.5	.4	5.2	6.8	7.5
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	1.3	1.3	1.9
ON ALL 3 OF 3 DAYS.....PCT.....	.3	.2	.1	.4	.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.7	1.3	1.3	1.4
STANDARD DEVIATION.....DO.....	.7	1.0	.5	.6	.8
MAXIMUM.....DO.....	8.0	3.0	3.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	15	4	10	12	14
STANDARD DEVIATION.....GM.....	15	2	9	10	13
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3		1	3	3
25TH.....GM.....	7		4	5	7
50TH.....GM.....	12		7	7	9
75TH.....GM.....	19		14	13	17
90TH.....GM.....	28		22	27	30
95TH.....GM.....	38		27	38	38
99TH.....GM.....	76		38	57	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	35	9	25	28	32
STANDARD DEVIATION.....GM.....	31	2	22	25	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	34	0	28	24	30
STANDARD DEVIATION.....GM.....	25	0	13	12	14
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	37	7	5	33	29
STANDARD DEVIATION.....GM.....	33	0	0	16	16
MAXIMUM IN ANY 1 DAY.....GM.....	340	11	113	113	340
MAXIMUM OVER 3 DAYS.....GM.....	542	21	120	170	340
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	33	8	24	27	29
STANDARD DEVIATION.....GM.....	30	2	20	22	26
PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		4	10	10
25TH.....GM.....	18		10	13	16
50TH.....GM.....	21		18	20	21
75TH.....GM.....	40		36	36	35
90TH.....GM.....	60		54	43	57
95TH.....GM.....	80		57	72	80
99TH.....GM.....	156		85	113	113
MAXIMUM.....GM.....	340	11	113	113	340
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	67.2	1.0	4.0	2.8	4.0
PROPORTION OF EATING OCCASIONS.....PCT.....	2.2	20.2	4.6	1.5	1.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
9.9	11.3	8.3	10.7	8.2	6.3	3.8	3.1	1.2	.5	1.4	1.7
6.9	8.0	6.5	8.5	6.0	5.1	2.8	2.6	1.2	.5	1.2	1.3
2.5	2.7	1.4	1.7	1.8	.9	.7	.4	0	0	.2	.2
.5	.6	.4	.6	.4	.3	.3	.1	0	0	0	.2
1.4	1.4	1.4	1.4	1.3	1.2	1.4	1.3	1.2	1.0	1.1	1.4
.7	.7	.8	.7	.6	.5	.7	.9	.4	.0	.3	.7
4.0	4.0	5.0	5.0	5.0	3.0	4.0	8.0	2.0	1.0	2.0	3.0
16	15	21	16	18	14	16	14	8	6	6	10
15	13	22	11	19	13	13	19	6	2	4	8
3	3	5	3	3	4	3	2				
7	7	7	7	7	7	7	6				
13	13	13	13	13	12	13	9				
19	20	27	20	20	14	23	13				
30	28	52	28	30	27	28	27				
41	40	71	38	40	38	33	40				
95	57	95	53	88	75	61	151				
37	37	50	39	39	32	39	33	25	19	19	21
34	37	48	26	26	20	31	35	19	7	13	15
33	32	52	34	38	46	25	27	0	0	5	18
27	18	48	20	20	36	14	14	0	0	0	0
36	26	45	34	56	26	35	74	0	0	0	28
16	9	25	8	59	8	21	65	0	0	0	0
284	340	227	128	340	252	184	240	57	28	40	56
288	340	397	213	542	280	280	454	57	28	40	85
33	32	46	34	39	34	33	32	21	19	15	23
29	30	44	22	36	28	26	30	11	7	13	12
7	9	14	10	10	10	10	7				
20	20	20	20	20	20	20	14				
22	21	30	28	28	28	21	21				
40	40	54	40	44	40	40	40				
60	57	85	60	80	57	72	57				
80	80	160	85	85	80	80	85				
128	170	213	113	170	156	120	170				
284	340	227	128	340	252	184	240	40	28	40	56
5.4	5.3	4.2	7.9	8.6	9.0	8.8	4.1	0	0	0	2.2
1.8	1.5	2.6	3.7	2.0	2.2	3.3	1.4	0	0	0	9.2

TABLE 5.35.--PASTA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	11.4	3.4	14.1	14.7	14.5
ON ONLY 1 OF 3 DAYS.....PCT.....	10.1	2.4	11.5	12.3	12.7
ON ONLY 2 OF 3 DAYS.....PCT.....	1.3	.7	2.5	2.3	1.8
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	.3	.1	(2)	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.5	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.4	.7	.5	.4	.4
MAXIMUM.....DO.....	6.0	3.0	4.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	59	28	33	39	45
STANDARD DEVIATION.....GM.....	46	32	26	26	31
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	12		5	12	9
25TH.....GM.....	27		13	23	23
50TH.....GM.....	47		27	27	40
75TH.....GM.....	80		47	47	53
90TH.....GM.....	107		70	70	93
95TH.....GM.....	140		93	93	103
99TH.....GM.....	233		123	123	160
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	156	50	83	100	119
STANDARD DEVIATION.....GM.....	110	30	58	61	75
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	159	52	82	100	122
STANDARD DEVIATION.....GM.....	106	42	45	41	61
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	141	98	192	157	0
STANDARD DEVIATION.....GM.....	61	48	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	980	160	490	420	480
MAXIMUM OVER 3 DAYS.....GM.....	1,960	413	577	472	620
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	153	58	82	99	116
STANDARD DEVIATION.....GM.....	108	42	59	58	74
PERCENTILES: <u>1/</u>					
5TH.....GM.....	35		14	35	27
25TH.....GM.....	70		40	70	70
50TH.....GM.....	140		70	80	105
75TH.....GM.....	210		105	140	140
90TH.....GM.....	280		160	160	210
95TH.....GM.....	320		210	210	280
99TH.....GM.....	560		280	280	320
MAXIMUM.....GM.....	980	160	350	400	480
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	22.6	0	.7	.7	3.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.4	.2	1.1

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.0	14.5	11.2	10.8	12.3	12.6	10.8	9.9	8.9	6.7	7.5	7.3
12.7	13.1	10.3	9.8	10.8	11.0	9.8	8.7	7.8	5.8	6.8	5.7
1.2	1.4	.9	1.0	1.4	1.5	1.0	1.1	.9	.9	.7	1.4
.1	(2)	0	0	.1	(2)	0	(2)	.3	0	0	.2
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.2	1.1	1.3
.4	.4	.4	.3	.4	.4	.3	.4	.8	.5	.3	.5
3.0	4.0	4.0	2.0	3.0	3.0	3.0	3.0	6.0	4.0	2.0	3.0
60	54	75	59	76	61	70	54	57	50	51	51
42	38	58	43	60	45	52	38	47	48	28	31
13	12	20	12	13	13	17	12	18	13		7
27	27	38	27	41	27	35	27	27	26		27
47	47	53	47	53	47	53	47	47	40		47
93	70	93	80	93	80	93	70	76	53		70
107	105	151	107	147	107	135	104	107	93		100
140	117	210	157	187	140	187	117	140	107		107
213	187	280	207	280	280	280	210	245	373		107
164	142	198	160	201	158	192	147	146	126	145	131
104	89	144	114	141	103	129	95	91	75	82	83
169	173	250	177	208	179	200	138	143	145	113	118
97	75	118	66	141	102	128	84	112	151	22	39
140	110	0	0	171	80	0	210	165	0	0	80
0	0	0	0	47	0	0	0	87	0	0	0
640	560	720	800	980	640	840	700	613	560	320	320
840	700	1,040	800	1,960	980	1,120	840	900	1,120	320	320
162	145	198	158	198	160	189	142	138	127	133	121
102	89	133	99	141	101	123	95	108	84	69	71
40	35	61	35	40	40	40	35	35	35		20
80	80	120	80	105	80	105	70	70	70		80
140	140	140	140	160	140	160	140	105	85		99
210	175	280	210	280	210	280	160	160	160		160
280	280	320	280	350	280	320	280	280	280		210
320	280	454	320	480	350	420	320	420	320		280
560	420	701	480	700	420	640	480	560	341		320
640	560	720	560	980	640	840	700	560	480	320	320
.8	0	1.4	.9	1.1	5.3	4.0	2.2	0	0	2.1	0
.2	0	.8	.5	.2	.7	.7	.3	0	0	4.5	0

TABLE 5.36.--SPAGHETTI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.6	1.6	5.8	7.1	7.9
ON ONLY 1 OF 3 DAYS.....PCT.....	5.2	1.3	5.2	6.2	7.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	.3	.7	.9	.6
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	1.1	1.3	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.3	.5	.3	.3	.3
MAXIMUM.....DO.....	4.0	2.0	2.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	63	22	31	40	43
STANDARD DEVIATION.....GM.....	48	25	21	26	25
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	12		9	12	9
25TH.....GM.....	35		18	23	23
50TH.....GM.....	47		23	35	47
75TH.....GM.....	93		47	47	47
90TH.....GM.....	107		47	93	82
95TH.....GM.....	140		82	93	93
99TH.....GM.....	245		93	117	117
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	173	42	85	104	124
STANDARD DEVIATION.....GM.....	118	33	58	64	69
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	185	82	72	111	105
STANDARD DEVIATION.....GM.....	124	52	40	44	52
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	245	0	0	0	0
STANDARD DEVIATION.....GM.....	0	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	980	140	280	280	280
MAXIMUM OVER 3 DAYS.....GM.....	1,960	253	280	350	350
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	172	53	83	104	118
STANDARD DEVIATION.....GM.....	117	39	55	59	66
PERCENTILES: <u>1</u> /					
5TH.....GM.....	35		26	35	28
25TH.....GM.....	87		35	70	70
50TH.....GM.....	140		70	105	140
75TH.....GM.....	210		140	140	140
90TH.....GM.....	280		140	140	210
95TH.....GM.....	420		140	210	280
99TH.....GM.....	561		280	280	280
MAXIMUM.....GM.....	980	130	280	280	280
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	10.1	0	.7	.7	2.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	1.0	.5	1.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.1	7.4	6.4	5.4	6.0	6.4	5.0	4.9	4.2	2.6	3.4	2.6
6.5	7.0	5.9	5.1	5.7	5.8	4.7	4.5	4.0	2.2	3.4	2.5
.6	.4	.5	.3	.3	.6	.2	.4	.1	.3	0	.1
0	0	0	0	0	0	0	0	.1	0	0	0
1.1	1.1	1.2	1.1	1.1	1.1	1.1	1.1	1.2	1.2	1.1	1.0
.3	.3	.5	.3	.2	.3	.2	.3	.6	.6	.3	.2
3.0	3.0	4.0	2.0	2.0	2.0	3.0	2.0	4.0	4.0	2.0	2.0
64	59	82	58	83	67	74	58	66	59	44	41
43	37	58	36	67	47	51	39	50	63	22	24
18	12	23	12	18	18	23	12				
38	38	47	35	47	47	38	35				
47	47	70	47	70	47	65	47				
93	70	93	70	93	93	93	93				
117	93	187	107	187	117	140	93				
140	117	213	140	187	140	187	140				
233	187	280	163	280	280	280	187				
172	165	222	164	233	177	212	162	175	139	133	121
96	94	150	97	163	103	138	102	109	82	66	72
207	194	269	174	267	222	218	166	175	214	0	92
104	77	95	61	215	111	145	80	0	199	0	0
0	0	0	0	0	0	0	0	245	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
560	560	700	490	980	640	840	700	613	560	280	280
840	700	840	490	1,960	840	980	840	736	1,120	280	280
176	166	213	162	236	183	210	161	170	148	119	118
97	93	129	87	170	108	135	99	124	86	38	70
53	35	70	53	53	53	70	35				
113	112	140	105	140	105	112	105				
140	140	140	140	210	140	158	140				
210	210	280	210	280	280	280	210				
280	280	420	280	448	280	350	280				
280	280	454	320	561	420	420	280				
560	420	641	420	841	560	701	560				
560	560	700	420	980	640	840	700	560	341	210	280
.8	0	1.4	.9	0	1.4	1.6	0	0	0	0	0
.5	0	1.4	1.0	0	.4	.6	0	0	0	0	0

TABLE 5.37.--NOODLES

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.7	1.5	5.3	4.8	4.6
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	.9	4.4	4.3	4.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	.5	.9	.5	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	.1	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.6	1.2	1.1	1.1
STANDARD DEVIATION.....DO.....	.3	.7	.4	.3	.3
MAXIMUM.....DO.....	3.0	3.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	52	34	29	34	42
STANDARD DEVIATION.....GM.....	41	20	18	21	32
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	13		3	13	13
25TH.....GM.....	27		13	27	27
50TH.....GM.....	53		27	27	27
75TH.....GM.....	53		40	53	53
90TH.....GM.....	107		53	53	80
95TH.....GM.....	107		60	67	107
99TH.....GM.....	213		80	133	160
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	143	89	77	97	121
STANDARD DEVIATION.....GM.....	103	41	50	59	86
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	141	60	64	80	118
STANDARD DEVIATION.....GM.....	105	44	26	30	80
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	119	40	0	0	0
STANDARD DEVIATION.....GM.....	60	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	800	160	240	400	480
MAXIMUM OVER 3 DAYS.....GM.....	1,120	240	240	400	480
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	140	65	71	93	117
STANDARD DEVIATION.....GM.....	98	47	49	56	86
PERCENTILES: <u>1/</u>					
5TH.....GM.....	40		10	40	40
25TH.....GM.....	80		40	61	80
50TH.....GM.....	120		61	80	80
75TH.....GM.....	160		80	120	160
90TH.....GM.....	320		160	160	200
95TH.....GM.....	320		160	160	320
99TH.....GM.....	480		240	400	480
MAXIMUM.....GM.....	720	160	240	400	480
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	6.7	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.9	4.6	3.0	3.0	4.3	3.7	3.5	3.2	3.6	3.0	2.4	4.2
3.6	4.3	2.9	2.8	4.0	3.4	3.2	3.1	3.3	2.9	2.4	3.2
.3	.3	.1	.2	.3	.4	.3	.1	.3	.1	0	.8
0	0	0	0	.1	(2)	0	0	0	0	0	.2
1.1	1.1	1.0	1.1	1.1	1.1	1.1	1.1	1.1	1.0	1.0	1.3
.3	.3	.2	.3	.4	.4	.4	.3	.3	.2	.0	.6
2.0	2.0	2.0	2.0	3.0	3.0	3.0	3.0	2.0	2.0	1.0	3.0
52	45	64	58	63	52	72	47	44	41	49	54
35	34	60	51	45	36	58	31	30	28	31	30
13	13			13	13	13	7		7		
27	27			27	27	40	27		27		
53	27			53	53	53	53		27		
53	53			80	53	80	53		53		
107	107			107	107	133	107		57		
107	107			160	120	187	107		107		
213	187			213	160	373	160		160		
152	124	171	168	165	144	186	140	118	122	148	144
107	87	124	155	105	97	124	91	82	86	92	87
96	142	520	139	205	126	252	93	129	60	0	105
38	72	0	42	75	72	150	47	39	0	0	42
0	0	0	0	171	80	0	0	0	0	0	80
0	0	0	0	47	0	0	0	0	0	0	0
640	480	720	800	480	480	800	480	480	480	320	320
640	560	1,040	800	640	640	1,120	480	480	480	320	320
142	126	186	159	168	136	193	133	120	118	148	121
103	85	148	122	103	79	118	88	81	85	92	75
40	40			40	40	40	20		20		
80	80			80	80	113	80		80		
120	80			160	120	160	150		80		
160	160			240	160	240	160		160		
320	240			320	240	320	240		170		
320	320			320	320	400	320		320		
640	480			480	320	640	480		480		
640	480	720	480	480	360	640	480	480	480	320	320
0	0	0	0	1.1	2.6	1.0	0	0	0	2.1	0
0	0	0	0	.6	1.1	.5	0	0	0	16.4	0

TABLE 5.38.--MACARONI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.5	.4	3.7	3.3	2.8
ON ONLY 1 OF 3 DAYS.....PCT.....	2.3	.4	3.2	2.9	2.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.2	0	.5	.5	.1
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.0	1.2	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.4	.4	.3
MAXIMUM.....DO.....	6.0	1.0	2.0	3.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	49	15	37	37	39
STANDARD DEVIATION.....GM.....	37	7	28	26	30
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	12			12	6
25TH.....GM.....	23			23	23
50TH.....GM.....	47			23	23
75TH.....GM.....	58			47	47
90TH.....GM.....	93			70	93
95TH.....GM.....	104			93	103
99TH.....GM.....	187			140	140
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	134	46	101	98	111
STANDARD DEVIATION.....GM.....	94	22	81	67	84
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	139	0	82	100	135
STANDARD DEVIATION.....GM.....	91	0	41	43	24
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	129	0	0	0	0
STANDARD DEVIATION.....GM.....	69	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	700	70	490	420	420
MAXIMUM OVER 3 DAYS.....GM.....	980	70	490	420	420
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	131	46	93	96	109
STANDARD DEVIATION.....GM.....	93	22	67	57	73
PERCENTILES: <u>1/</u>					
5TH.....GM.....	35			35	18
25TH.....GM.....	70			70	70
50TH.....GM.....	140			70	70
75TH.....GM.....	140			140	140
90TH.....GM.....	280			140	210
95TH.....GM.....	280			210	280
99TH.....GM.....	420			280	420
MAXIMUM.....GM.....	700	70	350	310	420
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	5.8	0	0	0	.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	0	1.4

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.4	3.1	1.8	2.8	2.4	2.7	2.5	2.2	1.6	1.3	2.4	.9
3.3	2.9	1.7	2.7	2.2	2.5	2.3	1.9	1.5	1.1	2.4	.9
.1	.2	.1	.1	.2	.3	.2	.2	0	.2	0	0
0	0	0	0	0	0	0	(2)	.1	0	0	0
1.0	1.1	1.1	1.0	1.1	1.1	1.1	1.2	1.4	1.2	1.0	1.0
.2	.2	.3	.2	.3	.3	.3	.4	1.3	.4	.0	.0
2.0	2.0	2.0	2.0	2.0	2.0	2.0	3.0	6.0	2.0	1.0	1.0
54	44	57	52	63	51	53	45	46	43	47	43
41	28	48	33	42	45	33	38	31	32	26	16
12	6			6	17	12	12				
23	23			47	23	23	23				
47	47			47	47	47	35				
93	47			93	47	93	47				
93	93			98	93	93	93				
140	93			140	140	93	93				
187	105			187	327	187	233				
153	124	155	155	166	129	158	117	132	112	140	129
112	79	127	100	97	92	99	83	95	73	78	47
247	127	181	105	204	185	98	117	0	118	0	0
35	29	96	0	75	121	51	91	0	69	0	0
0	0	0	0	0	0	0	210	70	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
560	280	700	560	420	560	560	560	420	280	280	210
560	315	700	560	560	980	560	700	420	350	280	210
156	124	159	149	168	139	144	115	100	108	140	129
107	76	124	96	100	101	93	87	90	74	78	47
35	18			17	52	35	26				
70	70			105	70	70	70				
140	140			140	140	140	85				
280	140			280	140	210	140				
280	280			280	280	280	210				
420	280			310	420	280	280				
420	280			420	560	560	560				
420	280	700	560	420	560	560	560	420	280	280	210
0	0	0	0	0	1.4	1.4	2.2	0	0	0	0
0	0	0	0	0	.9	1.0	1.2	0	0	0	0

TABLE 5.39.--SPAGHETTI WITH TOMATO SAUCE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.6	4.0	16.8	17.2	17.9
ON ONLY 1 OF 3 DAYS.....PCT.....	9.6	3.1	14.4	14.8	16.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	.9	2.3	2.2	1.4
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.2	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.1	1.2	1.2	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.4	.4	.4	.4
MAXIMUM.....DO.....	4.0	2.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	132	69	71	90	98
STANDARD DEVIATION.....GM.....	93	32	44	53	64
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	42		21	21	42
25TH.....GM.....	83		42	42	63
50TH.....GM.....	104		63	83	83
75TH.....GM.....	167		83	125	125
90TH.....GM.....	250		146	167	167
95TH.....GM.....	333		167	167	208
99TH.....GM.....	458		167	250	333
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	361	178	194	240	261
STANDARD DEVIATION.....GM.....	233	85	120	136	143
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	355	153	161	225	321
STANDARD DEVIATION.....GM.....	202	26	74	81	137
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	273	0	125	195	260
STANDARD DEVIATION.....GM.....	196	0	0	39	15
MAXIMUM IN ANY 1 DAY.....GM.....	2,500	250	750	1,000	1,500
MAXIMUM OVER 3 DAYS.....GM.....	4,500	375	750	1,000	1,500
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	351	169	184	227	264
STANDARD DEVIATION.....GM.....	225	75	104	121	142
PERCENTILES: <u>1/</u>					
5TH.....GM.....	95		47	63	125
25TH.....GM.....	250		125	125	188
50TH.....GM.....	250		155	250	250
75TH.....GM.....	500		250	250	250
90TH.....GM.....	500		250	375	500
95TH.....GM.....	751		375	500	500
99TH.....GM.....	1,001		500	500	750
MAXIMUM.....GM.....	2,500	250	500	1,000	1,500
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	67.5	0	2.1	6.1	2.9
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	1.0	1.7	.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.1	17.4	13.2	12.6	8.9	9.9	8.8	7.5	5.2	5.6	4.4	3.4
16.8	15.6	12.0	11.5	8.1	9.2	8.2	6.9	4.3	4.7	2.8	2.8
2.1	1.6	1.1	1.1	.6	.7	.6	.6	.8	.8	1.6	.6
.1	.1	.1	0	.1	.1	0	(2)	.1	.2	0	0
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.3	1.4	1.2
.4	.4	.4	.3	.4	.3	.4	.3	.5	.7	.5	.4
3.0	3.0	3.0	3.0	3.0	3.0	4.0	4.0	3.0	4.0	2.0	2.0
145	118	167	129	182	133	164	125	155	127	152	149
119	76	111	89	107	76	101	88	93	89	97	61
42	32	42	42	63	42	42	42	38	42		
83	83	83	83	83	83	83	83	83	83		
125	83	167	83	167	125	167	83	139	83		
167	167	167	167	250	167	208	167	250	167		
250	208	292	250	333	250	292	208	292	250		
333	250	417	258	375	250	333	319	333	333		
500	417	584	500	583	417	500	417	417	417		
380	323	460	342	500	372	470	337	406	328	374	390
243	198	291	200	278	192	297	191	250	199	147	127
432	317	452	422	523	351	403	387	377	377	300	353
343	138	220	203	163	170	97	165	119	160	201	87
212	253	333	0	439	317	0	1,167	208	42	0	0
44	47	0	0	13	62	0	0	0	0	0	0
2,500	1,500	2,000	1,750	2,000	1,250	2,500	2,000	1,000	1,250	750	500
4,500	1,500	2,000	2,000	2,000	1,500	2,500	3,500	1,250	1,250	1,000	1,000
381	314	446	348	491	363	442	342	375	298	334	378
268	185	280	198	258	190	279	192	198	189	196	121
125	95	125	125	188	125	125	125	125	31		
250	250	250	250	250	250	250	250	250	125		
250	250	500	250	500	375	375	250	375	250		
500	375	500	500	563	500	500	500	500	470		
625	500	751	500	751	500	751	500	750	500		
751	625	1,000	750	1,000	750	1,000	750	751	750		
1,501	1,000	1,501	1,001	1,500	1,000	1,501	1,124	1,000	750		
2,500	1,500	2,000	1,500	2,000	1,250	2,500	1,500	1,000	750	750	500
3.5	2.7	1.9	1.4	5.7	19.0	13.5	5.1	2.0	1.6	0	0
.8	.6	.9	.7	1.5	3.3	2.8	.9	2.8	1.3	0	0

TABLE 5.40.--MACARONI AND CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.7	3.0	14.5	14.7	13.5
ON ONLY 1 OF 3 DAYS.....PCT.....	7.9	2.5	12.8	13.5	11.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	.4	1.7	1.3	1.8
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.1	1.2	1.1	1.1	1.2
STANDARD DEVIATION.....DO.....	.3	.4	.3	.3	.4
MAXIMUM.....DO.....	3.0	2.0	2.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	76	37	46	58	69
STANDARD DEVIATION.....GM.....	55	19	31	47	48
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		5	10	17
25TH.....GM.....	33		25	33	33
50TH.....GM.....	67		40	40	67
75TH.....GM.....	83		67	80	80
90TH.....GM.....	133		80	100	133
95TH.....GM.....	167		100	160	160
99TH.....GM.....	267		160	267	240
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	209	107	132	154	190
STANDARD DEVIATION.....GM.....	139	53	85	100	129
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	199	70	95	195	144
STANDARD DEVIATION.....GM.....	125	31	62	132	86
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	208	0	0	0	136
STANDARD DEVIATION.....GM.....	106	0	0	0	67
MAXIMUM IN ANY 1 DAY.....GM.....	1,200	240	480	720	800
MAXIMUM OVER 3 DAYS.....GM.....	1,494	240	480	960	800
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	203	93	121	157	175
STANDARD DEVIATION.....GM.....	136	51	81	109	118
PERCENTILES: <u>1/</u>					
5TH.....GM.....	38		13	25	30
25TH.....GM.....	100		60	100	100
50TH.....GM.....	200		100	120	150
75TH.....GM.....	240		200	200	240
90TH.....GM.....	400		240	240	300
95TH.....GM.....	480		240	400	400
99TH.....GM.....	720		360	480	720
MAXIMUM.....GM.....	1,200	240	480	720	720
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	13.7	0	2.2	0	2.9
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	1.3	0	1.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
11.2	11.4	10.2	8.7	8.1	8.9	6.6	7.2	5.3	6.4	3.9	5.4
10.3	10.5	9.3	8.0	7.3	8.1	6.0	6.7	5.1	5.8	3.7	5.0
.9	.8	.7	.6	.8	.7	.6	.5	.2	.6	.2	.4
.1	(2)	.1	.1	0	.1	0	(2)	0	0	0	0
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
.3	.3	.4	.4	.3	.4	.3	.3	.2	.3	.4	.3
3.0	3.0	3.0	3.0	2.0	3.0	3.0	3.0	2.0	2.0	2.0	2.0
79	73	108	78	107	75	85	65	62	84	69	61
48	53	82	52	73	49	53	40	49	57	58	37
22	17	33	20	33	17	17	8	8	8		1
40	33	67	50	67	40	50	33	27	33		33
67	67	80	67	80	67	67	67	60	67		66
100	80	133	80	133	100	120	80	67	133		80
133	133	200	133	200	133	160	120	120	160		120
167	167	333	160	267	160	200	133	160	200		133
267	267	360	300	400	240	240	180	220	267		133
214	204	289	216	287	210	237	182	182	231	191	186
118	132	200	125	178	128	144	102	148	158	165	111
239	204	302	173	303	192	225	161	150	224	237	67
100	145	147	116	155	101	80	104	0	91	0	45
113	200	358	300	0	217	0	183	0	0	0	0
0	0	175	0	0	17	0	37	0	0	0	0
800	960	1,000	960	1,200	720	720	600	660	800	480	400
900	960	1,494	960	1,440	800	1,000	900	660	800	480	400
213	203	284	207	285	202	229	178	175	219	180	167
115	134	196	128	176	119	137	101	142	145	139	107
60	25	60	50	76	50	50	25	25	50		4
120	100	175	120	200	100	120	100	80	100		100
200	200	200	200	240	200	200	200	150	200		180
240	240	400	240	400	240	300	240	200	300		240
400	400	498	400	480	360	400	300	360	400		360
400	480	720	420	601	400	480	400	480	480		400
600	601	1,000	601	961	720	720	480	660	800		400
720	960	1,000	960	1,200	720	720	600	660	800	480	400
3.3	0	0	1.5	.9	0	1.5	1.4	0	0	0	0
1.3	0	0	1.1	.3	0	.4	.3	0	0	0	0

TABLE 5.41.--LASAGNE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.9	.2	1.6	1.5	2.8
ON ONLY 1 OF 3 DAYS.....PCT.....	1.8	0	1.2	1.5	2.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.1	.2	.3	0	.1
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.1	2.0	1.2	1.0	1.0
STANDARD DEVIATION.....DO.....	.3	.0	.4	.0	.2
MAXIMUM.....DO.....	3.0	2.0	2.0	1.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	121	83	56	56	82
STANDARD DEVIATION.....GM.....	96	0	46	24	39
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	28				34
25TH.....GM.....	63				63
50TH.....GM.....	83				83
75TH.....GM.....	167				83
90TH.....GM.....	230				136
95TH.....GM.....	272				167
99TH.....GM.....	584				230
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	319	0	101	167	235
STANDARD DEVIATION.....GM.....	195	0	46	71	100
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	420	125	208	0	279
STANDARD DEVIATION.....GM.....	266	0	26	0	58
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	514	0	0	0	0
STANDARD DEVIATION.....GM.....	125	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,632	125	250	250	500
MAXIMUM OVER 3 DAYS.....GM.....	2,040	250	500	250	689
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	333	125	139	167	238
STANDARD DEVIATION.....GM.....	218	0	68	71	100
PERCENTILES: <u>1/</u>					
5TH.....GM.....	85				102
25TH.....GM.....	188				188
50TH.....GM.....	250				250
75TH.....GM.....	500				250
90TH.....GM.....	510				375
95TH.....GM.....	751				500
99TH.....GM.....	1,001				500
MAXIMUM.....GM.....	1,632	125	250	250	500
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	19.8	0	2.0	2.0	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	2.5	0	10.1	7.7	3.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.9	2.6	2.8	2.7	2.8	1.9	1.7	1.4	.5	.7	1.4	.6
3.7	2.4	2.6	2.7	2.5	1.7	1.5	1.3	.3	.7	1.4	.6
.2	.2	.3	0	.3	.2	.2	.1	.2	0	0	0
(2)	0	0	0	.1	0	0	0	0	0	0	0
1.1	1.1	1.1	1.0	1.1	1.1	1.1	1.1	1.3	1.0	1.0	1.0
.3	.3	.3	.0	.4	.3	.3	.3	.5	.0	.0	.0
3.0	2.0	2.0	1.0	3.0	2.0	2.0	2.0	2.0	1.0	1.0	1.0
102	93	132	89	188	126	153	111	137	103	101	59
72	67	74	48	140	92	114	66	71	47	50	36
28	32			63	38	42	28				
63	42			104	83	83	63				
83	83			167	95	125	83				
125	95			230	166	167	167				
167	167			333	208	333	167				
272	250			596	255	408	250				
442	375			681	544	584	281				
280	237	376	267	454	320	390	320	460	310	304	176
168	129	215	144	217	163	239	184	248	140	149	109
387	389	294	0	679	425	524	244	156	0	0	0
191	114	105	0	318	231	240	150	0	0	0	0
323	0	0	0	595	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
816	750	1,250	750	1,632	1,000	1,250	816	750	500	500	250
1,326	1,125	1,250	750	2,040	1,632	1,750	1,000	750	500	500	250
289	259	355	267	498	332	416	305	308	310	304	176
171	145	218	144	287	174	245	180	241	140	149	109
85	95			188	113	125	85				
188	125			313	230	250	159				
250	250			500	250	375	250				
375	250			562	459	500	425				
500	500			817	500	817	500				
816	582			1,001	626	876	681				
816	750			1,632	875	1,250	816				
816	750	1,250	750	1,632	875	1,250	816	750	500	500	250
4.1	0	2.8	1.0	1.3	2.6	0	2.3	0	0	0	0
4.8	0	6.5	2.6	1.0	2.2	0	2.2	0	0	0	0

TABLE 5.42.--PIZZA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.8	.5	6.9	10.1	15.7
ON ONLY 1 OF 3 DAYS.....PCT.....	8.9	.5	6.4	9.4	14.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	0	.6	.6	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	(2)	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.1	1.0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.3	.6	.4
MAXIMUM.....DO.....	8.0	1.0	2.0	8.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	64	10	28	38	42
STANDARD DEVIATION.....GM.....	52	7	21	35	29
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	17		9	9	13
25TH.....GM.....	30		13	20	23
50TH.....GM.....	45		23	28	36
75TH.....GM.....	81		37	45	45
90TH.....GM.....	123		68	72	78
95TH.....GM.....	158		76	84	94
99TH.....GM.....	272		102	113	149
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	175	31	80	106	116
STANDARD DEVIATION.....GM.....	139	20	60	69	70
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	164	0	71	80	112
STANDARD DEVIATION.....GM.....	101	0	35	35	54
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	187	0	0	407	125
STANDARD DEVIATION.....GM.....	103	0	0	0	121
MAXIMUM IN ANY 1 DAY.....GM.....	1,945	57	305	477	461
MAXIMUM OVER 3 DAYS.....GM.....	1,945	57	305	1,222	840
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	171	31	78	102	115
STANDARD DEVIATION.....GM.....	133	20	59	62	70
PERCENTILES: ^{1/}					
5TH.....GM.....	43		26	28	38
25TH.....GM.....	85		34	60	61
50TH.....GM.....	136		68	85	108
75TH.....GM.....	227		91	136	136
90TH.....GM.....	338		128	181	216
95TH.....GM.....	408		213	227	255
99TH.....GM.....	639		305	268	377
MAXIMUM.....GM.....	1,945		305	457	461
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	1008.3	0	31.9	51.7	72.4
PROPORTION OF EATING OCCASIONS.....PCT.....	24.4	0	40.3	26.6	22.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.9	17.8	17.5	17.2	12.5	12.0	5.0	5.8	1.8	2.8	1.0	2.2
17.5	15.7	14.9	15.0	11.6	11.4	4.5	5.3	1.7	2.5	.8	2.0
2.2	1.9	2.3	2.0	.9	.6	.5	.4	.1	.2	.2	.2
.2	.2	.3	.1	(2)	(2)	0	(2)	0	0	0	0
1.2	1.2	1.2	1.2	1.1	1.1	1.1	1.1	1.0	1.1	1.2	1.1
.4	.5	.4	.5	.3	.3	.3	.3	.2	.3	.4	.3
4.0	5.0	3.0	7.0	3.0	4.0	3.0	3.0	2.0	2.0	2.0	2.0
61	52	84	57	93	62	87	55	59	50	62	60
44	40	68	39	74	44	59	37	42	30	13	35
19	14	20	18	20	20	22	14				
33	27	37	32	45	35	43	28				
45	40	69	45	76	54	81	45				
81	68	112	74	122	80	113	68				
122	102	163	102	170	113	181	110				
151	136	205	122	221	136	213	132				
221	204	321	181	362	279	288	181				
166	140	213	155	270	175	236	154	176	144	187	156
118	101	151	101	222	111	145	96	129	87	44	80
144	127	226	128	209	191	244	160	113	112	94	200
77	53	148	62	87	107	125	79	0	37	0	0
151	172	221	194	172	279	0	117	0	0	0	0
54	92	107	102	0	0	0	0	0	0	0	0
816	682	1,418	680	1,945	880	865	610	486	370	251	284
816	840	1,758	984	1,945	907	1,092	636	486	370	251	401
158	137	211	148	260	174	235	154	171	140	155	164
115	93	144	96	211	103	145	101	125	83	57	75
42	43	61	56	61	60	61	31				
81	69	108	85	122	94	122	84				
122	112	183	122	227	152	227	136				
216	170	272	183	338	227	305	204				
305	244	380	258	476	284	424	272				
377	324	525	340	567	377	510	340				
550	486	742	543	849	486	680	543				
816	682	851	680	1,945	785	865	610	486	370	251	284
92.7	107.2	65.7	79.7	111.0	196.0	67.2	120.6	1.7	5.9	0	4.6
19.4	24.2	22.6	27.4	21.0	28.4	24.7	27.3	8.0	11.3	0	19.2

TABLE 5.43.--RICE

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	18.5	4.3	20.9	22.2	23.4
ON ONLY 1 OF 3 DAYS.....PCT.....	13.4	3.4	14.0	15.3	17.2
ON ONLY 2 OF 3 DAYS.....PCT.....	3.6	.8	4.6	4.9	4.4
ON ALL 3 OF 3 DAYS.....PCT.....	1.5	.1	2.3	2.0	1.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.5	1.6	1.5	1.4
STANDARD DEVIATION.....DO.....	.9	1.0	1.0	.8	.8
MAXIMUM.....DO.....	9.0	6.0	7.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	71	26	42	47	57
STANDARD DEVIATION.....GM.....	73	22	38	42	52
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		5	7	11
25TH.....GM.....	29		15	19	29
50TH.....GM.....	58		29	29	29
75TH.....GM.....	88		58	58	58
90TH.....GM.....	146		97	117	117
95TH.....GM.....	229		131	117	175
99TH.....GM.....	350		190	204	233
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	57	76	87	108
STANDARD DEVIATION.....GM.....	84	52	49	57	76
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	169	69	102	105	167
STANDARD DEVIATION.....GM.....	103	20	65	63	100
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	208	87	95	124	120
STANDARD DEVIATION.....GM.....	129	0	47	57	53
MAXIMUM IN ANY 1 DAY.....GM.....	1,400	175	350	525	875
MAXIMUM OVER 3 DAYS.....GM.....	3,500	263	613	755	1,225
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	147	53	81	95	120
STANDARD DEVIATION.....GM.....	91	42	50	58	77
PERCENTILES: <u>1/</u>					
5TH.....GM.....	41		11	21	41
25TH.....GM.....	88		44	44	88
50TH.....GM.....	165		88	88	88
75TH.....GM.....	175		88	88	175
90TH.....GM.....	263		175	175	175
95TH.....GM.....	350		175	175	263
99TH.....GM.....	438		263	263	350
MAXIMUM.....GM.....	1,400	175	350	350	875
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	67.2	0	0	1.5	.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	.3	.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
18.9	22.4	20.9	19.0	18.5	18.3	17.7	18.4	15.4	14.1	18.8	12.0
13.9	16.7	15.4	14.1	13.2	13.0	13.3	13.1	10.4	10.7	14.6	8.6
4.3	3.6	3.2	3.1	3.6	3.7	3.0	3.9	3.1	2.3	2.8	2.8
.7	2.0	2.3	1.7	1.7	1.6	1.5	1.3	1.9	1.1	1.4	.6
1.4	1.4	1.5	1.4	1.5	1.5	1.4	1.5	1.6	1.5	1.6	1.5
.7	.9	.8	.8	.8	.9	.9	.8	1.0	1.0	1.2	.9
5.0	6.0	6.0	5.0	7.0	9.0	7.0	9.0	5.0	7.0	7.0	5.0
68	66	95	76	88	73	82	69	79	61	74	61
56	59	103	69	81	76	86	70	89	63	73	50
15	15	22	15	15	15	15	9	15	14	19	9
29	29	44	29	36	29	29	29	29	29	29	29
58	55	58	58	58	58	58	55	58	44	58	55
88	83	117	88	117	66	88	83	88	58	58	58
117	146	194	175	193	175	175	146	175	117	146	139
175	175	233	233	233	233	233	233	182	175	230	175
277	292	467	350	467	350	350	350	526	408	408	233
140	129	180	147	165	126	155	126	138	115	148	128
86	80	107	96	101	70	85	78	76	64	84	74
173	163	200	184	210	180	191	168	136	134	159	147
99	86	95	100	106	122	94	106	75	53	77	82
189	177	272	214	238	219	262	211	239	222	268	157
61	85	199	91	128	110	166	118	153	102	107	79
700	700	1,400	700	875	1,400	875	875	700	525	525	350
1,185	1,488	3,500	1,225	1,575	1,575	2,538	1,925	1,750	1,225	1,225	700
149	138	195	160	180	151	170	142	149	125	142	120
86	77	117	89	105	98	95	83	90	56	61	58
44	41	67	44	44	41	44	28	44	44	54	11
88	88	124	88	88	88	88	88	88	88	88	88
175	131	175	175	175	175	175	131	131	131	175	131
175	175	248	175	175	175	175	175	175	175	175	175
263	262	350	350	350	263	350	263	284	175	175	175
350	263	350	350	350	350	350	350	350	175	175	175
350	350	700	350	526	350	438	350	350	263	350	263
700	525	700	525	700	1,400	525	525	438	350	350	263
4.9	.8	2.2	4.2	5.3	14.5	13.0	8.8	3.1	2.0	2.8	3.6
.9	.1	.5	1.1	.5	1.0	1.0	.5	1.1	.6	1.8	2.0

TABLE 5.44.--COOKED CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.4	16.3	33.1	26.0	21.3
ON ONLY 1 OF 3 DAYS.....PCT.....	10.3	5.8	16.2	15.3	13.1
ON ONLY 2 OF 3 DAYS.....PCT.....	4.3	4.9	11.1	7.2	6.3
ON ALL 3 OF 3 DAYS.....PCT.....	2.7	5.5	5.8	3.6	1.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	2.1	1.7	1.6	1.5
STANDARD DEVIATION.....DO.....	.8	1.1	.8	.8	.7
MAXIMUM.....DO.....	6.0	6.0	5.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	107	82	85	92	98
STANDARD DEVIATION.....GM.....	81	74	64	68	66
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20	5	10	13	20
25TH.....GM.....	50	26	40	41	41
50TH.....GM.....	82	62	80	80	82
75TH.....GM.....	160	109	120	123	140
90TH.....GM.....	240	210	163	184	172
95TH.....GM.....	245	245	220	243	240
99TH.....GM.....	388	320	280	320	320
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	202	98	142	167	202
STANDARD DEVIATION.....GM.....	109	75	100	95	114
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	208	112	155	192	198
STANDARD DEVIATION.....GM.....	100	71	82	91	84
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	208	142	157	178	196
STANDARD DEVIATION.....GM.....	108	86	71	82	91
MAXIMUM IN ANY 1 DAY.....GM.....	1,200	480	660	735	960
MAXIMUM OVER 3 DAYS.....GM.....	2,640	960	980	980	1,470
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	203	116	149	177	198
STANDARD DEVIATION.....GM.....	110	82	87	97	104
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31	14	28	31	38
25TH.....GM.....	123	60	81	120	123
50TH.....GM.....	240	105	123	180	240
75TH.....GM.....	245	123	240	245	245
90TH.....GM.....	360	245	245	245	263
95TH.....GM.....	480	263	245	263	368
99TH.....GM.....	490	360	480	480	490
MAXIMUM.....GM.....	960	462	540	735	960
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	45.4	.7	3.7	3.7	3.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	.4	.6	.5	.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.5	17.3	14.3	12.1	10.5	12.2	16.0	15.0	26.4	24.7	33.1	33.6
12.2	11.9	9.8	9.2	6.8	9.0	8.4	9.5	13.4	13.1	12.3	15.8
5.1	4.0	2.6	1.7	2.7	2.5	4.2	3.1	5.3	6.8	10.3	8.1
2.2	1.5	1.9	1.1	1.1	.8	3.3	2.4	7.7	4.7	10.5	9.7
1.5	1.4	1.5	1.3	1.5	1.3	1.7	1.5	1.8	1.7	2.0	1.8
.7	.6	.7	.6	.7	.6	.8	.8	.9	.8	.8	.9
3.0	3.0	4.0	3.0	4.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0
111	99	127	102	118	88	130	97	135	108	140	120
78	70	107	67	91	63	93	70	92	84	102	88
20	30	31	40	20	17	21	15	40	21	28	20
60	54	80	80	73	41	80	41	80	41	80	54
82	82	82	82	82	80	88	80	99	82	120	82
160	123	160	123	162	88	163	123	163	160	200	163
220	163	283	163	240	163	245	179	263	240	243	242
280	240	351	245	286	245	323	245	327	245	286	245
360	368	490	360	490	325	480	327	480	480	551	435
223	217	246	238	232	190	230	188	222	184	220	183
125	99	111	114	126	100	128	93	91	97	93	79
244	211	247	204	237	217	237	195	231	189	234	183
121	86	129	74	101	110	102	89	76	85	93	98
193	206	308	233	277	188	228	191	226	213	201	215
98	133	152	100	139	77	116	86	103	106	136	87
960	960	1,200	735	735	960	789	526	490	608	551	490
1,440	1,920	2,640	1,470	1,960	1,103	2,160	1,440	1,470	1,470	1,654	1,450
223	212	259	229	242	197	230	188	224	192	213	198
126	107	132	106	126	104	122	94	102	100	119	95
35	43	61	110	61	43	41	31	60	46	28	29
123	123	184	123	161	123	123	123	123	123	123	123
240	240	245	240	245	240	240	220	240	184	240	240
245	245	306	245	245	245	245	245	245	245	245	245
368	360	490	368	480	263	368	245	360	245	360	245
490	420	490	480	490	368	490	360	480	368	480	360
612	490	527	490	700	490	601	490	490	490	551	490
960	960	960	735	735	960	789	526	490	490	551	490
2.6	4.9	1.8	2.0	2.6	2.8	8.2	5.7	1.8	0	0	1.3
.4	.9	.6	.8	.4	.3	.6	.4	.3	0	0	.2

TABLE 5.45.--GRITS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.7	2.4	9.2	8.4	5.3
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	1.3	3.9	5.3	3.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.4	.6	3.3	2.0	1.7
ON ALL 3 OF 3 DAYS.....PCT.....	.9	.5	2.0	1.1	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.7	1.9	1.5	1.5
STANDARD DEVIATION.....DO.....	.8	.9	.9	.8	.6
MAXIMUM.....DO.....	5.0	4.0	5.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	97	67	77	78	89
STANDARD DEVIATION.....GM.....	78	58	63	62	69
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		8	10	20
25TH.....GM.....	41		28	41	41
50TH.....GM.....	82		55	61	82
75TH.....GM.....	123		114	82	123
90TH.....GM.....	204		163	163	163
95TH.....GM.....	245		218	245	204
99TH.....GM.....	388		245	305	490
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	173	100	136	141	159
STANDARD DEVIATION.....GM.....	93	82	126	83	71
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	190	119	115	163	181
STANDARD DEVIATION.....GM.....	94	36	69	73	72
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	200	143	137	164	225
STANDARD DEVIATION.....GM.....	105	70	76	84	111
MAXIMUM IN ANY 1 DAY.....GM.....	980	245	660	490	490
MAXIMUM OVER 3 DAYS.....GM.....	1,960	735	980	980	1,470
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	184	117	123	152	179
STANDARD DEVIATION.....GM.....	103	76	84	87	87
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31		15	31	55
25TH.....GM.....	123		61	110	123
50TH.....GM.....	184		123	123	184
75TH.....GM.....	245		163	245	245
90TH.....GM.....	245		245	245	245
95TH.....GM.....	368		245	245	245
99TH.....GM.....	490		368	430	490
MAXIMUM.....GM.....	735	245	490	490	490
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	15.4	0	1.5	.8	1.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.8	.4	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.8	4.1	6.2	4.9	4.8	4.6	6.1	5.0	7.0	7.0	7.7	7.7
4.5	2.6	3.7	3.1	3.1	3.3	3.1	3.4	3.5	3.7	2.5	4.4
1.7	1.0	1.4	1.1	1.1	.9	1.8	.9	1.3	1.8	3.5	1.6
.6	.5	1.1	.8	.6	.3	1.2	.8	2.2	1.5	1.8	1.6
1.4	1.5	1.6	1.5	1.5	1.4	1.7	1.5	1.8	1.7	1.9	1.7
.7	.7	.8	.8	.7	.6	.8	.8	.9	.9	.7	.8
3.0	3.0	4.0	3.0	4.0	3.0	4.0	4.0	3.0	4.0	3.0	3.0
88	94	121	95	114	86	116	88	117	95	105	96
66	67	102	62	103	68	85	67	89	79	58	76
18	27	20	37	20	15	19	10	20	15		10
41	41	55	41	41	41	61	41	41	41		41
82	82	82	82	82	82	82	82	82	73		82
95	136	163	123	163	82	163	110	163	123		123
163	163	245	204	245	163	245	184	245	204		245
204	204	327	245	327	245	286	245	327	245		245
408	368	490	245	490	368	408	286	408	327		368
173	175	196	192	198	169	195	169	174	138	209	181
80	87	90	92	98	86	102	91	91	75	47	94
206	202	225	176	218	233	203	169	201	177	226	134
122	71	94	66	96	111	94	78	69	89	79	78
173	198	269	187	313	188	217	194	200	186	68	200
58	84	132	70	149	82	101	74	101	93	40	90
613	490	980	490	735	613	735	490	490	490	490	490
1,225	1,103	1,470	735	1,960	1,103	1,470	980	1,225	1,470	858	1,103
184	189	227	186	230	191	204	175	193	164	166	174
105	86	117	81	124	101	106	91	103	92	101	97
26	46	61	55	61	61	31	31	41	28		28
123	123	123	123	123	123	123	123	123	123		123
184	245	245	220	245	220	245	184	184	123		184
245	245	245	245	245	245	245	245	245	245		245
245	245	368	245	490	245	245	245	368	245		245
368	368	490	245	490	368	490	245	368	245		245
612	490	490	490	490	490	490	490	490	490		490
613	490	490	490	735	613	490	490	490	490	490	490
1.6	1.4	.8	2.0	0	0	2.1	2.0	1.8	0	0	0
.8	1.1	.6	1.8	0	0	.4	.4	1.3	0	0	0

TABLE 5.46.--OATMEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.3	9.0	18.1	13.1	10.9
ON ONLY 1 OF 3 DAYS.....PCT.....	5.7	3.4	11.5	9.0	7.5
ON ONLY 2 OF 3 DAYS.....PCT.....	1.7	3.3	4.8	3.0	2.8
ON ALL 3 OF 3 DAYS.....PCT.....	1.0	2.3	1.8	1.1	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	2.0	1.5	1.4	1.4
STANDARD DEVIATION.....DO.....	.7	.9	.7	.7	.6
MAXIMUM.....DO.....	4.0	4.0	3.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	102	79	79	87	95
STANDARD DEVIATION.....GM.....	76	82	59	62	64
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		13	20	26
25TH.....GM.....	54		40	40	40
50TH.....GM.....	80		80	80	80
75TH.....GM.....	120		100	97	120
90TH.....GM.....	200		160	160	172
95TH.....GM.....	240		195	240	240
99TH.....GM.....	360		295	320	335
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	216	82	154	184	212
STANDARD DEVIATION.....GM.....	112	51	89	97	118
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	209	106	174	192	206
STANDARD DEVIATION.....GM.....	99	70	91	101	92
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	215	166	154	184	196
STANDARD DEVIATION.....GM.....	117	106	76	77	112
MAXIMUM IN ANY 1 DAY.....GM.....	1,200	480	540	480	960
MAXIMUM OVER 3 DAYS.....GM.....	2,640	960	960	960	1,080
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	213	121	161	185	206
STANDARD DEVIATION.....GM.....	113	93	93	96	115
PERCENTILES: <u>1/</u>					
5TH.....GM.....	43		43	38	43
25TH.....GM.....	120		120	120	120
50TH.....GM.....	240		120	180	240
75TH.....GM.....	240		240	240	240
90TH.....GM.....	360		240	263	360
95TH.....GM.....	480		360	360	462
99TH.....GM.....	480		480	480	526
MAXIMUM.....GM.....	960	462	540	480	960
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	19.1	.7	0	2.9	2.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	.8	0	.9	.8

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
8.7	8.7	6.4	4.6	3.8	5.3	7.3	7.4	15.0	11.4	17.3	19.6
6.2	6.8	5.3	4.1	3.1	4.5	4.7	5.0	8.5	7.4	6.6	10.3
1.7	1.3	.8	.3	.4	.7	1.6	1.5	3.2	2.6	5.3	4.6
.8	.7	.3	.2	.3	.2	1.1	.8	3.3	1.4	5.3	4.7
1.4	1.3	1.2	1.2	1.3	1.2	1.5	1.4	1.7	1.5	2.0	1.7
.6	.6	.5	.5	.6	.5	.7	.7	.8	.7	.9	.8
3.0	3.0	4.0	3.0	3.0	3.0	3.0	4.0	3.0	3.0	3.0	3.0
115	94	115	102	106	79	126	93	129	104	149	115
80	75	97	65	62	55	93	64	82	85	88	79
19	30	40	40	40	19	20	19	40	26	40	28
80	53	80	80	80	40	80	40	80	54	80	54
80	80	80	80	80	80	80	80	88	80	140	80
160	100	120	120	120	80	160	120	160	120	240	160
220	160	165	160	200	160	240	160	240	200	240	240
263	200	240	240	240	170	280	200	280	240	240	240
408	400	480	360	320	320	480	360	480	480	480	320
255	223	272	253	260	198	253	196	252	200	275	195
135	96	109	102	113	105	138	96	82	85	116	86
245	199	256	275	319	218	238	204	228	185	213	200
123	103	148	72	102	131	86	69	87	77	84	96
239	228	400	302	182	154	256	185	221	260	226	207
112	172	227	60	73	78	138	103	105	126	101	78
960	960	1,200	480	720	960	789	526	480	480	480	480
1,440	1,920	2,640	1,080	960	1,080	2,160	1,440	1,440	1,440	1,440	1,306
249	218	282	262	254	198	250	195	233	211	228	202
131	119	137	98	120	113	129	95	100	103	101	89
35	35	88	120	60	35	60	43	60	60	43	60
180	120	240	240	180	120	180	120	180	120	180	120
240	240	240	240	240	240	240	240	240	240	240	240
263	240	360	263	277	240	263	240	240	240	240	240
480	360	480	480	480	263	480	263	360	360	360	240
480	480	526	480	480	360	480	360	480	480	480	346
527	480	721	480	480	480	721	480	480	480	480	480
960	960	960	480	720	960	789	526	480	480	480	480
1.0	1.5	1.0	0	0	2.8	6.1	1.0	0	0	0	0
.4	.6	.9	0	0	.8	1.1	.1	0	0	0	0

TABLE 5.47.--COOKED WHEAT CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.2	6.4	7.9	6.2	6.2
ON ONLY 1 OF 3 DAYS.....PCT.....	3.0	3.0	5.1	4.1	4.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	2.2	1.8	1.6	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.4	1.1	.9	.5	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.8	1.5	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.8	.7	.6	.6
MAXIMUM.....DO.....	5.0	5.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	99	65	78	94	88
STANDARD DEVIATION.....GM.....	71	55	59	70	51
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		10	20	20
25TH.....GM.....	54		41	41	41
50TH.....GM.....	82		61	82	82
75TH.....GM.....	123		96	116	102
90TH.....GM.....	163		163	184	163
95TH.....GM.....	245		204	245	194
99TH.....GM.....	327		245	327	255
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	221	102	140	183	211
STANDARD DEVIATION.....GM.....	112	65	75	89	101
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	202	101	163	196	214
STANDARD DEVIATION.....GM.....	98	68	55	104	82
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	217	143	201	245	159
STANDARD DEVIATION.....GM.....	113	55	45	50	63
MAXIMUM IN ANY 1 DAY.....GM.....	735	245	306	735	490
MAXIMUM OVER 3 DAYS.....GM.....	1,654	735	735	980	766
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	214	110	160	198	204
STANDARD DEVIATION.....GM.....	112	68	71	103	97
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31		31	57	38
25TH.....GM.....	123		123	123	123
50TH.....GM.....	245		163	245	245
75TH.....GM.....	245		245	245	245
90TH.....GM.....	368		245	245	245
95TH.....GM.....	490		245	368	368
99TH.....GM.....	490		245	490	490
MAXIMUM.....GM.....	735	245	306	735	490
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	10.9	0	2.2	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	1.8	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.9	5.3	2.4	2.7	2.5	2.5	3.3	3.2	5.8	8.3	9.7	9.5
3.4	4.4	1.9	2.4	2.0	1.7	2.2	2.4	4.2	5.6	6.0	5.5
1.3	.8	.3	.3	.4	.6	.7	.6	.7	1.6	1.1	2.6
.3	.1	.2	.1	.2	.2	.4	.3	.9	1.0	2.5	1.4
1.4	1.2	1.3	1.1	1.3	1.4	1.5	1.3	1.4	1.4	1.6	1.6
.6	.4	.6	.4	.6	.6	.8	.6	.7	.7	.9	.7
3.0	3.0	3.0	3.0	3.0	3.0	4.0	4.0	3.0	3.0	3.0	3.0
106	89	131	107	108	98	120	87	119	91	122	110
69	49	117	72	87	57	70	56	87	64	124	73
31	30			9	15	27	19	41	20	15	25
49	61			82	71	82	41	82	41	41	61
82	82			82	82	102	82	82	82	82	82
163	115			123	123	163	82	163	123	163	163
190	163			184	175	204	163	163	163	245	245
245	204			306	245	245	163	368	245	551	245
327	245			490	245	327	327	490	286	552	327
244	233	244	278	231	217	293	213	243	188	205	227
126	99	100	127	104	96	140	96	92	102	107	114
239	216	365	248	245	209	208	176	230	190	254	186
126	102	122	84	112	77	119	68	40	72	108	78
174	180	455	490	338	219	207	174	265	186	232	224
86	61	55	0	116	30	78	113	122	85	186	88
613	490	490	735	571	490	613	490	490	490	551	490
980	735	1,470	1,470	1,470	735	980	1,225	1,470	858	1,654	980
234	227	306	284	252	214	243	195	248	188	225	212
126	102	130	129	123	87	129	96	101	94	155	102
57	56			31	61	29	29	123	46	28	31
123	173			216	173	173	123	164	123	123	123
245	245			245	245	245	245	245	245	245	245
245	245			306	245	245	245	245	245	245	245
490	368			490	306	490	245	368	245	490	306
490	368			490	368	490	368	490	368	551	490
612	490			571	490	612	490	490	490	551	490
613	490	490	735	571	490	613	490	490	490	551	490
0	2.0	0	0	2.6	0	0	2.8	0	0	0	1.3
0	1.5	0	0	2.1	0	0	.9	0	0	0	.9

TABLE 5.48.--READY-TO-EAT CEREALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	43.4	68.7	68.0	75.8	76.8
ON ONLY 1 OF 3 DAYS.....PCT.....	17.4	7.5	23.5	25.5	24.2
ON ONLY 2 OF 3 DAYS.....PCT.....	13.5	10.4	23.8	27.0	26.7
ON ALL 3 OF 3 DAYS.....PCT.....	12.4	50.7	20.8	23.3	25.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	2.0	3.8	2.1	2.1	2.1
STANDARD DEVIATION.....DO.....	1.0	2.1	1.0	1.0	.9
MAXIMUM.....DO.....	15.0	15.0	9.0	10.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	23	17	16	20	23
STANDARD DEVIATION.....GM.....	20	15	12	14	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4	2	3	5	5
25TH.....GM.....	10	7	8	9	10
50TH.....GM.....	18	13	13	17	19
75TH.....GM.....	30	20	21	27	30
90TH.....GM.....	46	40	31	38	43
95TH.....GM.....	58	43	38	47	54
99TH.....GM.....	100	66	55	67	78
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	37	18	22	30	34
STANDARD DEVIATION.....GM.....	25	11	14	17	18
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	36	18	24	30	32
STANDARD DEVIATION.....GM.....	22	12	13	16	18
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	38	19	25	32	35
STANDARD DEVIATION.....GM.....	25	16	13	16	18
MAXIMUM IN ANY 1 DAY.....GM.....	495	158	150	200	224
MAXIMUM OVER 3 DAYS.....GM.....	935	342	261	400	383
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	36	13	23	29	33
STANDARD DEVIATION.....GM.....	25	11	14	17	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	8	2	6	10	9
25TH.....GM.....	22	5	13	19	25
50TH.....GM.....	29	10	21	27	28
75TH.....GM.....	45	20	28	35	40
90TH.....GM.....	60	28	40	50	56
95TH.....GM.....	84	40	50	60	65
99TH.....GM.....	120	46	70	90	100
MAXIMUM.....GM.....	440	80	110	170	224
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	130.9	21.5	8.7	4.8	7.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	1.7	.6	.2	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
69.8	64.0	50.4	43.7	28.8	30.5	30.3	30.4	47.5	46.6	48.2	47.4
24.0	25.5	19.5	22.7	14.1	16.5	12.8	14.6	14.4	16.5	15.3	13.9
24.3	21.4	15.0	13.0	8.5	9.5	8.4	8.8	13.4	14.8	16.0	13.5
21.6	17.2	15.9	8.0	6.2	4.4	9.1	7.0	19.7	15.4	17.0	20.0
2.1	1.9	2.0	1.7	1.8	1.6	1.9	1.8	2.2	2.0	2.1	2.2
1.0	.9	1.0	.8	.9	.8	.9	.9	1.0	.9	1.0	1.0
7.0	6.0	6.0	6.0	6.0	5.0	5.0	6.0	6.0	6.0	6.0	7.0
28	23	33	21	29	20	27	20	27	22	24	24
24	17	27	16	27	16	22	16	23	17	18	20
6	5	7	5	5	4	4	4	7	4	5	4
13	10	15	10	12	9	11	9	12	10	12	10
23	19	27	17	20	17	20	17	22	17	20	20
37	30	45	28	37	25	33	27	35	28	30	30
53	43	67	41	57	38	54	40	50	43	42	43
67	51	83	50	80	52	70	50	65	52	57	52
127	85	129	79	136	81	100	84	133	100	110	83
41	37	51	38	48	36	40	34	36	32	33	31
28	24	43	22	33	24	26	22	19	20	18	18
42	36	47	37	44	37	43	35	35	33	34	32
26	19	27	20	29	22	24	20	20	20	18	18
46	37	55	38	58	40	44	35	41	34	37	34
29	21	30	20	37	20	27	20	28	20	22	24
440	275	495	220	320	220	226	220	240	180	168	300
935	500	688	385	581	352	678	380	530	398	371	758
41	36	49	37	49	37	42	34	37	32	34	32
28	21	31	22	35	23	28	22	25	21	22	20
13	11	15	13	10	10	10	10	10	8	9	8
25	25	28	25	25	25	25	21	25	20	20	21
35	30	43	30	42	30	37	28	30	28	28	28
50	45	60	45	60	50	50	45	50	40	40	38
74	57	86	60	90	60	75	60	68	55	57	54
90	75	110	80	113	83	100	75	84	60	70	64
128	111	167	110	172	112	131	112	120	120	120	113
440	226	330	220	280	220	226	220	224	180	168	150
7.2	3.0	8.2	0	24.9	10.2	12.0	11.1	3.4	3.3	1.5	3.7
.2	.1	.6	0	1.2	.4	.4	.3	.3	.2	.3	.4

TABLE 5.49.--READY-TO-EAT CEREALS, 15-21 GM PER CUP, ALL SUGAR LEVELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.6	.3	3.4	4.7	4.9
ON ONLY 1 OF 3 DAYS.....PCT.....	2.2	.3	2.5	3.6	3.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	0	.7	.8	1.1
ON ALL 3 OF 3 DAYS.....PCT.....	.5	0	.2	.3	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.3	1.4	1.4
STANDARD DEVIATION.....DO.....	.8	.0	.6	.7	.7
MAXIMUM.....DO.....	5.0	1.0	3.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	11	2	8	9	9
STANDARD DEVIATION.....GM.....	7	0	4	6	6
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3			2	2
25TH.....GM.....	5			5	5
50TH.....GM.....	8			7	7
75TH.....GM.....	13			11	13
90TH.....GM.....	20			17	19
95TH.....GM.....	25			23	21
99TH.....GM.....	38			30	35
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	7	19	20	21
STANDARD DEVIATION.....GM.....	12	0	10	11	11
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	0	15	19	15
STANDARD DEVIATION.....GM.....	10	0	6	9	9
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	0	14	21	22
STANDARD DEVIATION.....GM.....	9	0	6	7	6
MAXIMUM IN ANY 1 DAY.....GM.....	84	8	40	57	63
MAXIMUM OVER 3 DAYS.....GM.....	180	8	60	90	105
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	20	7	17	19	19
STANDARD DEVIATION.....GM.....	11	0	9	10	11
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5			4	5
25TH.....GM.....	15			11	11
50TH.....GM.....	20			20	20
75TH.....GM.....	21			23	21
90TH.....GM.....	40			32	32
95TH.....GM.....	42			42	42
99TH.....GM.....	56			43	45
MAXIMUM.....GM.....	80	8	40	43	63
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2.5	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.5	4.7	4.2	2.8	1.4	2.0	2.7	3.2	6.9	7.8	8.5	6.5
3.6	3.0	2.7	1.8	.8	1.5	1.5	2.0	3.3	3.6	4.6	2.6
1.3	1.3	1.0	1.0	.5	.5	.7	.7	1.8	2.3	2.0	1.6
.7	.5	.5	0	.1	0	.6	.5	1.7	1.9	1.9	2.2
1.5	1.5	1.5	1.4	1.4	1.3	1.7	1.5	1.9	1.8	1.7	1.9
.7	.7	.7	.5	.6	.4	.8	.8	1.0	.9	.8	.9
4.0	3.0	3.0	2.0	3.0	2.0	3.0	3.0	5.0	4.0	3.0	3.0
13	11	14	11	11	9	12	10	12	11	10	11
9	7	9	5	9	5	8	7	8	8	9	7
3	3	4		3	2	3	2	3	3		3
7	7	7		5	7	7	5	7	5		5
10	10	13		9	7	10	7	10	8		8
18	14	20		14	13	13	13	15	15		16
25	20	27		20	14	20	19	25	20		20
28	20	30		28	19	28	20	28	27		20
40	30	47		53	28	40	38	37	40		40
27	24	29	29	21	22	20	18	22	16	15	16
14	12	17	14	11	12	11	8	8	8	8	4
22	20	27	18	27	21	20	22	19	19	15	19
12	9	9	8	18	9	11	9	11	7	6	4
27	23	29	0	21	0	22	18	20	19	23	17
11	8	10	0	2	0	8	7	8	9	10	8
84	48	80	63	80	60	80	57	45	57	40	40
180	126	140	70	160	85	120	120	112	120	120	120
25	22	28	24	24	21	21	19	19	18	18	17
12	11	14	12	15	10	10	8	10	9	9	7
5	5	15		8	7	8	8	5	5		7
16	16	20		15	15	15	15	15	10		11
21	20	21		20	20	20	20	20	20		20
40	30	40		28	23	23	20	20	20		20
42	42	42		42	40	30	28	36	30		28
42	42	50		43	43	40	30	42	40		30
60	43	80		80	43	56	56	42	40		40
63	43	80	63	80	43	80	57	42	57	40	40
0	0	1.0	0	0	0	0	0	0	0	1.5	0
0	0	1.2	0	0	0	0	0	0	0	1.9	0

TABLE 5.50.--READY-TO-EAT CEREALS, 25-32 GM PER CUP, 30 PCT OR MORE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.1	1.1	18.5	23.3	22.3
ON ONLY 1 OF 3 DAYS.....PCT.....	3.5	.5	10.2	13.5	10.6
ON ONLY 2 OF 3 DAYS.....PCT.....	1.7	.6	4.9	5.6	7.8
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	3.5	4.2	3.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.6	1.7	1.7	1.8
STANDARD DEVIATION.....DO.....	.8	.5	1.0	.9	.8
MAXIMUM.....DO.....	6.0	2.0	6.0	6.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	17	6	13	15	18
STANDARD DEVIATION.....GM.....	13	4	10	12	12
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		3	4	5
25TH.....GM.....	9		5	9	9
50TH.....GM.....	14		9	10	14
75TH.....GM.....	20		18	20	25
90TH.....GM.....	33		21	30	37
95TH.....GM.....	40		28	37	43
99TH.....GM.....	60		60	60	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	32	4	22	26	30
STANDARD DEVIATION.....GM.....	18	3	12	12	16
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	33	14	22	30	33
STANDARD DEVIATION.....GM.....	18	3	9	14	17
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	33	0	25	31	32
STANDARD DEVIATION.....GM.....	15	0	15	15	13
MAXIMUM IN ANY 1 DAY.....GM.....	168	28	96	150	168
MAXIMUM OVER 3 DAYS.....GM.....	336	42	244	276	224
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	31	12	22	28	31
STANDARD DEVIATION.....GM.....	17	6	13	14	15
PERCENTILES: <u>1/</u>					
5TH.....GM.....	8		7	8	8
25TH.....GM.....	21		14	16	25
50TH.....GM.....	28		16	28	28
75TH.....GM.....	32		28	30	32
90TH.....GM.....	56		30	45	56
95TH.....GM.....	60		56	57	57
99TH.....GM.....	90		64	90	64
MAXIMUM.....GM.....	168	28	96	96	112
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	3.4	0	.9	.9	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	.3	.1	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.5	16.9	7.6	7.8	2.3	2.1	.8	.9	.4	1.4	1.3	1.6
10.4	9.3	4.0	5.4	1.6	1.6	.5	.6	.4	.8	.8	1.0
4.9	4.9	2.2	1.5	.5	.5	.2	.2	0	.3	.2	.3
2.2	2.7	1.3	.9	.2	.1	.1	.1	0	.3	.3	.3
1.6	1.6	1.7	1.4	1.4	1.3	1.4	1.4	1.0	1.7	1.5	1.6
.8	.8	.8	.8	.6	.5	.6	.7	.0	.9	.8	.8
4.0	5.0	6.0	6.0	3.0	3.0	3.0	3.0	1.0	3.0	3.0	3.0
19	18	25	15	18	14	14	13	11	16	12	14
14	12	18	9	17	9	8	8	2	9	4	12
5	5	5	5	3	4		2				
9	9	9	9	9	9		9				
19	14	20	10	10	10		10				
24	21	35	20	20	19		17				
37	33	56	28	30	28		28				
47	40	59	30	56	33		30				
63	56	84	56	103	37		37				
39	33	44	33	33	33	37	30	33	30	38	23
23	17	25	14	21	17	17	10	7	15	13	14
38	32	44	28	44	34	27	26	0	28	14	15
23	13	20	11	38	14	15	10	0	13	0	(2)
36	34	51	32	43	33	9	25	0	27	9	36
14	13	19	10	16	5	4	13	0	3	0	7
168	120	126	84	168	84	56	56	42	60	56	57
336	252	294	176	308	112	112	112	42	90	56	126
36	32	45	31	38	33	28	28	33	27	22	26
20	15	22	13	28	15	18	12	7	11	16	13
14	11	14	14	8	11		6				
28	28	28	28	24	28		28				
28	28	42	28	30	28		28				
45	32	60	30	53	42		30				
60	56	70	56	75	56		42				
64	60	84	60	90	60		45				
112	84	120	84	168	84		56				
168	120	120	84	168	84	56	56	42	45	56	57
0	0	0	0	0	1.6	0	0	0	0	0	0
0	0	0	0	0	1.1	0	0	0	0	0	0

TABLE 5.51.--READY-TO-EAT CEREALS, 25-32 GM PER CUP, LESS THAN 15 PCT SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	19.4	6.2	31.8	36.1	35.1
ON ONLY 1 OF 3 DAYS.....PCT.....	10.9	2.0	16.6	18.4	17.4
ON ONLY 2 OF 3 DAYS.....PCT.....	5.5	2.7	10.0	12.0	11.3
ON ALL 3 OF 3 DAYS.....PCT.....	3.1	1.5	5.2	5.7	6.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	2.0	1.7	1.7	1.7
STANDARD DEVIATION.....DO.....	.8	.9	.8	.9	.8
MAXIMUM.....DO.....	10.0	4.0	6.0	10.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	16	9	11	14	16
STANDARD DEVIATION.....GM.....	12	9	8	10	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4		2	4	4
25TH.....GM.....	8		5	8	8
50TH.....GM.....	13		8	10	13
75TH.....GM.....	20		15	17	20
90TH.....GM.....	30		21	25	30
95TH.....GM.....	38		27	33	38
99TH.....GM.....	58		33	46	50
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	13	19	25	27
STANDARD DEVIATION.....GM.....	16	8	11	12	12
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	13	20	24	28
STANDARD DEVIATION.....GM.....	15	11	9	11	15
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	31	15	20	27	28
STANDARD DEVIATION.....GM.....	16	13	10	14	11
MAXIMUM IN ANY 1 DAY.....GM.....	275	60	75	200	125
MAXIMUM OVER 3 DAYS.....GM.....	510	120	180	400	200
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	29	13	19	24	27
STANDARD DEVIATION.....GM.....	16	13	10	12	13
PERCENTILES: <u>1/</u>					
5TH.....GM.....	8		6	10	8
25TH.....GM.....	25		13	15	25
50TH.....GM.....	25		15	25	25
75TH.....GM.....	30		25	30	30
90TH.....GM.....	50		30	38	50
95TH.....GM.....	57		30	50	50
99TH.....GM.....	85		50	60	75
MAXIMUM.....GM.....	275	60	75	112	100
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	52.1	0	4.0	1.6	5.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	.7	.1	.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
34.0	28.6	24.5	19.1	12.5	13.4	13.8	13.1	24.0	20.0	25.2	21.8
17.9	16.4	13.8	11.9	7.3	9.0	7.7	8.4	10.6	10.3	9.5	11.3
11.1	8.4	5.6	5.2	3.4	3.5	2.9	3.3	6.9	5.8	9.0	5.9
5.0	3.8	5.1	2.0	1.8	.9	3.2	1.4	6.5	3.9	6.7	4.6
1.7	1.6	1.7	1.5	1.6	1.4	1.7	1.5	1.9	1.7	2.0	1.8
.8	.8	.9	.8	.7	.6	.8	.7	.9	.8	1.0	.9
6.0	5.0	6.0	5.0	4.0	5.0	4.0	5.0	4.0	4.0	5.0	7.0
19	15	23	15	19	13	18	13	17	14	18	14
15	10	19	10	14	8	15	9	11	11	12	10
4	4	5	4	4	4	4	2	4	4	5	3
8	8	9	8	8	8	8	8	8	8	8	8
17	13	17	13	17	10	13	9	15	13	15	10
25	20	30	20	25	17	24	17	25	18	25	18
38	27	50	30	38	21	33	25	33	25	30	25
50	33	53	33	50	27	45	30	35	30	45	30
67	50	95	49	70	42	70	45	50	50	60	60
33	30	39	32	34	29	29	26	28	25	27	25
23	14	25	16	18	14	17	13	11	12	11	12
36	29	39	32	37	28	31	27	26	25	30	23
18	16	17	14	20	13	11	12	12	7	13	11
38	29	46	27	39	28	33	26	30	27	28	26
19	9	23	10	17	8	20	12	10	16	14	13
275	180	188	125	168	170	180	120	75	170	125	90
375	330	510	150	300	270	380	240	180	365	200	210
35	29	40	30	36	28	31	26	28	26	27	23
20	13	21	14	18	12	18	13	12	13	14	11
11	10	13	13	11	10	6	6	11	8	13	6
25	25	25	25	25	25	25	19	25	19	19	15
30	25	30	25	30	25	28	25	25	25	25	25
45	30	50	38	50	30	38	30	30	30	30	30
60	50	60	50	60	50	50	45	50	38	45	30
60	50	75	57	75	50	60	50	50	47	50	38
100	75	120	84	100	60	120	60	60	60	75	60
275	100	150	99	120	125	180	100	75	170	125	60
2.9	.9	1.7	0	16.6	4.2	10.2	2.9	0	2.0	0	0
.2	.1	.3	0	2.2	.4	.9	.2	0	.3	0	0

TABLE 5.52.--READY-TO-EAT CEREALS, 35-45 GM PER CUP, 30 PCT OR MORE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.7	.7	11.6	15.5	16.8
ON ONLY 1 OF 3 DAYS.....PCT.....	3.5	.7	6.8	9.3	9.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.5	0	2.7	4.6	5.0
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	2.1	1.5	2.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.7	1.5	1.6
STANDARD DEVIATION.....DO.....	.8	.0	.9	.7	.8
MAXIMUM.....DO.....	7.0	1.0	5.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	8	16	18	21
STANDARD DEVIATION.....GM.....	16	5	11	11	14
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	6		3	5	6
25TH.....GM.....	12		7	12	12
50TH.....GM.....	15		12	13	18
75TH.....GM.....	27		23	25	26
90TH.....GM.....	42		33	37	40
95TH.....GM.....	53		40	40	49
99TH.....GM.....	80		47	53	74
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	43	24	29	36	40
STANDARD DEVIATION.....GM.....	24	14	23	19	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	43	0	29	35	40
STANDARD DEVIATION.....GM.....	22	0	11	15	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	0	31	38	37
STANDARD DEVIATION.....GM.....	22	0	10	9	16
MAXIMUM IN ANY 1 DAY.....GM.....	320	37	140	140	148
MAXIMUM OVER 3 DAYS.....GM.....	520	37	148	200	240
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	42	24	29	36	39
STANDARD DEVIATION.....GM.....	23	14	15	16	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	12		9	12	10
25TH.....GM.....	28		18	23	28
50TH.....GM.....	37		28	35	37
75TH.....GM.....	50		37	40	40
90TH.....GM.....	74		40	57	70
95TH.....GM.....	80		53	70	80
99TH.....GM.....	120		80	80	111
MAXIMUM.....GM.....	175	37	90	120	148
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2.2	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.4	13.2	10.0	8.5	3.7	2.7	1.3	1.5	1.9	1.9	1.3	2.0
10.5	8.0	6.0	5.6	2.2	2.0	.8	1.1	1.1	1.1	.3	1.1
4.7	3.4	2.7	2.0	1.0	.6	.3	.2	.6	.5	.8	.3
2.2	1.8	1.3	.9	.5	.1	.2	.2	.2	.3	.2	.6
1.6	1.6	1.6	1.5	1.5	1.3	1.6	1.5	1.7	1.6	1.9	1.7
.8	.8	.8	.7	.8	.6	.8	.8	.9	.8	.6	.9
7.0	5.0	5.0	3.0	4.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
25	22	30	21	25	19	22	18	23	20	30	16
19	17	20	14	20	14	17	13	18	12	20	6
6	6	6	9	5	6	3	6				
12	12	13	12	12	12	12	12				
20	16	25	13	20	13	13	13				
30	27	41	26	30	23	27	23				
49	41	58	38	57	35	40	40				
58	53	67	53	67	49	70	43				
89	93	93	80	100	75	80	60				
48	42	65	44	50	45	40	38	31	33	35	41
23	20	39	18	28	24	23	14	8	8	0	14
48	45	53	42	52	38	38	39	54	39	40	40
23	25	27	22	31	18	13	14	16	11	0	0
52	44	56	44	46	57	50	40	45	42	80	16
31	23	21	16	28	19	21	14	24	3	0	5
320	170	225	140	170	180	100	80	80	60	80	80
520	383	320	280	300	225	240	200	240	140	240	80
47	43	57	43	49	44	42	38	39	38	47	29
23	23	31	19	32	21	21	14	15	10	16	15
14	13	18	18	5	13	10	19				
35	35	37	35	28	35	30	30				
40	37	53	40	40	40	40	40				
60	53	74	45	70	53	40	40				
80	70	90	74	80	70	80	60				
80	80	120	80	120	80	80	70				
111	140	167	120	170	120	100	80				
175	170	167	140	170	120	100	80	80	60	80	80
0	1.3	0	0	0	0	.9	0	0	0	0	0
0	.3	0	0	0	0	.9	0	0	0	0	0

TABLE 5.53.--READY-TO-EAT CEREALS, 35-45 GM PER CUP, LESS THAN 20 PCT SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.9	.8	3.4	4.8	4.7
ON ONLY 1 OF 3 DAYS.....PCT.....	2.8	.5	1.8	3.0	3.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.3	0	.8	1.3	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.8	.3	.7	.5	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.8	1.7	1.5	1.5
STANDARD DEVIATION.....DO.....	.8	1.0	.8	.7	.8
MAXIMUM.....DO.....	6.0	3.0	3.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	20	9	19	17	20
STANDARD DEVIATION.....GM.....	16	6	13	13	14
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4			3	5
25TH.....GM.....	12			9	12
50TH.....GM.....	15			13	17
75TH.....GM.....	27			25	22
90TH.....GM.....	38			29	37
95TH.....GM.....	47			39	55
99TH.....GM.....	83			62	72
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	39	14	34	33	43
STANDARD DEVIATION.....GM.....	22	4	21	18	23
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	38	0	30	34	35
STANDARD DEVIATION.....GM.....	19	0	12	18	11
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	39	16	35	36	46
STANDARD DEVIATION.....GM.....	23	5	12	16	22
MAXIMUM IN ANY 1 DAY.....GM.....	215	22	86	138	129
MAXIMUM OVER 3 DAYS.....GM.....	409	65	165	193	215
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	38	15	34	33	40
STANDARD DEVIATION.....GM.....	21	5	17	19	20
PERCENTILES: <u>1/</u>					
5TH.....GM.....	9			5	11
25TH.....GM.....	23			21	22
50TH.....GM.....	35			32	38
75TH.....GM.....	43			43	55
90TH.....GM.....	65			55	65
95TH.....GM.....	83			65	74
99TH.....GM.....	110			86	86
MAXIMUM.....GM.....	172	22	86	138	111
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	10.1	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.1	4.4	4.8	3.5	4.0	4.1	4.9	4.7	8.7	9.1	7.2	8.7
3.2	2.9	2.6	2.7	2.7	2.4	2.2	2.6	5.5	5.1	4.1	3.0
1.4	.9	1.2	.5	1.0	1.3	1.7	1.4	.8	1.8	2.2	2.5
.4	.6	1.0	.3	.3	.4	.9	.7	2.4	2.2	.9	3.1
1.5	1.5	1.7	1.3	1.4	1.5	1.8	1.6	1.6	1.7	1.6	2.1
.7	.8	.9	.6	.8	.7	.8	.7	.9	.9	.7	1.0
4.0	3.0	4.0	3.0	6.0	5.0	5.0	3.0	3.0	4.0	3.0	5.0
24	22	37	20	26	19	23	17	19	17	17	21
20	13	28	13	22	12	18	11	13	10	12	14
4	6	9	5	4	4	4	3	5	2		3
12	13	18	12	13	9	12	9	9	9		12
19	22	28	14	22	14	18	13	13	13		13
29	29	49	27	31	25	31	23	25	23		27
46	37	72	37	46	37	43	35	38	33		43
55	50	110	43	57	43	70	38	41	37		53
128	57	129	70	136	55	83	47	73	43		58
46	48	58	43	53	36	38	33	34	29	33	28
25	23	24	23	28	19	19	17	16	16	14	14
48	42	74	50	45	37	37	32	34	35	28	35
15	19	38	12	16	18	17	15	4	12	9	19
67	38	68	45	78	39	45	29	34	27	37	29
36	12	31	13	34	7	23	13	14	9	19	14
165	111	215	110	175	110	123	86	75	75	70	140
387	185	387	210	409	172	258	180	219	129	210	175
50	43	64	45	54	37	39	31	34	29	32	29
25	19	27	20	28	18	21	16	16	13	15	17
11	18	28	14	13	11	9	4	12	4		4
35	28	43	35	35	28	26	20	19	18		19
43	40	57	43	43	35	37	35	35	30		28
65	55	86	53	70	43	53	40	38	37		39
83	70	110	70	86	60	70	43	57	43		43
105	86	110	85	110	74	83	57	70	53		53
129	95	129	110	129	86	86	84	75	57		105
129	110	129	110	172	110	110	86	75	70	70	105
.7	0	1.3	0	2.5	0	0	1.8	0	0	0	3.7
.5	0	1.1	0	1.1	0	0	.3	0	0	0	2.1

TABLE 5.54.--READY-TO-EAT CEREALS, 50-75 GM PER CUP, ALL SUGAR LEVELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.8	.2	6.2	7.8	7.8
ON ONLY 1 OF 3 DAYS.....PCT.....	4.5	0	3.6	4.6	5.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.9	.2	1.6	2.1	2.0
ON ALL 3 OF 3 DAYS.....PCT.....	1.4	0	1.0	1.1	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	2.0	1.6	1.6	1.4
STANDARD DEVIATION.....DO.....	.8	.0	.8	.8	.7
MAXIMUM.....DO.....	6.0	2.0	4.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	25	17	18	21	22
STANDARD DEVIATION.....GM.....	20	0	13	14	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	6		4	8	4
25TH.....GM.....	13		8	9	17
50TH.....GM.....	17		17	17	17
75TH.....GM.....	33		25	25	30
90TH.....GM.....	50		38	47	40
95TH.....GM.....	60		42	50	60
99TH.....GM.....	100		50	67	85
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	0	30	39	46
STANDARD DEVIATION.....GM.....	28	0	14	17	21
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	47	25	38	43	43
STANDARD DEVIATION.....GM.....	23	0	13	21	22
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	47	0	37	39	61
STANDARD DEVIATION.....GM.....	28	0	10	16	19
MAXIMUM IN ANY 1 DAY.....GM.....	400	25	80	100	120
MAXIMUM OVER 3 DAYS.....GM.....	600	50	150	200	270
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	46	25	35	39	47
STANDARD DEVIATION.....GM.....	26	0	14	18	22
PERCENTILES: <u>1/</u>					
5TH.....GM.....	13		13	13	13
25TH.....GM.....	25		25	25	25
50TH.....GM.....	50		30	38	50
75TH.....GM.....	50		50	50	50
90TH.....GM.....	75		50	60	75
95TH.....GM.....	100		50	73	100
99TH.....GM.....	142		60	100	120
MAXIMUM.....GM.....	400	25	80	100	120
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	12.5	0	0	1.0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	.5	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.0	6.8	6.8	5.5	5.8	6.4	8.4	8.0	13.9	13.8	13.4	13.6
4.9	4.7	4.2	3.9	3.4	3.8	4.8	4.6	6.0	6.7	6.1	5.4
1.4	1.6	1.5	.9	1.5	1.6	2.2	1.8	3.3	4.5	2.4	3.3
.6	.6	1.2	.7	.9	.9	1.4	1.6	4.6	2.7	4.8	4.8
1.5	1.4	1.6	1.4	1.6	1.6	1.6	1.6	2.0	1.7	1.9	2.0
.8	.7	.9	.7	.8	.8	.8	.8	1.1	.8	1.0	1.0
5.0	4.0	5.0	3.0	4.0	4.0	4.0	5.0	6.0	4.0	5.0	4.0
31	26	36	24	32	23	27	23	28	21	24	26
26	18	30	15	25	15	19	16	27	17	17	23
8	8	9	4	10	7	6	5	5	4	5	5
17	17	17	17	17	13	15	13	9	10	13	13
25	17	25	17	25	17	20	17	19	17	17	17
42	33	40	33	37	33	33	30	33	25	33	33
60	50	75	50	58	50	50	46	60	40	50	45
67	60	111	50	75	50	67	55	75	50	63	60
133	100	133	67	150	67	100	83	120	100	80	83
66	57	61	54	66	46	50	43	37	37	40	37
48	36	26	29	34	23	27	23	17	16	17	18
67	52	57	54	52	48	51	43	46	37	46	38
34	21	21	21	24	18	22	21	25	22	29	17
74	59	88	41	67	40	51	41	46	37	34	41
37	25	33	17	37	18	22	20	36	24	17	31
400	275	200	168	200	150	200	200	240	180	120	300
510	350	540	225	450	225	300	338	530	360	240	600
65	54	67	50	61	44	50	42	42	37	38	38
40	26	31	25	33	21	25	22	26	22	21	24
25	25	25	15	17	14	13	13	9	7	8	6
50	50	50	30	40	25	30	25	25	25	19	25
50	50	50	50	50	50	50	43	38	30	38	30
75	56	90	50	75	50	60	50	50	50	50	50
100	100	100	75	100	75	83	60	75	60	60	60
125	100	120	100	142	75	100	75	100	60	63	75
200	150	180	120	150	120	113	110	120	120	120	150
400	200	200	168	150	150	200	200	170	180	120	150
3.6	.8	0	0	0	0	0	2.4	3.4	1.3	0	0
1.7	.4	0	0	0	0	0	.3	1.1	.3	0	0

TABLE 5.55.--READY-TO-EAT CEREALS, 80 GM OR MORE PER CUP, ALL SUGAR LEVELS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.5	.8	2.4	2.8	2.9
ON ONLY 1 OF 3 DAYS.....PCT.....	2.0	.2	1.6	2.0	2.2
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	.4	.7	.6	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.6	.2	.1	.2	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.6	2.1	1.4	1.4	1.4
STANDARD DEVIATION.....DO.....	.8	.7	.6	.9	.7
MAXIMUM.....DO.....	6.0	3.0	3.0	6.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	32	29	14	28	28
STANDARD DEVIATION.....GM.....	29	14	10	23	25
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	2				2
25TH.....GM.....	12				14
50TH.....GM.....	28				18
75TH.....GM.....	39				37
90TH.....GM.....	73				73
95TH.....GM.....	92				75
99TH.....GM.....	131				112
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	64	27	41	53	58
STANDARD DEVIATION.....GM.....	49	0	30	33	42
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	61	47	23	73	71
STANDARD DEVIATION.....GM.....	45	16	17	30	46
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	58	42	23	68	69
STANDARD DEVIATION.....GM.....	43	0	0	35	27
MAXIMUM IN ANY 1 DAY.....GM.....	495	65	110	170	224
MAXIMUM OVER 3 DAYS.....GM.....	678	129	110	339	337
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	60	42	31	59	61
STANDARD DEVIATION.....GM.....	46	12	26	35	42
PERCENTILES: ^{1/}					
5TH.....GM.....	5				7
25TH.....GM.....	28				28
50TH.....GM.....	55				55
75TH.....GM.....	84				84
90TH.....GM.....	112				112
95TH.....GM.....	126				113
99TH.....GM.....	220				224
MAXIMUM.....GM.....	440	65	110	170	224
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	20.4	0	2.3	1.3	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.0	0	6.6	1.9	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.2	2.3	2.2	2.3	3.5	3.6	3.7	3.9	5.6	4.8	3.9	6.2
1.2	1.6	1.7	1.7	2.1	2.2	2.0	2.3	2.4	2.5	1.8	3.1
.7	.4	.3	.3	.9	1.0	1.0	1.0	1.0	1.2	1.7	1.4
.2	.4	.2	.2	.5	.4	.7	.7	2.2	1.1	.4	1.7
1.6	1.5	1.4	1.4	1.6	1.5	1.7	1.6	2.0	1.7	1.6	1.9
.8	.8	.7	.7	.8	.7	.8	.8	.9	.8	.7	1.1
4.0	3.0	3.0	3.0	4.0	4.0	4.0	4.0	4.0	3.0	3.0	5.0
51	36	51	32	42	29	38	27	34	28	31	21
42	27	33	26	38	26	34	22	32	23	35	19
	5			3	2	4	2	2	2		2
	18			9	9	14	9	14	9		9
	28			37	19	28	19	28	21		19
	55			56	37	56	37	37	37		28
	73			85	73	84	57	84	56		37
	73			113	84	110	75	120	84		37
	151			194	112	147	85	149	121		120
101	76	128	72	75	55	66	55	67	69	34	59
68	45	89	39	60	44	42	42	28	41	21	30
100	67	91	67	83	59	73	53	41	34	60	21
80	36	26	48	53	40	42	29	24	26	54	16
94	74	98	77	84	58	70	44	49	44	79	30
36	33	30	37	50	33	51	30	44	34	40	29
440	226	495	220	280	220	226	220	224	180	168	120
660	452	495	385	581	336	678	360	448	363	336	360
95	73	111	72	79	56	69	50	51	49	56	33
71	41	57	43	57	41	47	36	41	39	48	31
	28			10	5	7	7	2	2		2
	55			28	28	28	28	21	21		7
	56			63	43	63	43	36	42		28
	110			110	84	110	78	84	57		49
	112			165	112	112	105	112	112		84
	113			220	112	149	113	120	120		110
	226			280	220	226	128	121	126		120
440	226	330	220	280	220	226	220	224	180	168	120
0	0	0	0	7.2	4.5	0	3.2	0	0	0	1.9
0	0	0	0	3.3	1.5	0	.7	0	0	0	1.6

TABLE 6.01.--COOKED DRIED BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.8	2.8	11.7	13.4	12.3
ON ONLY 1 OF 3 DAYS.....PCT.....	6.9	2.3	7.6	9.3	8.7
ON ONLY 2 OF 3 DAYS.....PCT.....	2.2	.5	3.2	2.6	2.9
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	.8	1.4	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.3	1.5	1.5	1.5
STANDARD DEVIATION.....DO.....	.8	.8	.7	.9	.8
MAXIMUM.....DO.....	6.0	4.0	4.0	6.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	78	24	40	55	58
STANDARD DEVIATION.....GM.....	65	29	28	53	45
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		8	8	12
25TH.....GM.....	32		16	24	32
50TH.....GM.....	63		32	32	48
75TH.....GM.....	118		62	63	63
90TH.....GM.....	154		85	123	123
95TH.....GM.....	190		95	157	127
99TH.....GM.....	317		123	247	247
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	167	39	86	110	132
STANDARD DEVIATION.....GM.....	118	30	61	78	86
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	174	113	93	125	125
STANDARD DEVIATION.....GM.....	105	55	45	79	80
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	175	0	64	126	126
STANDARD DEVIATION.....GM.....	102	0	22	93	60
MAXIMUM IN ANY 1 DAY.....GM.....	1,140	185	370	740	555
MAXIMUM OVER 3 DAYS.....GM.....	2,420	370	555	1,758	925
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	156	53	81	108	119
STANDARD DEVIATION.....GM.....	99	45	54	74	80
PERCENTILES: <u>1/</u>					
5TH.....GM.....	36		24	24	24
25TH.....GM.....	93		48	48	70
50TH.....GM.....	185		70	95	95
75TH.....GM.....	190		95	139	185
90TH.....GM.....	285		185	190	190
95TH.....GM.....	380		190	190	231
99TH.....GM.....	475		285	380	380
MAXIMUM.....GM.....	760	185	370	380	555
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	67.4	0	3.9	9.6	8.8
PROPORTION OF EATING OCCASIONS.....PCT.....	1.2	0	2.1	2.7	2.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
12.1	12.4	11.1	10.9	9.1	9.3	8.5	9.4	10.1	6.6	9.8	8.6
8.1	8.6	8.0	7.5	6.7	6.5	5.7	6.5	7.8	5.0	7.1	6.1
3.1	2.4	2.6	2.5	1.6	2.2	2.3	2.2	1.8	1.4	2.0	2.1
.9	1.3	.5	.9	.8	.6	.5	.7	.5	.2	.7	.5
1.5	1.5	1.4	1.5	1.5	1.5	1.6	1.5	1.5	1.4	1.6	1.5
.8	.9	.7	.8	.9	.8	.9	.9	.8	.8	.9	.8
5.0	5.0	4.0	6.0	6.0	6.0	6.0	6.0	4.0	4.0	4.0	4.0
80	75	101	79	96	78	98	76	79	65	85	67
60	60	75	53	74	66	82	62	71	51	75	44
16	8	19	16	15	15	24	15	12	12	8	15
32	32	62	32	42	32	59	32	40	31	32	31
63	63	63	63	63	62	63	62	62	62	63	63
100	95	127	123	127	123	127	123	93	95	123	94
135	127	190	158	190	143	185	135	127	127	190	127
190	187	247	185	253	190	247	190	190	127	190	143
317	317	377	278	317	380	380	315	507	317	370	190
170	159	217	167	211	161	210	163	188	145	156	159
106	97	135	99	150	110	128	118	114	101	101	101
160	156	239	178	218	184	199	173	217	164	202	136
84	83	110	58	111	106	105	110	188	91	76	67
196	168	246	165	205	184	250	157	98	179	266	142
76	89	118	67	85	96	166	66	51	14	89	42
760	555	760	570	855	760	925	1,140	760	570	555	380
950	1,130	1,425	855	1,213	1,500	2,420	1,520	1,520	950	1,110	570
156	150	213	161	194	159	190	149	161	136	160	130
89	89	119	82	126	88	109	90	106	85	93	64
46	24	69	48	38	35	46	28	24	46	23	36
93	95	185	95	95	95	95	93	93	95	93	93
185	139	190	185	190	185	185	143	185	95	185	95
190	190	278	190	190	190	190	190	190	190	190	190
278	190	380	190	380	278	370	238	370	190	285	190
370	380	380	370	380	370	380	370	380	380	370	190
380	380	571	380	571	380	570	380	380	380	416	278
570	555	740	570	760	570	740	760	555	380	416	380
.8	9.9	.9	2.6	3.8	11.2	4.6	11.3	0	0	0	0
.2	2.4	.4	1.1	.7	1.5	.7	1.1	0	0	0	0

TABLE 6.02.--BAKED BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.4	2.4	10.9	13.4	14.0
ON ONLY 1 OF 3 DAYS.....PCT.....	9.3	1.8	9.7	11.8	12.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	.2	1.1	1.5	1.5
ON ALL 3 OF 3 DAYS.....PCT.....	.1	.4	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.4	1.1	1.1	1.2
STANDARD DEVIATION.....DO.....	.4	.8	.4	.4	.4
MAXIMUM.....DO.....	4.0	3.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	67	32	37	42	53
STANDARD DEVIATION.....GM.....	53	37	35	29	39
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11		5	5	11
25TH.....GM.....	43		16	21	32
50TH.....GM.....	43		39	43	43
75TH.....GM.....	85		43	53	79
90TH.....GM.....	128		76	85	85
95TH.....GM.....	170		85	85	128
99TH.....GM.....	298		255	128	170
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	179	60	100	117	141
STANDARD DEVIATION.....GM.....	121	43	100	81	84
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	172	48	95	93	132
STANDARD DEVIATION.....GM.....	112	0	51	48	70
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	223	84	75	82	167
STANDARD DEVIATION.....GM.....	145	57	0	64	102
MAXIMUM IN ANY 1 DAY.....GM.....	1,140	190	765	510	510
MAXIMUM OVER 3 DAYS.....GM.....	1,710	475	765	510	1,140
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	177	68	97	110	138
STANDARD DEVIATION.....GM.....	119	52	92	78	84
PERCENTILES: <u>1/</u>					
5TH.....GM.....	32		16	16	31
25TH.....GM.....	128		43	64	80
50TH.....GM.....	128		84	127	128
75TH.....GM.....	255		128	128	158
90TH.....GM.....	255		128	255	255
95TH.....GM.....	383		255	255	255
99TH.....GM.....	510		765	383	383
MAXIMUM.....GM.....	1,020	190	765	510	510
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	28.0	0	0	1.6	4.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	.6	1.4

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.7	13.5	12.2	9.9	11.2	9.6	12.2	8.0	8.6	5.5	9.2	5.3
12.7	12.1	11.3	9.2	9.7	8.6	10.5	7.5	8.1	4.6	8.4	5.1
1.9	1.1	.7	.7	1.3	.9	1.5	.5	.4	.9	.8	.2
.2	.3	.2	0	.1	.1	.2	(2)	.1	0	0	0
1.2	1.1	1.1	1.1	1.2	1.1	1.2	1.1	1.1	1.2	1.1	1.0
.4	.4	.4	.3	.4	.4	.4	.3	.3	.4	.4	.2
3.0	4.0	4.0	3.0	4.0	3.0	4.0	3.0	3.0	2.0	3.0	2.0
75	64	75	59	86	60	89	54	74	67	88	53
60	57	59	36	62	42	67	36	44	34	49	35
21	11	11	11	21	11	21	11	16	11		5
43	43	43	43	43	43	43	32	43	43		43
64	43	64	43	85	43	85	43	85	85		43
85	85	85	85	85	85	85	85	85	85		75
128	127	128	106	170	85	170	85	128	92		85
170	170	170	128	170	128	234	127	170	127		85
340	340	298	170	340	221	340	170	213	190		170
194	171	202	163	230	166	227	152	213	183	233	144
127	124	125	98	141	114	138	96	116	85	107	76
180	136	191	172	222	141	221	148	168	148	291	255
98	75	96	55	155	84	130	72	111	64	49	0
359	238	337	0	174	116	340	223	212	0	0	0
160	167	175	0	60	50	93	32	0	0	0	0
1,020	1,020	765	510	1,140	1,020	1,020	765	510	383	510	255
1,710	1,520	1,710	638	1,520	1,020	1,290	765	638	570	765	510
194	168	203	164	222	158	226	151	207	171	238	153
129	123	129	94	137	98	137	92	120	82	100	79
64	32	32	32	64	32	64	32	32	32		16
128	84	128	128	128	120	128	84	128	128		128
128	128	190	128	255	128	255	128	191	143		128
255	255	255	255	255	255	255	255	255	255		255
255	255	383	255	383	255	383	255	383	255		255
383	383	510	319	510	281	510	255	510	255		255
638	570	571	510	637	510	571	510	510	380		255
1,020	1,020	765	510	1,020	663	1,020	765	510	383	510	255
.6	3.5	0	2.1	3.0	6.7	5.0	1.3	0	0	0	0
.2	1.0	0	1.3	.6	1.2	.7	.2	0	0	0	0

TABLE 6.03.--NUTS

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.0	0	1.9	3.2	5.8
ON ONLY 1 OF 3 DAYS.....PCT.....	3.9	0	1.3	2.4	4.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	0	.5	.8	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.1	0	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER....	1.3	0	1.3	1.3	1.3
STANDARD DEVIATION.....DO.....	.7	0	.5	.5	.6
MAXIMUM.....DO.....	8.0	0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	19	0	8	17	18
STANDARD DEVIATION.....GM.....	21	0	8	19	23
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	2			1	2
25TH.....GM.....	6			6	6
50TH.....GM.....	12			11	12
75TH.....GM.....	24			24	24
90TH.....GM.....	48			37	43
95TH.....GM.....	57			57	52
99TH.....GM.....	109			117	121
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	44	0	20	33	47
STANDARD DEVIATION.....GM.....	46	0	18	33	60
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	52	0	15	51	44
STANDARD DEVIATION.....GM.....	44	0	9	39	49
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	43	0	36	0	42
STANDARD DEVIATION.....GM.....	30	0	0	0	9
MAXIMUM IN ANY 1 DAY.....GM.....	432	0	57	288	363
MAXIMUM OVER 3 DAYS.....GM.....	495	0	108	351	363
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	43	0	19	39	44
STANDARD DEVIATION.....GM.....	43	0	16	36	48
PERCENTILES: <u>1</u> /					
5TH.....GM.....	5			4	6
25TH.....GM.....	16			14	17
50TH.....GM.....	32			28	28
75TH.....GM.....	57			57	47
90TH.....GM.....	85			85	85
95TH.....GM.....	144			113	144
99TH.....GM.....	189			144	252
MAXIMUM.....GM.....	426	0	57	144	288
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER....	114.2	0	.8	.9	8.9
PROPORTION OF EATING OCCASIONS.....PCT.....	4.5	0	2.8	1.3	6.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.9	4.1	3.2	3.2	6.7	5.1	6.8	5.8	4.6	4.6	2.0	3.0
3.3	3.6	2.8	2.8	5.3	4.2	5.2	4.4	3.7	2.9	1.5	2.2
.5	.4	.3	.4	1.1	.7	1.1	1.0	.7	1.2	0	.7
.1	(2)	0	.1	.3	.2	.4	.4	.2	.5	.5	0
1.2	1.2	1.1	1.2	1.3	1.3	1.3	1.4	1.2	1.6	2.0	1.2
.5	.5	.3	.5	.6	.6	.7	.9	.5	.9	1.8	.4
4.0	4.0	2.0	3.0	4.0	4.0	5.0	8.0	3.0	5.0	6.0	2.0
22	16	22	18	24	18	23	17	13	13	22	12
30	19	19	18	24	19	24	19	9	12	25	22
3	2			2	2	2	1	2	2		
6	6			9	6	7	6	6	4		
12	11			15	12	16	12	12	9		
24	23			30	24	24	24	18	19		
48	42			48	48	57	44	28	30		
96	48			57	57	72	48	30	33		
144	120			142	78	109	93	36	62		
45	38	59	41	58	49	44	39	31	31	24	17
51	35	58	35	60	54	33	40	24	29	16	14
97	71	59	58	62	38	77	39	28	19	0	49
71	55	11	39	50	21	48	26	14	14	0	55
62	31	0	72	39	38	54	41	30	28	64	0
35	0	0	0	24	24	28	36	2	18	6	0
432	216	340	221	426	324	288	280	108	145	85	144
495	360	340	246	432	324	435	424	108	185	218	288
54	43	59	44	55	44	51	36	30	24	34	30
57	41	53	36	54	43	41	36	21	23	19	39
6	6			6	5	7	5	5	4		
18	18			18	14	24	9	17	6		
36	32			36	32	36	28	27	14		
63	57			72	57	72	48	36	33		
144	85			144	85	113	72	57	57		
144	144			144	144	144	112	72	72		
288	216			227	216	168	168	108	136		
288	216	340	166	426	233	288	240	108	136	73	144
5.7	8.4	2.8	3.4	15.1	14.5	19.7	26.4	1.1	6.3	0	0
5.6	8.3	5.8	5.9	4.5	4.2	4.4	4.5	1.7	4.8	0	0

TABLE 6.04.--PEANUTS

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.8	0	1.1	2.2	4.3
ON ONLY 1 OF 3 DAYS.....PCT.....	2.3	0	.8	1.8	3.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.2	.4	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	0	1.3	1.2	1.2
STANDARD DEVIATION.....DO.....	.6	0	.6	.5	.5
MAXIMUM.....DO.....	4.0	0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	0	11	15	20
STANDARD DEVIATION.....GM.....	23	0	9	20	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3				5
25TH.....GM.....	9				6
50TH.....GM.....	14				12
75TH.....GM.....	24				24
90TH.....GM.....	48				48
95TH.....GM.....	63				84
99TH.....GM.....	120				121
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	52	0	26	34	54
STANDARD DEVIATION.....GM.....	46	0	17	34	64
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	62	0	21	53	47
STANDARD DEVIATION.....GM.....	51	0	10	54	58
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	55	0	36	0	34
STANDARD DEVIATION.....GM.....	35	0	0	0	3
MAXIMUM IN ANY 1 DAY.....GM.....	432	0	57	288	363
MAXIMUM OVER 3 DAYS.....GM.....	495	0	108	351	363
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	53	0	26	38	49
STANDARD DEVIATION.....GM.....	44	0	16	37	53
PERCENTILES: <u>1/</u>					
5TH.....GM.....	9				9
25TH.....GM.....	20				17
50TH.....GM.....	36				36
75TH.....GM.....	72				63
90TH.....GM.....	113				111
95TH.....GM.....	144				144
99TH.....GM.....	216				288
MAXIMUM.....GM.....	340	0	57	144	288
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	40.0	0	.8	0	3.4
PROPORTION OF EATING OCCASIONS.....PCT.....	3.1	0	5.1	0	3.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.7	2.7	1.7	1.6	3.6	2.6	3.8	2.9	3.1	1.7	.9	1.5
2.2	2.3	1.4	1.3	2.9	2.4	3.0	2.4	2.2	1.2	.9	1.0
.5	.3	.3	.2	.6	.1	.7	.3	.7	.4	0	.5
0	(2)	0	0	.1	(2)	.2	.2	.2	.1	0	0
1.2	1.2	1.2	1.2	1.3	1.1	1.3	1.3	1.4	1.4	1.0	1.3
.5	.5	.4	.5	.5	.4	.6	.7	.6	.7	.0	.5
3.0	4.0	2.0	3.0	3.0	4.0	4.0	4.0	3.0	4.0	1.0	2.0
27	18	29	18	27	21	26	20	15	17	12	17
32	21	22	16	23	20	24	21	9	15	4	30
3	2			6	3	5	3				
9	6			12	6	12	6				
12	12			23	12	21	12				
32	24			36	24	25	24				
64	48			48	48	57	48				
96	48			63	60	72	48				
165	120			144	108	144	141				
53	42	81	44	61	58	53	48	38	37	35	18
54	36	68	32	41	59	31	41	25	37	13	20
97	73	60	63	79	56	80	39	28	30	0	63
71	61	12	32	52	16	51	24	14	13	0	65
0	31	0	0	60	70	55	60	30	61	0	0
0	0	0	0	15	4	35	45	2	0	0	0
432	216	340	144	288	324	288	280	108	145	57	144
495	360	340	216	432	324	435	424	108	185	57	288
64	47	75	47	64	56	59	47	33	34	35	39
61	44	59	34	42	47	41	39	21	30	13	52
12	6			17	8	16	9				
28	18			36	19	28	18				
48	32			57	36	48	36				
72	48			72	72	72	72				
144	126			144	126	113	108				
144	144			144	144	144	144				
288	216			189	233	168	168				
288	216	340	144	189	233	288	170	108	136	57	144
4.7	6.0	1.1	1.0	3.5	1.8	5.6	11.0	1.1	0	0	0
6.8	8.9	4.1	3.5	2.0	1.1	2.3	4.2	2.3	0	0	0

TABLE 6.05.--PEANUT BUTTER
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	15.2	2.5	24.7	32.4	32.6
ON ONLY 1 OF 3 DAYS.....PCT.....	10.5	2.1	15.8	20.7	22.2
ON ONLY 2 OF 3 DAYS.....PCT.....	3.3	.3	7.1	8.3	7.1
ON ALL 3 OF 3 DAYS.....PCT.....	1.4	.2	1.8	3.4	3.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.2	1.5	1.5	1.5
STANDARD DEVIATION.....DO.....	.8	.6	.7	.8	.8
MAXIMUM.....DO.....	8.0	3.0	5.0	6.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	13	8	10	11	13
STANDARD DEVIATION.....GM.....	13	4	10	10	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		2	2	2
25TH.....GM.....	5		4	5	5
50TH.....GM.....	11		5	11	11
75TH.....GM.....	16		11	14	16
90TH.....GM.....	27		22	22	27
95TH.....GM.....	34		27	32	32
99TH.....GM.....	65		51	43	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	21	20	24	26
STANDARD DEVIATION.....GM.....	25	8	17	16	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	24	20	23	28
STANDARD DEVIATION.....GM.....	20	0	13	14	17
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	29	9	21	25	26
STANDARD DEVIATION.....GM.....	20	0	21	17	16
MAXIMUM IN ANY 1 DAY.....GM.....	454	32	194	194	161
MAXIMUM OVER 3 DAYS.....GM.....	454	48	274	323	242
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	27	19	20	23	25
STANDARD DEVIATION.....GM.....	22	9	18	16	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		5	5	5
25TH.....GM.....	16		8	16	16
50TH.....GM.....	16		16	16	16
75TH.....GM.....	32		32	32	32
90TH.....GM.....	48		32	32	48
95TH.....GM.....	65		48	48	65
99TH.....GM.....	129		98	65	97
MAXIMUM.....GM.....	454	32	194	194	161
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	229.7	0	6.1	13.0	32.9
PROPORTION OF EATING OCCASIONS.....PCT.....	2.7	0	1.6	1.5	3.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
27.0	24.1	19.3	12.5	12.8	12.2	10.6	9.2	12.9	9.3	10.1	10.7
18.7	16.9	12.4	9.1	8.6	9.2	6.8	7.3	8.2	6.3	5.4	7.3
5.9	5.4	5.4	2.4	2.6	2.4	2.3	1.4	2.7	2.5	2.6	2.0
2.4	1.8	1.6	1.0	1.6	.6	1.4	.5	1.9	.6	2.1	1.4
1.5	1.4	1.5	1.4	1.6	1.3	1.6	1.3	1.6	1.4	1.7	1.5
.8	.7	.8	.8	.9	.6	.9	.6	1.1	.6	.8	.8
6.0	4.0	5.0	4.0	6.0	4.0	8.0	5.0	6.0	3.0	3.0	4.0
15	13	19	12	20	11	16	10	16	9	15	11
12	10	18	12	21	9	14	8	12	7	13	11
3	3	4	2	4	2	4	2	4	2	5	2
5	5	11	5	9	5	5	5	5	5	7	5
11	11	16	11	11	9	11	7	11	5	11	8
22	16	22	16	22	13	22	11	22	11	16	13
32	22	40	22	43	22	32	22	32	18	32	27
43	32	48	32	59	32	43	22	43	22	48	32
65	59	97	65	108	43	65	38	54	43	57	57
32	30	39	27	40	26	32	25	34	21	26	22
25	24	29	20	43	20	24	20	29	18	11	14
32	27	41	32	44	25	33	19	31	19	25	28
18	17	26	26	28	15	27	11	15	12	15	20
31	24	45	25	42	26	28	22	26	12	32	21
18	12	32	12	31	17	15	15	15	6	16	16
194	258	290	145	454	129	194	145	145	145	97	98
258	258	387	258	454	199	290	274	161	145	170	170
30	27	38	26	38	25	30	23	29	19	28	22
22	20	26	21	35	19	23	18	22	16	14	18
5	5	11	5	11	5	5	5	5	5	11	5
16	16	16	16	16	16	16	16	16	11	16	8
32	24	32	16	32	16	32	16	32	16	28	16
32	32	48	32	48	32	32	32	32	31	32	32
65	48	65	48	65	48	57	32	52	32	48	48
65	65	97	65	93	65	65	65	65	48	57	57
129	97	129	129	145	97	129	97	129	65	57	98
194	258	129	145	454	129	194	145	145	145	65	98
22.6	23.1	10.1	10.7	27.7	14.3	28.7	28.6	5.1	1.5	1.7	3.7
2.7	3.1	2.5	4.1	3.5	1.7	3.5	3.4	2.2	.6	1.8	2.3

TABLE 7.01.--MEAT, POULTRY, FISH, AND THEIR MIXTURES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST CNCE IN 3 DAYS.....PCT.....	98.6	55.0	98.5	99.3	99.6
ON ONLY 1 OF 3 DAYS.....PCT.....	2.8	13.0	4.4	3.3	1.6
ON ONLY 2 OF 3 DAYS.....PCT.....	14.2	13.2	20.5	19.0	16.6
ON ALL 3 OF 3 DAYS.....PCT.....	81.5	28.8	73.7	77.0	81.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	4.8	3.3	4.3	4.5	4.7
STANDARD DEVIATION.....DO.....	1.7	1.8	1.7	1.6	1.5
MAXIMUM.....DO.....	12.0	10.0	9.0	9.0	10.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	193	87	96	114	146
STANDARD DEVIATION.....GM.....	114	77	59	64	75
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	51	8	23	33	48
25TH.....GM.....	113	37	54	70	93
50TH.....GM.....	172	64	82	102	133
75TH.....GM.....	249	128	125	145	185
90TH.....GM.....	338	191	172	199	244
95TH.....GM.....	405	255	216	233	283
99TH.....GM.....	562	336	283	309	379
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	146	84	73	125	129
STANDARD DEVIATION.....GM.....	133	62	57	109	135
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	173	104	102	111	136
STANDARD DEVIATION.....GM.....	113	64	69	64	72
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	211	122	108	127	159
STANDARD DEVIATION.....GM.....	112	85	58	63	74
MAXIMUM IN ANY 1 DAY.....GM.....	2,350	638	980	745	992
MAXIMUM OVER 3 DAYS.....GM.....	4,010	1,696	1,116	1,435	1,720
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	121	80	66	77	93
STANDARD DEVIATION.....GM.....	107	53	57	64	76
PERCENTILES: ^{1/}					
5TH.....GM.....	16	14	8	14	16
25TH.....GM.....	56	40	28	30	44
50TH.....GM.....	97	63	56	57	84
75TH.....GM.....	162	115	85	97	112
90TH.....GM.....	235	133	125	142	180
95TH.....GM.....	324	184	170	205	240
99TH.....GM.....	510	224	268	336	383
MAXIMUM.....GM.....	1,792	425	528	720	960
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	2704.5	5.9	52.3	103.2	127.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	.7	1.2	1.4	1.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
99.7	99.6	99.8	99.5	99.3	98.3	99.5	99.3	98.6	98.3	98.8	98.4
1.9	2.5	1.4	3.5	1.8	3.9	1.0	2.9	2.6	4.3	3.7	5.1
13.9	16.9	11.4	17.5	10.6	17.3	7.4	14.8	10.7	16.9	11.1	18.0
83.8	80.2	87.0	78.5	86.9	77.0	91.0	81.6	85.3	77.0	84.0	75.3
4.9	4.6	5.3	4.6	5.3	4.5	5.5	4.6	5.0	4.2	4.7	4.1
1.6	1.6	1.7	1.6	1.7	1.6	1.7	1.6	1.8	1.6	1.8	1.6
10.0	11.0	12.0	10.0	12.0	10.0	12.0	10.0	12.0	9.0	10.0	9.0
191	162	253	175	277	179	261	177	207	153	182	136
92	86	126	89	143	96	122	93	108	87	100	79
65	49	84	56	94	52	94	53	64	43	51	35
124	104	170	112	180	110	174	113	131	93	112	85
178	148	235	162	254	164	244	163	189	141	166	124
243	204	311	222	343	228	323	225	266	196	230	173
310	273	417	289	453	304	423	301	343	258	332	229
363	319	498	343	537	353	489	354	407	312	374	297
460	417	647	462	755	473	639	456	557	440	490	410
171	111	215	141	192	173	225	129	144	129	233	127
138	89	191	124	119	140	156	96	119	121	391	96
184	154	225	171	269	170	237	174	182	145	169	138
110	86	125	93	167	93	143	105	102	84	130	89
205	178	270	194	294	199	271	193	223	172	195	152
88	83	123	85	139	93	119	89	104	85	94	76
1,536	1,376	1,708	1,225	2,350	1,329	1,957	1,501	1,360	1,869	1,502	1,128
2,018	2,791	2,957	1,879	4,010	2,260	3,401	2,695	2,627	2,360	1,665	1,540
117	105	143	115	158	120	143	117	125	108	116	99
97	87	123	92	135	99	123	99	106	92	107	83
17	20	24	19	24	16	18	16	16	16	16	14
56	54	57	56	64	56	57	56	56	54	44	43
90	85	112	97	113	97	112	92	108	88	90	85
148	119	180	142	220	160	184	150	168	135	150	113
227	210	273	224	314	227	278	227	235	224	227	218
291	250	368	270	442	304	368	288	340	252	294	245
490	480	612	480	705	490	600	490	490	480	490	460
1,020	960	1,596	1,225	1,792	1,275	1,596	1,440	1,020	1,588	1,470	750
178.7	121.3	101.2	119.3	295.9	325.6	495.3	514.0	69.2	115.8	30.6	48.7
1.7	1.2	1.4	1.8	1.4	1.4	1.8	1.6	1.3	1.6	1.2	1.2

TABLE 7.02.--MEAT, POULTRY, AND FISH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	96.9	35.5	95.8	97.7	98.7
ON ONLY 1 OF 3 DAYS.....PCT.....	8.8	12.7	12.1	9.5	9.8
ON ONLY 2 OF 3 DAYS.....PCT.....	27.2	7.5	29.8	31.6	32.2
ON ALL 3 OF 3 DAYS.....PCT.....	61.0	15.3	53.9	56.6	56.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	3.9	2.7	3.6	3.7	3.7
STANDARD DEVIATION.....DO.....	1.7	1.7	1.7	1.6	1.6
MAXIMUM.....DO.....	12.0	10.0	9.0	9.0	10.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	134	51	65	80	96
STANDARD DEVIATION.....GM.....	87	55	41	47	55
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	28	4	13	19	23
25TH.....GM.....	73	16	33	47	56
50TH.....GM.....	117	38	57	73	85
75TH.....GM.....	176	61	87	103	127
90TH.....GM.....	245	118	122	138	168
95TH.....GM.....	296	151	142	165	200
99TH.....GM.....	413	283	183	232	269
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	126	59	64	84	99
STANDARD DEVIATION.....GM.....	97	49	46	60	65
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	142	59	74	94	109
STANDARD DEVIATION.....GM.....	83	41	39	54	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	165	83	83	98	120
STANDARD DEVIATION.....GM.....	87	67	41	45	54
MAXIMUM IN ANY 1 DAY.....GM.....	2,350	383	651	559	992
MAXIMUM OVER 3 DAYS.....GM.....	3,480	1,021	889	1,030	1,178
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	102	56	54	64	77
STANDARD DEVIATION.....GM.....	82	39	39	46	54
PERCENTILES: <u>1/</u>					
5TH.....GM.....	16	9	8	13	16
25TH.....GM.....	44	28	28	28	40
50TH.....GM.....	85	47	44	56	61
75TH.....GM.....	129	78	71	85	110
90TH.....GM.....	209	128	112	112	140
95TH.....GM.....	234	128	113	144	170
99TH.....GM.....	393	135	176	224	276
MAXIMUM.....GM.....	1,792	255	336	476	907
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2152.3	4.0	43.2	90.4	97.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	.8	1.2	1.4	1.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
98.5	98.1	97.8	97.2	98.4	96.3	98.9	97.9	98.1	97.2	97.4	96.1
8.7	9.3	7.3	12.3	7.7	10.8	4.4	8.9	8.3	9.3	5.4	10.7
31.1	33.6	27.4	29.8	25.6	29.4	19.9	27.5	21.0	28.1	22.4	26.2
58.7	55.2	63.1	55.2	65.0	56.0	74.6	61.5	68.9	59.8	69.6	59.2
3.9	3.7	4.2	3.6	4.2	3.6	4.6	3.8	4.3	3.7	4.2	3.6
1.7	1.6	1.8	1.6	1.8	1.6	1.8	1.7	1.8	1.6	1.8	1.6
10.0	9.0	12.0	9.0	11.0	9.0	12.0	10.0	12.0	9.0	10.0	9.0
125	106	164	117	186	122	187	127	150	112	138	102
71	62	95	71	108	73	99	75	85	65	83	61
29	26	38	24	46	28	54	28	37	28	28	24
74	62	96	65	113	68	117	75	88	65	76	56
113	98	149	107	168	110	173	115	140	103	123	95
163	139	211	154	240	160	238	166	194	150	192	135
219	181	283	208	320	217	313	223	260	201	241	176
259	207	328	245	375	257	366	264	295	233	295	212
340	288	475	371	509	345	490	372	426	321	364	335
126	107	159	128	181	129	178	121	130	107	131	111
94	75	103	112	143	89	109	87	78	61	165	75
137	120	175	126	208	137	186	141	159	120	124	110
69	60	95	66	107	70	89	79	89	58	55	56
154	134	197	151	219	153	211	155	176	140	163	127
69	62	93	67	108	73	97	72	80	63	82	60
952	1,376	1,411	952	2,350	1,072	1,438	1,501	937	688	897	1,128
1,455	2,605	2,957	1,427	3,480	2,028	2,208	2,005	1,854	1,274	1,563	1,311
96	86	117	96	132	100	122	99	105	92	98	85
70	64	90	72	106	75	95	78	80	67	78	63
16	16	16	16	24	16	16	16	16	14	14	13
46	44	56	48	56	48	56	45	50	44	36	38
85	84	97	85	112	85	112	85	86	84	85	82
113	112	168	113	170	124	168	123	140	113	126	112
180	168	224	170	255	194	227	184	224	170	210	168
224	196	280	224	336	227	288	225	227	224	227	218
336	302	448	344	480	368	448	368	392	336	368	304
896	960	1,344	896	1,792	960	1,131	1,440	768	624	678	564
148.8	94.8	79.4	93.9	225.3	265.9	393.7	406.4	47.1	85.9	29.2	46.6
1.9	1.2	1.4	1.8	1.4	1.4	1.8	1.5	1.0	1.4	1.3	1.3

TABLE 7.03.--BEEF, PORK, LAMB, AND VEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,900	524	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	84.6	23.2	78.2	82.8	84.6
ON ONLY 1 OF 3 DAYS.....PCT.....	29.8	12.7	33.3	34.5	34.7
ON ONLY 2 OF 3 DAYS.....PCT.....	33.8	4.7	31.2	33.0	35.6
ON ALL 3 OF 3 DAYS.....PCT.....	21.0	5.8	13.7	15.2	14.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.4	2.0	2.1	2.1	2.1
STANDARD DEVIATION.....DO.....	1.3	1.4	1.2	1.1	1.1
MAXIMUM.....DO.....	10.0	8.0	8.0	7.0	9.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	85	39	37	46	57
STANDARD DEVIATION.....GM.....	67	49	28	34	41
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11	3	5	5	8
25TH.....GM.....	37	9	17	21	28
50TH.....GM.....	69	24	29	37	48
75TH.....GM.....	112	46	52	62	75
90TH.....GM.....	168	85	77	88	111
95TH.....GM.....	211	135	89	108	131
99TH.....GM.....	317	340	121	161	206
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	128	56	61	75	96
STANDARD DEVIATION.....GM.....	98	47	45	50	74
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	53	64	82	99
STANDARD DEVIATION.....GM.....	79	28	37	47	52
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	139	88	63	75	95
STANDARD DEVIATION.....GM.....	83	74	33	38	48
MAXIMUM IN ANY 1 DAY.....GM.....	1,792	383	418	559	624
MAXIMUM OVER 3 DAYS.....GM.....	3,328	1,021	579	856	972
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	107	58	53	66	82
STANDARD DEVIATION.....GM.....	85	42	40	46	55
PERCENTILES: <u>1/</u>					
5TH.....GM.....	16	9	8	8	14
25TH.....GM.....	46	28	23	28	43
50TH.....GM.....	86	47	46	56	84
75TH.....GM.....	140	85	80	85	112
90TH.....GM.....	224	128	112	112	142
95TH.....GM.....	252	128	115	140	171
99TH.....GM.....	432	135	170	224	280
MAXIMUM.....GM.....	1,792	255	280	448	504
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	1097.1	1.6	29.4	52.6	42.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.4	.6	1.7	1.8	1.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
87.1	84.2	87.9	82.6	87.5	82.1	90.7	86.1	85.7	84.5	87.5	80.7
33.4	33.9	27.0	32.2	27.8	32.5	24.6	30.1	24.2	30.6	21.6	28.8
35.5	34.5	37.1	32.5	34.8	32.3	35.2	34.7	33.5	33.6	32.4	33.0
18.2	15.7	23.8	17.9	24.9	17.2	31.0	21.3	28.0	20.4	33.4	18.8
2.2	2.1	2.6	2.2	2.6	2.2	2.8	2.4	2.7	2.3	2.9	2.3
1.2	1.1	1.4	1.2	1.4	1.2	1.5	1.3	1.5	1.2	1.6	1.3
9.0	8.0	9.0	8.0	8.0	9.0	10.0	9.0	9.0	9.0	8.0	7.0
76	67	105	76	119	78	117	80	92	68	93	65
55	52	73	56	87	58	82	58	63	49	69	50
13	11	17	11	16	13	20	11	16	9	11	5
37	32	56	37	57	37	58	37	46	35	42	35
65	56	92	61	102	65	99	67	76	56	75	56
103	90	140	105	160	106	157	108	130	92	129	85
149	127	197	143	224	149	219	155	169	134	187	136
181	151	241	184	272	181	270	193	208	168	224	156
266	233	340	276	405	285	409	278	302	234	303	224
120	109	152	113	177	129	172	128	142	108	138	96
76	81	99	88	134	97	106	97	101	66	110	63
129	115	158	127	180	129	167	126	130	112	122	111
69	59	81	67	95	66	95	70	70	62	70	63
123	113	164	128	185	131	171	126	136	106	135	102
65	70	84	62	103	70	90	68	71	58	79	62
896	1,376	1,025	896	1,792	960	1,344	972	720	688	840	1,128
1,308	2,605	2,375	1,176	3,328	1,512	2,118	1,527	1,420	965	1,204	1,249
103	94	123	102	140	107	126	102	103	89	97	86
71	69	90	73	109	80	99	79	79	66	81	66
16	16	16	16	16	16	16	14	16	10	10	8
56	56	56	56	56	56	56	43	40	35	24	32
86	84	112	88	112	92	112	86	86	84	84	84
138	112	168	129	194	140	168	138	140	112	138	112
184	168	224	170	276	224	226	188	224	168	224	168
224	224	280	224	336	232	291	224	226	224	226	224
336	336	448	344	468	368	448	384	368	280	368	322
896	960	896	896	1,792	960	1,080	972	552	540	602	564
62.0	50.1	48.5	47.6	119.7	144.1	194.2	191.3	30.1	40.1	18.8	24.5
1.5	1.3	1.5	1.8	1.4	1.5	1.6	1.3	1.2	1.2	1.4	1.3

TABLE 7.04.--BEEF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	67.3	15.6	60.1	65.5	67.2
ON ONLY 1 OF 3 DAYS.....PCT.....	38.5	9.4	38.6	41.6	40.8
ON ONLY 2 OF 3 DAYS.....PCT.....	22.4	4.0	17.0	19.7	22.0
ON ALL 3 OF 3 DAYS.....PCT.....	6.4	2.2	4.4	4.2	4.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER....	1.7	1.7	1.5	1.5	1.6
STANDARD DEVIATION.....DO.....	.9	1.1	.8	.8	.8
MAXIMUM.....DO.....	8.0	6.0	5.0	6.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	75	32	33	40	51
STANDARD DEVIATION.....GM.....	59	32	24	30	35
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	19	5	7	9	14
25TH.....GM.....	37	9	18	19	28
50TH.....GM.....	57	19	28	34	38
75TH.....GM.....	95	43	43	55	65
90TH.....GM.....	149	85	67	75	96
95TH.....GM.....	187	100	81	93	114
99TH.....GM.....	283	151	108	143	173
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	142	53	67	82	104
STANDARD DEVIATION.....GM.....	95	40	40	46	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	147	65	71	86	105
STANDARD DEVIATION.....GM.....	80	37	36	46	46
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	159	72	72	90	107
STANDARD DEVIATION.....GM.....	83	49	29	39	48
MAXIMUM IN ANY 1 DAY.....GM.....	1,792	228	418	448	561
MAXIMUM OVER 3 DAYS.....GM.....	3,312	454	579	816	928
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	133	56	64	79	97
STANDARD DEVIATION.....GM.....	85	41	38	43	52
PERCENTILES: 1/					
5TH.....GM.....	41	9	14	28	28
25TH.....GM.....	84	27	35	56	56
50TH.....GM.....	112	46	56	80	85
75TH.....GM.....	168	84	84	112	112
90TH.....GM.....	224	128	112	112	162
95TH.....GM.....	280	128	126	149	184
99TH.....GM.....	448	135	170	224	288
MAXIMUM.....GM.....	1,792	198	280	448	504
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER....	683.2	1.6	13.4	31.2	29.2
PROPORTION OF EATING OCCASIONS.....PCT.....	1.6	1.2	1.4	1.8	1.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
69.0	68.2	70.3	65.9	72.2	65.1	75.2	68.1	64.2	63.4	65.2	57.3
39.2	40.3	36.4	38.7	37.5	40.1	38.1	39.1	34.6	40.7	39.5	34.3
24.5	23.2	25.2	21.0	26.6	19.9	26.9	22.8	21.4	17.6	18.3	18.4
5.3	4.7	8.7	6.1	8.0	5.0	10.2	6.2	8.2	5.1	7.4	4.6
1.7	1.6	1.8	1.7	1.8	1.6	1.8	1.7	1.8	1.6	1.7	1.6
.9	.8	1.0	.9	.9	.8	1.0	.8	1.0	.8	1.0	.8
8.0	5.0	7.0	5.0	7.0	6.0	7.0	8.0	6.0	6.0	6.0	5.0
69	60	93	68	105	69	99	71	79	59	81	61
50	47	65	49	78	49	71	51	53	42	58	44
19	19	27	19	27	19	24	19	19	19	28	14
37	29	47	37	56	37	47	37	37	29	37	37
56	47	75	56	84	56	75	56	66	47	61	47
92	75	121	90	138	93	131	93	112	75	93	75
131	112	171	127	203	131	191	137	149	112	150	121
164	138	211	167	243	160	233	173	177	149	187	159
246	227	311	240	357	241	336	252	271	215	303	224
129	121	175	133	196	141	175	139	149	121	173	117
72	77	100	93	128	89	103	91	85	69	115	57
138	121	171	131	192	144	181	142	145	122	148	125
62	81	77	54	94	67	91	72	66	59	77	63
161	128	179	147	212	141	190	141	154	133	158	132
70	50	88	61	115	60	86	57	57	49	76	61
896	1,376	896	896	1,792	960	1,344	972	560	672	840	448
1,120	2,421	2,246	1,176	3,312	1,460	1,891	1,221	948	896	1,120	840
124	111	152	123	174	131	160	128	133	113	140	112
66	70	87	73	110	79	96	79	70	62	77	60
43	42	56	43	56	43	56	43	56	40	43	35
84	84	92	84	112	84	92	84	85	84	85	81
112	112	113	112	161	112	138	112	112	112	112	112
162	113	184	140	224	168	224	168	168	138	170	138
224	170	230	192	288	224	276	224	224	184	224	210
224	224	312	224	368	276	336	270	227	224	270	224
336	368	448	392	553	448	504	448	368	345	420	336
896	960	896	896	1,792	960	896	972	552	540	602	336
37.2	30.2	26.2	25.7	85.7	80.3	127.8	132.7	16.3	17.3	13.5	14.9
1.5	1.3	1.5	1.6	1.7	1.5	1.9	1.7	1.3	1.0	2.2	1.6

TABLE 7.05.--BEEF STEAKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	27.6	2.3	19.5	24.2	25.6
ON ONLY 1 OF 3 DAYS.....PCT.....	23.4	2.1	16.8	21.6	23.6
ON ONLY 2 OF 3 DAYS.....PCT.....	3.9	.1	2.3	2.4	1.8
ON ALL 3 OF 3 DAYS.....PCT.....	.4	.1	.4	.2	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.2	1.2	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.5	.5	.4	.3
MAXIMUM.....DO.....	6.0	3.0	3.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	65	17	24	29	39
STANDARD DEVIATION.....GM.....	51	10	16	24	29
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		5	8	12
25TH.....GM.....	31		12	15	23
50TH.....GM.....	56		19	27	31
75TH.....GM.....	75		31	37	46
90TH.....GM.....	123		46	51	68
95TH.....GM.....	151		56	61	95
99TH.....GM.....	245		75	112	153
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	163	44	62	75	106
STANDARD DEVIATION.....GM.....	107	22	41	43	63
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	172	69	61	86	126
STANDARD DEVIATION.....GM.....	88	0	28	72	87
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	200	28	54	91	87
STANDARD DEVIATION.....GM.....	149	0	11	38	12
MAXIMUM IN ANY 1 DAY.....GM.....	1,792	85	255	448	504
MAXIMUM OVER 3 DAYS.....GM.....	3,312	138	269	816	816
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	162	45	60	77	107
STANDARD DEVIATION.....GM.....	104	21	37	50	67
PERCENTILES: <u>1/</u>					
5TH.....GM.....	46		14	23	30
25TH.....GM.....	92		28	46	69
50TH.....GM.....	138		56	70	92
75TH.....GM.....	216		84	92	115
90TH.....GM.....	276		112	112	184
95TH.....GM.....	368		122	168	224
99TH.....GM.....	552		168	252	368
MAXIMUM.....GM.....	1,792	85	255	448	504
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	498.5	1.6	9.8	22.8	19.9
PROPORTION OF EATING OCCASIONS.....PCT.....	4.0	11.6	4.1	4.8	3.9

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
24.5	25.9	27.3	23.7	31.8	27.1	35.0	30.3	25.0	23.3	24.8	18.1
21.8	23.0	24.0	21.0	26.2	23.4	27.4	25.2	20.6	19.8	21.9	14.9
2.6	2.6	3.1	2.6	5.0	3.3	6.6	4.8	3.9	3.3	2.0	3.0
0	.2	.2	.1	.6	.4	1.0	.3	.5	.1	.8	.1
1.1	1.1	1.2	1.1	1.2	1.2	1.3	1.2	1.3	1.2	1.3	1.2
.3	.4	.4	.4	.5	.4	.6	.5	.6	.5	.6	.5
3.0	3.0	3.0	3.0	6.0	4.0	6.0	5.0	5.0	4.0	4.0	3.0
51	47	61	56	90	63	84	64	67	53	71	56
34	31	51	40	74	41	56	45	40	34	43	41
15	14	19	18	24	19	28	19	23	18	23	14
31	28	37	31	54	37	47	37	37	31	37	28
38	37	60	46	75	56	75	56	56	46	61	46
61	56	84	65	112	75	112	75	77	67	82	75
85	75	117	107	160	120	149	123	122	87	123	107
112	112	149	131	216	141	187	149	155	123	165	136
168	157	262	224	328	205	276	240	204	184	224	224
135	126	181	149	223	165	202	163	166	140	193	135
78	73	96	99	139	95	110	105	83	87	116	79
148	126	172	159	217	157	206	164	169	134	168	165
71	55	90	71	103	62	94	76	70	55	56	83
0	149	341	178	309	184	209	176	162	65	131	103
0	81	212	24	267	101	82	77	39	0	56	0
896	504	896	693	1,792	736	1,344	972	552	672	672	448
896	800	1,798	740	3,312	1,460	1,540	1,040	644	672	672	672
137	125	178	147	217	161	196	159	157	132	169	137
78	71	105	91	140	91	105	97	82	72	85	76
46	43	56	54	68	56	72	56	56	54	56	45
85	84	112	92	113	105	112	92	92	85	96	84
112	112	168	115	184	140	170	138	138	112	161	112
168	149	224	168	276	184	224	184	210	168	224	184
224	224	300	224	368	276	336	271	230	224	276	252
280	252	368	324	448	360	414	336	306	224	368	306
368	448	576	553	673	540	553	540	448	448	368	336
896	504	896	693	1,792	736	720	972	552	460	448	336
23.1	23.4	17.5	18.0	66.8	61.7	93.3	91.3	10.6	14.0	11.5	13.2
4.0	3.7	4.0	4.5	4.3	3.6	4.2	3.5	3.0	2.9	6.9	6.0

TABLE 7.06.--BEEF ROASTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST CNCE IN 3 DAYS.....PCT.....	19.1	2.1	12.6	14.8	16.2
ON ONLY 1 OF 3 DAYS.....PCT.....	15.6	2.1	11.4	13.1	14.7
ON ONLY 2 OF 3 DAYS.....PCT.....	3.2	0	1.2	1.6	1.5
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	0	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	1.0	1.1	1.2	1.1
STANDARD DEVIATION.....DO.....	.6	.0	.3	.4	.4
MAXIMUM.....DO.....	7.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	52	9	20	26	33
STANDARD DEVIATION.....GM.....	39	7	12	20	22
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		5	9	9
25TH.....GM.....	28		9	14	19
50TH.....GM.....	37		19	19	28
75TH.....GM.....	68		27	30	37
90TH.....GM.....	101		37	56	57
95TH.....GM.....	131		47	65	75
99TH.....GM.....	205		56	112	112
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	130	26	54	70	90
STANDARD DEVIATION.....GM.....	90	21	31	48	55
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	129	0	54	67	100
STANDARD DEVIATION.....GM.....	71	0	19	43	44
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	138	0	0	113	21
STANDARD DEVIATION.....GM.....	60	0	0	19	0
MAXIMUM IN ANY 1 DAY.....GM.....	896	70	168	420	336
MAXIMUM OVER 3 DAYS.....GM.....	1,312	70	168	560	396
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	124	26	54	68	89
STANDARD DEVIATION.....GM.....	82	21	29	46	53
PERCENTILES: <u>1/</u>					
5TH.....GM.....	35		14	28	28
25TH.....GM.....	72		28	43	56
50TH.....GM.....	112		56	56	84
75TH.....GM.....	168		70	84	112
90TH.....GM.....	224		84	112	168
95TH.....GM.....	280		112	144	168
99TH.....GM.....	448		140	280	288
MAXIMUM.....GM.....	896	70	144	336	336
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	55.0	0	1.7	1.4	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	1.2	.5	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.6	18.0	22.5	18.6	21.2	16.5	25.7	20.0	21.8	18.2	20.2	13.8
15.1	16.0	19.1	15.3	17.5	13.8	19.9	16.0	16.7	14.2	13.2	9.4
2.4	2.0	3.1	3.0	3.4	2.4	5.1	3.5	4.4	3.7	6.1	3.5
.1	.1	.3	.3	.4	.2	.7	.5	.6	.3	.9	.8
1.2	1.2	1.2	1.2	1.2	1.2	1.3	1.3	1.3	1.3	1.5	1.5
.5	.4	.5	.5	.5	.5	.6	.6	.7	.6	.8	.9
5.0	3.0	4.0	5.0	4.0	4.0	7.0	5.0	5.0	4.0	4.0	5.0
50	45	61	48	63	49	66	49	54	47	65	53
30	38	37	34	45	34	49	33	38	37	49	42
14	14	19	14	19	14	19	14	14	14	14	14
29	28	37	28	37	28	37	28	28	24	28	27
43	37	56	37	47	37	56	37	47	37	48	37
56	56	75	56	75	59	75	56	75	56	75	68
93	75	108	93	115	93	131	93	108	93	149	112
112	108	132	103	149	112	149	112	113	115	149	149
149	240	216	172	224	174	233	159	205	187	201	205
134	118	167	121	154	129	161	123	130	107	148	106
73	92	95	73	93	84	112	78	75	71	111	60
122	131	134	111	169	119	153	112	119	122	135	114
50	86	62	40	88	58	84	54	55	57	79	50
137	87	131	197	155	112	147	114	175	162	135	156
79	65	51	60	50	50	64	45	43	67	50	56
560	896	720	720	864	718	896	896	448	540	602	336
645	1,120	720	792	972	718	1,312	896	688	730	602	616
127	116	151	117	153	120	149	114	121	108	127	104
66	86	88	64	95	75	100	67	66	68	81	59
43	36	56	43	43	28	43	42	28	43	28	28
84	70	86	84	86	70	84	72	84	56	56	56
112	112	129	112	114	112	112	112	112	86	112	86
168	129	196	140	196	162	168	140	168	129	168	135
224	216	243	172	243	224	252	172	224	168	224	210
224	224	288	224	336	280	336	224	224	224	224	224
336	448	448	336	448	336	540	336	336	450	420	224
448	896	720	448	864	576	896	896	448	540	602	336
4.0	.8	4.0	4.5	3.5	4.4	16.0	8.9	3.5	1.4	0	.8
.9	.2	1.1	1.3	.3	.4	1.0	.5	1.1	.3	0	.4

TABLE 7.07.--GROUND BEEF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	36.3	3.3	35.8	39.2	41.5
ON ONLY 1 OF 3 DAYS.....PCT.....	28.8	2.9	29.3	31.2	33.3
ON ONLY 2 OF 3 DAYS.....PCT.....	6.6	.5	5.5	7.4	7.6
ON ALL 3 OF 3 DAYS.....PCT.....	1.0	0	1.0	.6	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.1	1.3	1.3	1.3
STANDARD DEVIATION.....DO.....	.6	.4	.5	.6	.5
MAXIMUM.....DO.....	6.0	2.0	4.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	53	18	29	36	41
STANDARD DEVIATION.....GM.....	37	11	18	23	23
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	19		7	9	19
25TH.....GM.....	28		19	19	28
50TH.....GM.....	37		25	28	37
75TH.....GM.....	75		37	42	51
90TH.....GM.....	93		56	63	75
95TH.....GM.....	117		67	75	80
99TH.....GM.....	187		80	112	131
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	127	48	74	88	101
STANDARD DEVIATION.....GM.....	71	27	42	44	47
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	130	41	70	85	100
STANDARD DEVIATION.....GM.....	63	21	31	29	37
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	138	0	63	132	98
STANDARD DEVIATION.....GM.....	58	0	16	53	29
MAXIMUM IN ANY 1 DAY.....GM.....	1,045	112	280	448	399
MAXIMUM OVER 3 DAYS.....GM.....	1,525	136	308	560	560
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	123	46	70	84	97
STANDARD DEVIATION.....GM.....	66	26	36	36	41
PERCENTILES: <u>1/</u>					
5TH.....GM.....	56		19	28	42
25TH.....GM.....	84		44	56	71
50TH.....GM.....	112		56	84	85
75TH.....GM.....	149		85	112	112
90TH.....GM.....	224		112	113	142
95TH.....GM.....	224		113	148	170
99TH.....GM.....	339		170	224	224
MAXIMUM.....GM.....	960	112	280	224	336
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	82.7	0	1.8	5.4	7.3
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.4	.6	.8

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
45.2	41.4	43.5	38.3	40.1	34.4	37.1	32.9	31.3	29.5	32.8	30.4
35.1	33.5	32.6	27.6	29.8	27.6	29.6	27.2	25.2	25.2	27.7	23.0
9.0	7.5	8.5	9.0	8.8	6.2	6.4	5.1	5.5	3.3	4.5	6.2
1.1	.5	2.4	1.6	1.5	.7	1.1	.7	.7	1.0	.6	1.2
1.3	1.3	1.4	1.4	1.4	1.3	1.3	1.2	1.3	1.2	1.2	1.3
.6	.5	.7	.7	.7	.5	.6	.5	.6	.6	.5	.6
5.0	4.0	5.0	5.0	4.0	4.0	5.0	4.0	5.0	6.0	4.0	4.0
53	44	67	53	74	50	63	48	55	45	54	49
36	32	44	38	50	30	41	29	31	26	27	29
19	19	19	19	24	19	24	19	24	19	28	19
28	28	37	28	37	28	37	28	37	28	37	37
37	37	56	37	57	37	56	37	47	37	47	37
65	56	84	66	85	65	75	56	75	56	75	56
95	75	131	99	149	84	112	75	93	75	93	75
112	93	149	112	187	103	149	112	112	93	112	107
187	149	242	207	243	156	224	149	179	149	149	163
121	109	155	116	172	122	152	121	136	114	141	114
59	59	86	69	100	59	79	63	54	45	58	39
133	115	156	124	172	121	155	115	129	110	136	118
52	74	64	60	72	44	80	44	45	45	44	56
151	108	145	127	178	127	145	132	163	121	114	105
74	28	54	39	70	57	48	39	40	45	2	42
480	1,045	725	896	896	454	672	800	336	448	336	448
1,014	1,525	924	1,008	1,120	922	1,344	800	616	672	448	616
120	106	142	114	162	118	146	116	130	109	132	111
59	62	74	64	82	57	74	57	51	45	49	46
56	56	56	42	56	56	56	56	56	50	84	56
84	84	85	84	112	84	112	84	98	84	112	84
112	112	112	112	142	112	112	112	112	112	112	112
142	112	170	112	224	113	170	113	168	112	150	112
224	168	224	168	226	170	224	168	224	168	224	168
224	213	255	224	336	224	255	224	224	224	224	224
336	336	448	320	448	336	340	336	227	224	336	255
480	960	640	896	672	454	672	800	336	448	336	336
6.3	4.7	3.1	3.1	13.9	9.0	9.0	14.1	2.2	2.0	0	.9
.5	.4	.4	.4	.6	.4	.4	.5	.5	.3	0	.2

TABLE 7.08.--PORK

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	49.9	10.1	44.2	46.0	46.7
ON ONLY 1 OF 3 DAYS.....PCT.....	30.5	9.2	29.2	31.3	33.5
ON ONLY 2 OF 3 DAYS.....PCT.....	13.5	.6	11.6	11.6	10.4
ON ALL 3 OF 3 DAYS.....PCT.....	5.9	.3	3.4	3.2	2.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.7	1.3	1.5	1.5	1.5
STANDARD DEVIATION.....DO.....	1.0	1.1	.8	.8	.8
MAXIMUM.....DO.....	8.0	8.0	6.0	6.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	39	28	19	24	28
STANDARD DEVIATION.....GM.....	39	53	17	21	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u> :					
5TH.....GM.....	5	3	3	3	3
25TH.....GM.....	11	9	5	8	9
50TH.....GM.....	28	19	13	19	23
75TH.....GM.....	54	33	28	32	37
90TH.....GM.....	84	43	42	49	56
95TH.....GM.....	111	45	54	63	75
99TH.....GM.....	178	340	77	97	119
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	84	55	46	56	68
STANDARD DEVIATION.....GM.....	77	40	43	46	60
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	76	64	35	48	56
STANDARD DEVIATION.....GM.....	62	41	25	34	35
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	70	340	36	46	50
STANDARD DEVIATION.....GM.....	63	0	26	37	39
MAXIMUM IN ANY 1 DAY.....GM.....	1,128	383	243	389	512
MAXIMUM OVER 3 DAYS.....GM.....	1,564	1,021	312	543	512
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	69	66	37	47	57
STANDARD DEVIATION.....GM.....	69	44	36	44	49
PERCENTILES: <u>1/</u> :					
5TH.....GM.....	8	8	5	8	8
25TH.....GM.....	16	28	8	16	16
50TH.....GM.....	44	57	24	30	43
75TH.....GM.....	92	116	56	70	85
90TH.....GM.....	160	128	86	97	112
95TH.....GM.....	194	128	112	135	142
99TH.....GM.....	320	135	160	188	224
MAXIMUM.....GM.....	1,080	135	243	389	432
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	379.0	0	16.0	17.6	12.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.2	0	2.2	1.5	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
48.8	47.0	56.1	46.2	52.6	47.0	57.6	51.2	53.8	47.9	59.2	45.4
33.0	34.4	34.4	30.8	31.4	31.2	29.4	31.5	25.1	25.7	25.0	26.2
12.0	9.7	15.3	11.2	14.8	11.3	18.6	13.6	17.7	15.3	20.1	12.1
3.8	3.0	6.4	4.2	6.4	4.5	9.5	6.2	10.9	6.9	14.1	7.0
1.6	1.5	1.7	1.6	1.7	1.6	1.9	1.7	2.0	1.8	2.2	1.9
.9	.8	1.0	.8	1.0	.9	1.1	1.0	1.1	1.0	1.3	1.1
7.0	6.0	7.0	5.0	6.0	6.0	8.0	7.0	7.0	7.0	7.0	6.0
35	31	45	35	51	37	51	36	45	34	44	33
33	28	39	31	47	39	48	35	50	32	43	35
4	3	5	5	5	5	5	5	5	5	5	4
11	11	15	12	16	11	16	11	16	11	15	9
28	27	36	28	37	28	38	28	32	27	33	27
47	41	61	47	71	49	72	48	57	47	56	47
75	65	94	75	111	76	104	79	95	75	92	73
96	81	120	103	143	100	149	99	130	96	141	79
165	124	168	145	224	166	221	158	230	134	235	124
82	75	97	76	109	83	105	78	95	74	85	67
73	69	78	66	98	70	90	71	92	67	81	58
76	67	82	77	101	72	90	72	75	61	75	69
59	44	56	55	77	52	77	54	65	50	55	44
58	61	86	66	83	77	83	59	67	55	64	52
44	41	59	36	65	79	66	54	77	44	60	64
648	945	988	630	816	648	1,112	864	720	464	468	1,128
689	945	1,020	673	1,238	1,512	1,564	1,347	1,420	965	784	1,249
68	64	79	68	88	70	79	64	67	56	59	53
65	57	75	60	84	67	81	62	72	56	67	57
8	8	14	8	16	8	12	8	8	8	8	8
16	24	24	21	24	20	24	16	16	16	16	16
53	53	56	52	56	56	56	43	40	32	28	35
92	87	112	92	113	92	112	92	86	85	84	84
158	135	170	144	194	160	184	142	162	117	144	112
194	170	224	184	264	194	227	184	194	172	194	140
291	272	336	280	408	312	368	276	368	230	288	258
648	630	648	426	680	576	1,080	504	460	432	454	564
20.8	19.9	21.3	21.0	34.0	63.8	61.8	52.0	13.8	14.4	5.3	4.7
1.3	1.3	1.6	2.0	.9	1.6	1.1	.8	1.1	1.0	.7	.6

TABLE 7.09.--PORK CHOPS AND STEAKS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	14.3	1.6	13.7	16.8	15.9
ON ONLY 1 OF 3 DAYS.....PCT.....	13.4	1.6	13.0	16.1	15.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	0	.6	.6	.4
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	(2)	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.1	1.0	1.1	1.1	1.0
STANDARD DEVIATION.....DO.....	.3	.0	.3	.2	.2
MAXIMUM.....DO.....	4.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	46	16	24	29	33
STANDARD DEVIATION.....GM.....	28	12	14	17	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		8	9	15
25TH.....GM.....	27		15	18	23
50TH.....GM.....	38		23	27	31
75TH.....GM.....	61		32	32	37
90TH.....GM.....	78		41	50	53
95TH.....GM.....	97		47	61	65
99TH.....GM.....	149		74	97	97
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	129	48	69	81	94
STANDARD DEVIATION.....GM.....	72	36	38	39	44
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	132	0	68	110	110
STANDARD DEVIATION.....GM.....	60	0	25	34	45
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	131	0	0	112	77
STANDARD DEVIATION.....GM.....	65	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	552	116	243	360	291
MAXIMUM OVER 3 DAYS.....GM.....	895	116	243	403	368
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	128	48	67	83	95
STANDARD DEVIATION.....GM.....	70	36	37	41	45
PERCENTILES: <u>1/</u>					
5TH.....GM.....	45		23	28	44
25TH.....GM.....	80		43	53	69
50TH.....GM.....	104		58	80	92
75TH.....GM.....	160		92	97	106
90TH.....GM.....	203		116	138	156
95TH.....GM.....	276		138	176	184
99TH.....GM.....	368		188	194	276
MAXIMUM.....GM.....	552	116	243	360	291
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	83.2	0	5.5	3.5	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	1.4	0	3.6	1.2	.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.5	15.3	16.4	14.7	14.6	14.5	15.0	13.9	14.4	10.1	17.4	10.2
13.8	14.8	15.7	14.1	13.6	13.8	13.5	12.8	12.5	8.7	16.2	9.6
.7	.4	.7	.7	1.0	.6	1.4	.9	1.6	1.2	1.2	.6
0	.1	.1	0	0	(2)	.1	.1	.3	.1	0	0
1.1	1.0	1.1	1.0	1.1	1.1	1.1	1.1	1.2	1.2	1.1	1.1
.2	.2	.3	.2	.3	.2	.4	.4	.5	.5	.3	.3
2.0	3.0	3.0	2.0	2.0	3.0	3.0	4.0	3.0	4.0	3.0	2.0
41	36	50	42	63	43	62	44	54	46	47	37
23	18	25	23	35	26	32	24	33	26	31	22
15	15	20	15	23	15	27	15	19	20	19	9
27	27	31	27	36	27	39	27	31	31	28	23
32	32	46	32	61	34	53	38	46	32	32	32
53	46	61	53	77	53	77	53	61	59	56	47
65	61	84	65	97	65	97	69	92	85	92	63
86	65	96	80	129	92	124	93	123	107	129	65
129	107	129	138	184	142	184	123	186	123	156	120
118	106	144	117	176	124	173	122	145	118	126	108
61	49	66	56	87	65	80	58	84	57	78	65
121	105	129	143	189	108	138	128	120	124	163	85
49	51	44	47	66	47	68	54	43	43	39	20
0	38	135	0	0	259	189	76	135	108	0	0
0	0	0	0	0	27	5	22	49	1	0	0
473	368	368	426	552	552	464	426	485	368	468	360
473	381	414	426	650	895	582	596	558	434	468	360
116	105	140	119	175	124	163	120	137	115	125	104
56	50	64	57	87	66	79	57	74	49	55	62
46	44	59	46	69	46	69	46	53	56	57	20
80	80	92	80	97	80	92	80	85	88	85	69
97	97	138	97	180	97	160	97	122	97	106	92
145	117	184	160	224	160	194	154	184	138	160	116
194	184	234	194	284	194	276	184	194	184	194	184
232	194	276	225	320	240	291	224	291	194	234	194
291	284	348	368	468	368	448	300	368	284	276	360
322	368	368	426	552	552	464	368	460	368	276	360
3.2	3.7	5.5	3.2	13.0	12.8	15.5	5.7	4.2	3.4	2.1	0
1.0	1.1	2.3	1.4	2.1	1.6	1.8	.5	2.2	1.6	2.0	0

TABLE 7.10.--HAM

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	16.3	5.4	9.5	11.7	12.7
ON ONLY 1 OF 3 DAYS.....PCT.....	12.6	4.5	7.7	9.8	10.8
ON ONLY 2 OF 3 DAYS.....PCT.....	3.2	.6	1.6	2.0	1.8
ON ALL 3 OF 3 DAYS.....PCT.....	.5	.3	.2	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.5	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.7	1.5	.5	.5	.4
MAXIMUM.....DO.....	8.0	8.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	39	40	20	23	27
STANDARD DEVIATION.....GM.....	38	69	14	15	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		5	7	9
25TH.....GM.....	19		9	12	14
50TH.....GM.....	29		19	19	23
75TH.....GM.....	48		28	28	37
90TH.....GM.....	75		37	38	47
95TH.....GM.....	103		56	56	56
99TH.....GM.....	192		75	75	75
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	90	70	55	61	72
STANDARD DEVIATION.....GM.....	69	35	37	36	40
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	99	64	39	53	65
STANDARD DEVIATION.....GM.....	76	41	25	31	26
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	104	340	46	0	24
STANDARD DEVIATION.....GM.....	79	0	20	0	3
MAXIMUM IN ANY 1 DAY.....GM.....	1,080	383	224	194	224
MAXIMUM OVER 3 DAYS.....GM.....	1,512	1,021	224	280	252
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	87	80	50	55	66
STANDARD DEVIATION.....GM.....	67	40	35	33	37
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28		9	14	22
25TH.....GM.....	43		28	28	42
50TH.....GM.....	70		43	56	56
75TH.....GM.....	112		56	70	86
90TH.....GM.....	168		86	112	112
95TH.....GM.....	224		112	112	135
99TH.....GM.....	336		224	170	180
MAXIMUM.....GM.....	1,080	135	224	189	224
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	64.7	0	2.2	3.0	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	0	1.8	1.2	.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
14.6	13.4	19.1	14.9	18.8	15.2	22.1	16.4	17.0	16.2	14.8	17.3
11.9	10.9	14.7	11.9	13.2	12.4	15.7	13.3	12.4	12.2	10.6	12.0
2.5	2.1	3.8	2.3	4.8	2.6	5.5	2.5	3.3	3.4	4.2	4.4
.2	.4	.6	.7	.8	.2	.9	.5	1.3	.7	0	.8
1.3	1.3	1.3	1.3	1.5	1.3	1.4	1.3	1.5	1.4	1.4	1.5
.5	.6	.6	.6	.7	.6	.8	.7	.8	.7	.7	.8
4.0	4.0	5.0	4.0	5.0	5.0	8.0	5.0	4.0	5.0	4.0	5.0
37	32	46	32	50	37	48	35	47	39	41	34
33	29	38	26	44	42	45	35	44	30	26	21
9	9	12	9	9	9	9	9	9	9	9	9
19	14	23	14	19	19	19	15	19	19	18	19
28	24	37	28	37	28	37	28	36	28	38	29
47	37	56	38	57	46	57	38	56	56	56	47
75	65	85	63	102	70	93	69	93	75	75	70
96	76	112	84	137	86	142	85	115	96	84	75
187	149	196	132	227	164	224	159	230	131	125	93
91	82	114	80	110	86	102	81	104	90	96	76
70	85	92	69	87	65	73	54	80	64	62	39
101	74	99	74	116	86	118	102	115	89	96	80
75	34	70	37	78	50	89	96	101	44	33	28
69	63	100	69	108	233	112	90	83	98	0	69
24	20	40	27	67	183	65	40	46	54	0	21
560	945	972	630	680	648	1,080	864	560	336	342	224
689	945	972	630	1,238	1,512	1,123	1,339	1,064	648	374	280
88	75	103	73	103	86	100	81	96	84	88	67
66	60	79	51	78	66	78	59	70	55	54	33
28	28	28	28	28	28	28	28	28	28	28	28
56	43	56	43	56	43	56	43	56	43	45	43
72	56	84	56	85	67	84	57	84	72	84	56
112	86	113	86	113	112	112	112	112	112	112	86
168	113	172	129	180	168	196	168	170	168	162	112
224	168	224	168	245	224	224	170	224	216	196	114
340	336	448	315	448	432	425	315	360	270	288	168
560	630	648	410	680	576	1,080	448	432	288	288	170
2.8	3.8	4.4	2.6	.9	9.7	11.8	10.9	4.4	3.8	0	2.9
.7	1.0	1.3	.9	.1	.9	.8	.7	1.6	1.0	0	1.1

TABLE 7.11.--PORK ROASTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.5	.3	2.1	1.6	2.6
ON ONLY 1 OF 3 DAYS.....PCT.....	2.2	.3	1.9	1.5	2.5
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	.2	.1	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER.....	1.2	1.0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.0	.4	.3	.2
MAXIMUM.....DO.....	5.0	1.0	3.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	47	23	18	21	30
STANDARD DEVIATION.....GM.....	37	14	10	12	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	9				
25TH.....GM.....	27				
50TH.....GM.....	37				
75TH.....GM.....	56				
90TH.....GM.....	84				
95TH.....GM.....	112				
99TH.....GM.....	205				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	121	68	47	63	84
STANDARD DEVIATION.....GM.....	78	42	24	36	69
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	122	0	56	23	88
STANDARD DEVIATION.....GM.....	76	0	(2)	0	21
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	178	0	0	0	0
STANDARD DEVIATION.....GM.....	79	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,080	112	112	144	432
MAXIMUM OVER 3 DAYS.....GM.....	1,116	112	113	144	432
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	117	68	47	58	84
STANDARD DEVIATION.....GM.....	78	42	22	33	65
PERCENTILES: 1/					
5TH.....GM.....	28				
25TH.....GM.....	56				
50TH.....GM.....	112				
75TH.....GM.....	162				
90TH.....GM.....	224				
95TH.....GM.....	227				
99TH.....GM.....	432				
MAXIMUM.....GM.....	648	112	112	140	432
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER.....	16.2	0	0	0	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.4	0	0	0	2.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.8	2.1	3.0	2.4	1.9	1.8	3.3	3.1	2.7	2.6	4.2	2.3
2.7	1.9	2.9	2.0	1.6	1.7	2.7	2.6	1.8	2.0	3.1	1.7
(2)	.2	.2	.4	.3	.1	.6	.4	.9	.6	.7	.5
0	0	0	0	(2)	0	.1	(2)	0	0	.4	.2
1.1	1.1	1.1	1.2	1.3	1.1	1.2	1.2	1.4	1.3	1.7	1.6
.3	.4	.4	.4	.6	.3	.5	.5	.5	.5	1.2	.9
3.0	3.0	3.0	3.0	4.0	2.0	3.0	4.0	2.0	2.0	5.0	4.0
43	38	48	50	65	35	59	46	65	38	82	42
32	31	25	33	45	19	46	29	63	27	55	27
9				19	9	19	14				
28				37	19	29	28				
37				56	37	46	37				
47				75	38	75	56				
75				103	64	112	75				
93				149	75	149	108				
216				299	93	205	168				
130	97	139	126	167	101	136	124	111	100	217	106
94	61	74	71	78	51	75	74	55	82	126	80
9	131	107	140	124	92	160	108	180	79	86	101
0	80	16	61	54	38	94	44	122	30	58	22
0	0	0	0	299	0	205	149	0	0	184	43
0	0	0	0	0	0	0	0	0	0	44	0
648	504	392	420	560	227	1,080	504	360	432	432	258
648	585	392	448	896	280	1,116	504	720	432	672	258
120	102	129	127	155	98	141	116	139	88	140	81
89	63	66	75	82	52	80	70	111	68	112	54
28				36	28	43	35				
72				90	56	86	70				
105				168	86	113	112				
140				224	112	170	140				
196				280	184	224	203				
224				280	224	258	224				
648				448	224	448	454				
648	420	280	420	448	227	540	504	360	432	432	258
1.8	0	2.8	0	1.0	0	5.4	3.7	0	0	0	0
2.9	0	5.9	0	1.0	0	2.7	1.4	0	0	0	0

TABLE 7.12.--BACON
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	25.3	1.7	23.5	22.0	20.7
ON ONLY 1 OF 3 DAYS.....PCT.....	16.7	1.7	15.9	16.2	15.5
ON ONLY 2 OF 3 DAYS.....PCT.....	5.8	0	5.9	4.1	4.0
ON ALL 3 OF 3 DAYS.....PCT.....	2.9	0	1.7	1.6	1.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	1.5	1.0	1.4	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.0	.6	.6	.6
MAXIMUM.....DO.....	6.0	1.0	3.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	12	4	7	7	9
STANDARD DEVIATION.....GM.....	11	3	7	7	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	3		2	3	3
25TH.....GM.....	5		3	3	5
50TH.....GM.....	8		5	5	5
75TH.....GM.....	13		8	9	11
90TH.....GM.....	21		11	13	16
95TH.....GM.....	32		16	19	21
99TH.....GM.....	57		38	53	43
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	25	12	14	17	20
STANDARD DEVIATION.....GM.....	22	8	10	11	16
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	23	0	16	17	20
STANDARD DEVIATION.....GM.....	18	0	15	16	12
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	23	0	12	16	17
STANDARD DEVIATION.....GM.....	17	0	5	11	9
MAXIMUM IN ANY 1 DAY.....GM.....	454	28	112	112	142
MAXIMUM OVER 3 DAYS.....GM.....	624	28	168	196	142
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	24	12	14	16	20
STANDARD DEVIATION.....GM.....	20	8	13	13	15
PERCENTILES: <u>1</u> /					
5TH.....GM.....	8		4	5	8
25TH.....GM.....	16		8	8	12
50TH.....GM.....	16		8	16	16
75TH.....GM.....	24		16	16	24
90TH.....GM.....	40		24	24	32
95TH.....GM.....	57		28	36	48
99TH.....GM.....	113		85	84	85
MAXIMUM.....GM.....	454	28	112	112	142
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	125.2	0	6.9	4.1	3.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.9	0	2.0	.8	.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
23.4	20.7	28.6	21.2	26.9	21.6	31.7	26.7	32.2	26.5	40.1	24.0
17.4	16.5	18.4	15.8	17.8	14.9	18.7	17.7	16.9	14.9	18.8	14.4
4.3	3.4	7.7	4.3	6.1	4.6	8.2	5.8	8.6	7.8	11.4	5.5
1.7	.8	2.6	1.1	3.0	2.1	4.8	3.2	6.8	3.8	9.9	4.0
1.4	1.3	1.5	1.3	1.5	1.4	1.6	1.5	1.7	1.7	1.9	1.6
.6	.6	.7	.6	.7	.7	.8	.7	.9	.9	.9	.8
4.0	5.0	4.0	4.0	4.0	5.0	5.0	5.0	4.0	6.0	6.0	5.0
10	10	13	10	16	11	14	10	14	11	14	10
8	9	11	8	14	10	13	10	13	11	17	8
3	3	4	3	4	3	4	3	4	3	3	3
5	5	5	5	8	5	7	5	5	5	5	5
8	8	11	8	11	8	11	8	11	8	11	8
13	11	16	13	19	13	16	11	16	13	16	12
19	19	27	19	32	21	27	19	24	21	28	19
27	27	32	27	41	32	38	27	37	33	38	27
46	56	57	53	76	57	62	57	76	57	76	38
24	22	28	22	35	24	30	22	24	22	27	24
20	16	23	14	33	19	23	19	16	17	32	26
21	24	27	26	31	24	24	21	26	19	29	18
12	20	16	16	24	18	14	15	26	13	37	9
21	30	25	22	30	24	28	20	22	24	19	14
13	18	15	12	18	16	22	14	12	18	11	7
170	227	227	144	454	144	227	251	170	256	454	170
255	340	227	178	454	240	624	284	284	304	470	170
23	23	27	23	32	23	26	21	24	20	23	18
18	17	21	17	27	18	21	17	18	17	32	18
8	8	8	8	10	8	8	8	8	8	8	8
16	16	16	16	16	16	16	15	16	10	16	8
16	16	24	16	24	16	24	16	16	16	16	16
24	24	32	24	32	24	32	24	24	24	24	20
40	40	48	40	57	40	48	32	40	32	38	28
57	57	57	57	84	57	57	57	57	57	57	36
112	113	113	85	113	113	113	85	113	85	113	113
170	113	227	144	454	144	227	227	146	240	454	170
6.3	7.2	3.9	8.1	8.8	29.6	15.4	21.7	.9	5.5	1.8	1.8
1.0	1.3	.7	2.0	.6	1.8	.6	.8	.1	.7	.4	.5

TABLE 7.13.--LAMB
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.5	2.6	1.4	.6	.5
ON ONLY 1 OF 3 DAYS.....PCT.....	1.4	1.7	1.4	.6	.5
ON ONLY 2 OF 3 DAYS.....PCT.....	.1	.9	0	0	0
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER....	1.1	1.4	1.0	1.0	1.0
STANDARD DEVIATION.....DO.....	.3	.5	.0	.0	.0
MAXIMUM.....DO.....	3.0	2.0	1.0	1.0	1.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	54	24	24	30	46
STANDARD DEVIATION.....GM.....	33	13	15	20	29
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	15				
25TH.....GM.....	36				
50TH.....GM.....	46				
75TH.....GM.....	73				
90TH.....GM.....	93				
95TH.....GM.....	112				
99TH.....GM.....	149				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	149	56	72	90	139
STANDARD DEVIATION.....GM.....	88	34	46	59	86
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	141	49	0	0	0
STANDARD DEVIATION.....GM.....	63	15	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	64	0	0	0	0
STANDARD DEVIATION.....GM.....	0	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	864	128	128	218	336
MAXIMUM OVER 3 DAYS.....GM.....	864	136	128	218	336
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	146	52	72	90	139
STANDARD DEVIATION.....GM.....	84	29	46	59	86
PERCENTILES: 1/					
5TH.....GM.....	43				
25TH.....GM.....	88				
50TH.....GM.....	123				
75TH.....GM.....	184				
90TH.....GM.....	227				
95TH.....GM.....	280				
99TH.....GM.....	448				
MAXIMUM.....GM.....	864	128	128	218	336
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER....	9.6	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
.9	.7	.5	1.0	1.2	1.4	1.8	1.8	3.5	3.0	2.6	2.8
.8	.7	.5	.9	.9	1.3	1.8	1.8	3.0	2.7	2.6	1.9
.1	0	.1	.1	.3	.1	.1	.1	.5	.4	0	.9
0	0	0	0	0	(2)	0	0	0	0	0	0
1.1	1.0	1.1	1.1	1.3	1.1	1.1	1.0	1.2	1.1	1.0	1.3
.3	.0	.3	.3	.5	.3	.3	.2	.4	.4	.0	.5
2.0	1.0	2.0	2.0	3.0	3.0	3.0	2.0	2.0	2.0	1.0	2.0
61	42	58	41	63	50	64	48	70	61	42	55
42	23	30	17	36	25	43	27	38	28	14	26
					19	28	15		31		
					31	38	28		36		
					37	56	46		48		
					73	73	65		79		
					79	97	76		112		
					109	149	93		113		
					112	288	149		118		
154	127	157	116	167	145	186	135	193	160	126	135
73	68	83	47	91	74	126	65	106	65	43	56
280	0	154	96	130	112	190	199	157	172	0	113
0	0	0	0	65	0	47	29	55	4	0	40
0	0	0	0	0	64	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
336	280	280	218	504	336	864	280	448	336	218	218
560	280	308	218	504	336	864	448	448	354	218	356
171	127	156	112	150	139	182	140	175	158	126	124
80	68	81	43	87	73	121	66	89	60	43	53
					40	84	45		92		
					92	112	85		109		
					112	168	138		144		
					168	218	196		210		
					237	280	224		236		
					253	290	276		276		
					336	864	280		336		
336	280	280	218	504	336	864	280	448	336	218	218
.8	0	0	0	0	0	2.1	1.9	0	0	0	4.8
3.7	0	0	0	0	0	2.2	1.4	0	0	0	13.0

TABLE 7.14.--VEAL
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.3	3.2	1.2	1.6	2.0
ON ONLY 1 OF 3 DAYS.....PCT.....	2.1	1.9	1.2	1.6	2.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.2	.5	0	0	(2)
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	.8	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	2.0	1.0	1.0	1.0
STANDARD DEVIATION.....DO.....	.4	1.2	.0	.0	.2
MAXIMUM.....DO.....	5.0	5.0	1.0	1.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	49	36	27	25	40
STANDARD DEVIATION.....GM.....	27	30	9	11	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15				
25TH.....GM.....	28				
50TH.....GM.....	38				
75TH.....GM.....	59				
90TH.....GM.....	90				
95TH.....GM.....	95				
99TH.....GM.....	144				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	138	67	80	75	118
STANDARD DEVIATION.....GM.....	74	44	28	33	78
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	121	48	0	0	110
STANDARD DEVIATION.....GM.....	45	19	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	72	72	0	0	0
STANDARD DEVIATION.....GM.....	35	35	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	448	128	128	138	336
MAXIMUM OVER 3 DAYS.....GM.....	504	340	128	138	336
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	130	54	80	75	115
STANDARD DEVIATION.....GM.....	71	37	28	33	72
PERCENTILES: <u>1/</u>					
5TH.....GM.....	42				
25TH.....GM.....	84				
50TH.....GM.....	112				
75TH.....GM.....	168				
90TH.....GM.....	224				
95TH.....GM.....	276				
99TH.....GM.....	352				
MAXIMUM.....GM.....	448	128	128	138	336
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	26.2	0	0	3.9	.8
PROPORTION OF EATING OCCASIONS.....PCT.....	2.7	0	0	13.9	2.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.5	1.5	1.5	2.1	2.1	1.9	2.6	2.7	2.4	4.2	3.8	3.2
1.4	1.4	1.3	2.0	2.0	1.7	2.4	2.5	1.9	3.2	3.1	2.9
.1	.1	.2	.1	.1	.1	.2	.2	.5	1.0	.7	.3
0	0	0	0	0	0	0	0	0	0	0	0
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.3	1.4	1.1
.3	.3	.3	.3	.5	.3	.4	.3	.4	.6	.9	.3
2.0	2.0	2.0	2.0	4.0	2.0	3.0	3.0	2.0	3.0	4.0	2.0
45	35	64	48	57	50	60	48	61	49	42	44
28	21	34	25	27	21	32	24	29	24	24	29
				19	28	24	19		9		
				37	37	37	32		36		
				56	43	56	39		44		
				75	72	75	59		59		
				93	77	93	84		89		
				112	93	117	90		90		
				135	93	168	114		112		
124	99	175	136	165	142	168	140	171	127	101	113
81	61	98	67	78	56	83	70	93	56	27	47
122	112	152	142	164	130	169	105	114	103	114	159
9	0	2	35	21	12	53	27	22	42	51	68
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
448	280	405	336	405	280	448	405	360	270	172	224
448	280	405	337	405	280	504	405	360	336	344	448
124	96	170	131	156	140	164	131	150	111	93	121
75	46	87	62	76	53	84	63	86	57	25	54
				56	84	71	43		28		
				88	112	112	84		65		
				138	116	140	112		112		
				224	168	220	168		140		
				276	224	280	224		211		
				280	232	336	264		220		
				405	280	432	336		270		
448	224	405	336	405	280	448	342	360	270	168	224
3.2	0	.9	.9	0	0	3.5	4.7	0	8.3	0	0
9.4	0	3.9	2.8	0	0	2.5	2.2	0	8.7	0	0

TABLE 7.15.--POULTRY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,900	524	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	42.8	18.2	42.2	42.6	45.1
ON ONLY 1 OF 3 DAYS.....PCT.....	33.0	11.6	32.1	34.1	36.1
ON ONLY 2 OF 3 DAYS.....PCT.....	8.8	5.3	8.4	8.1	7.9
ON ALL 3 OF 3 DAYS.....PCT.....	1.1	1.2	1.7	.4	1.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.6	1.4	1.3	1.3
STANDARD DEVIATION.....DO.....	.6	.8	.7	.5	.6
MAXIMUM.....DO.....	7.0	4.0	4.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	57	31	33	38	43
STANDARD DEVIATION.....GM.....	39	25	23	24	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	17	3	9	12	14
25TH.....GM.....	32	14	18	20	27
50TH.....GM.....	46	28	28	33	37
75TH.....GM.....	73	43	38	47	56
90TH.....GM.....	109	58	66	69	75
95TH.....GM.....	132	84	85	82	100
99TH.....GM.....	191	134	112	114	141
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	140	64	77	96	110
STANDARD DEVIATION.....GM.....	84	45	43	52	59
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	65	82	90	102
STANDARD DEVIATION.....GM.....	68	34	42	44	48
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	122	75	65	93	97
STANDARD DEVIATION.....GM.....	55	42	30	27	45
MAXIMUM IN ANY 1 DAY.....GM.....	1,282	227	336	485	679
MAXIMUM OVER 3 DAYS.....GM.....	1,282	411	418	582	679
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	128	60	73	90	103
STANDARD DEVIATION.....GM.....	77	38	44	50	56
PERCENTILES: <u>1/</u>					
5TH.....GM.....	42	9	20	28	32
25TH.....GM.....	82	33	40	56	60
50TH.....GM.....	112	56	60	85	97
75TH.....GM.....	168	84	100	112	113
90TH.....GM.....	224	128	113	150	186
95TH.....GM.....	280	128	140	194	220
99TH.....GM.....	388	142	222	234	291
MAXIMUM.....GM.....	990	213	336	448	567
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	115.1	1.7	2.5	4.0	7.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	1.1	.4	.4	.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
44.3	44.0	43.8	43.7	39.9	43.2	43.1	44.6	42.4	45.1	36.1	40.3
35.8	35.9	34.4	36.0	30.4	33.3	33.7	33.8	29.1	32.7	24.2	27.6
7.9	7.6	8.4	6.9	8.2	9.0	8.4	9.7	11.5	10.8	9.9	10.4
.7	.5	1.0	.9	1.2	.9	1.1	1.0	1.8	1.7	1.9	2.3
1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.4	1.5	1.4	1.5	1.5
.6	.6	.6	.5	.6	.6	.6	.6	.7	.7	.7	.8
4.0	6.0	5.0	4.0	5.0	5.0	7.0	6.0	5.0	6.0	4.0	6.0
56	47	68	51	76	54	71	55	65	55	59	55
37	29	45	31	49	33	45	35	41	35	36	37
19	17	19	19	19	17	20	19	20	18	16	18
32	28	37	30	38	32	38	32	37	31	32	32
43	37	56	42	65	45	60	44	56	42	50	46
71	61	93	65	97	66	91	68	85	70	75	71
109	82	129	93	140	98	129	97	132	97	118	94
124	104	159	109	170	119	151	118	151	124	131	122
187	141	208	162	241	173	231	175	189	187	151	253
141	120	166	133	186	132	173	134	148	126	139	125
80	61	104	73	109	70	94	77	90	71	82	66
136	115	161	125	178	127	168	125	148	121	118	121
67	50	68	52	82	55	78	60	63	54	48	71
133	122	160	96	132	125	161	116	115	125	119	100
43	64	64	38	63	44	66	47	31	45	38	34
681	509	1,282	582	990	595	968	968	568	560	454	531
867	837	1,282	634	1,029	709	1,136	1,132	673	691	638	782
131	112	153	123	169	124	158	121	134	114	115	108
75	58	85	68	101	66	88	69	70	61	58	60
51	43	54	43	56	43	56	43	51	43	46	38
84	70	97	82	98	84	97	84	86	80	84	62
112	101	130	112	149	112	140	111	113	102	112	97
170	140	194	152	220	164	200	142	170	138	131	114
224	194	279	220	291	209	276	209	227	194	173	194
291	222	320	254	346	246	321	228	280	227	209	227
385	302	420	388	485	336	440	376	340	340	284	340
681	485	679	520	990	595	968	968	456	454	454	393
4.5	5.7	5.5	3.3	19.1	22.1	20.0	6.5	.8	6.5	2.2	3.0
.4	.5	.7	.4	.9	.7	.7	.2	.1	.6	.7	.5

TABLE 7.16.--CHICKEN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	38.7	15.6	38.8	39.3	41.4
ON ONLY 1 OF 3 DAYS.....PCT.....	31.2	10.5	30.7	32.7	34.9
ON ONLY 2 OF 3 DAYS.....PCT.....	6.8	4.1	6.9	6.2	6.0
ON ALL 3 OF 3 DAYS.....PCT.....	.6	1.1	1.2	.4	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.5	1.3	1.2	1.2
STANDARD DEVIATION.....DO.....	.6	.8	.6	.5	.5
MAXIMUM.....DO.....	7.0	4.0	4.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	56	32	32	37	43
STANDARD DEVIATION.....GM.....	36	26	22	23	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	17	5	9	12	14
25TH.....GM.....	32	14	17	21	27
50TH.....GM.....	45	24	27	32	37
75TH.....GM.....	70	38	37	43	55
90TH.....GM.....	102	58	65	65	75
95TH.....GM.....	129	85	81	78	99
99TH.....GM.....	185	137	103	114	129
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	141	65	76	97	111
STANDARD DEVIATION.....GM.....	83	39	42	52	58
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	66	83	93	105
STANDARD DEVIATION.....GM.....	66	37	41	46	46
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	124	81	60	88	121
STANDARD DEVIATION.....GM.....	55	44	33	25	46
MAXIMUM IN ANY 1 DAY.....GM.....	1,282	213	336	485	390
MAXIMUM OVER 3 DAYS.....GM.....	1,282	411	418	582	630
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	131	62	73	92	106
STANDARD DEVIATION.....GM.....	76	39	43	50	55
PERCENTILES: <u>1/</u>					
5TH.....GM.....	43	14	20	28	38
25TH.....GM.....	84	35	40	56	61
50TH.....GM.....	112	56	61	89	97
75TH.....GM.....	170	85	101	112	116
90TH.....GM.....	224	128	113	150	194
95TH.....GM.....	280	128	131	194	220
99TH.....GM.....	388	135	222	234	291
MAXIMUM.....GM.....	990	213	336	448	390
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	80.2	1.7	1.6	4.0	5.9
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	1.4	.3	.5	.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
39.8	39.6	38.9	39.5	34.7	39.3	38.8	40.3	38.7	41.4	33.3	37.8
33.7	33.4	32.5	34.4	27.6	31.6	31.9	32.1	28.5	31.2	24.8	26.3
6.0	6.0	6.1	4.9	6.5	7.2	6.2	7.6	9.1	8.9	6.9	9.4
.2	.2	.3	.2	.6	.5	.7	.6	1.1	1.3	1.6	2.1
1.2	1.2	1.2	1.2	1.3	1.3	1.3	1.3	1.4	1.4	1.5	1.5
.5	.5	.5	.4	.5	.5	.6	.6	.6	.7	.7	.7
3.0	4.0	4.0	4.0	3.0	5.0	7.0	6.0	5.0	6.0	4.0	6.0
55	47	66	51	75	53	69	52	63	53	58	55
35	27	43	29	46	32	42	32	40	35	34	36
19	18	20	19	27	18	23	19	19	18	16	18
32	28	37	32	41	32	38	32	34	31	32	32
42	37	56	41	65	43	60	42	50	42	48	41
70	59	86	65	97	65	87	65	80	70	75	71
102	78	129	87	135	95	124	93	129	94	115	93
122	102	153	102	166	113	150	113	151	122	131	113
173	138	191	162	228	168	202	161	187	187	151	253
144	121	170	136	193	135	177	133	148	126	139	127
79	60	103	73	106	69	91	74	90	72	82	66
140	118	167	129	175	126	164	123	145	124	135	117
72	50	72	50	83	54	75	56	62	51	44	70
117	118	154	98	140	135	169	111	122	127	106	102
31	17	29	17	57	44	67	49	35	50	25	35
681	485	1,282	582	990	520	907	880	568	560	454	531
867	539	1,282	582	990	709	1,136	907	582	691	500	782
136	115	160	128	177	127	162	122	134	116	120	109
77	57	87	70	99	65	85	67	70	60	58	59
52	45	56	51	60	49	57	47	51	45	49	38
85	78	97	85	111	84	97	84	85	80	85	65
112	105	140	112	162	112	148	112	112	104	112	97
170	140	200	164	224	168	209	149	170	140	145	114
224	194	284	224	304	209	279	209	227	194	182	189
291	224	328	255	352	244	320	227	280	227	224	226
388	284	448	388	485	336	440	352	340	336	454	340
681	485	679	520	990	520	907	680	456	454	454	393
3.2	2.8	3.7	2.5	13.4	19.3	9.7	3.5	.8	4.8	2.2	1.1
.3	.3	.6	.4	.8	.7	.4	.1	.1	.5	.8	.2

TABLE 7.17.--TURKEY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.8	5.1	4.4	4.5	5.7
ON ONLY 1 OF 3 DAYS.....PCT.....	4.9	4.6	3.9	3.8	4.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	.2	.4	.7	.7
ON ALL 3 OF 3 DAYS.....PCT.....	.1	.2	.1	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	1.3	1.2	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.6	.5	.5	.5	.5
MAXIMUM.....DO.....	5.0	3.0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	45	21	30	29	30
STANDARD DEVIATION.....GM.....	35	15	23	17	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	9			9	9
25TH.....GM.....	19			19	19
50TH.....GM.....	37			28	28
75TH.....GM.....	57			38	38
90TH.....GM.....	85			53	54
95TH.....GM.....	113			75	60
99TH.....GM.....	161			84	106
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	114	58	79	78	80
STANDARD DEVIATION.....GM.....	85	42	64	43	46
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	116	50	84	73	62
STANDARD DEVIATION.....GM.....	66	0	37	30	29
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	92	47	54	0	67
STANDARD DEVIATION.....GM.....	40	0	0	0	33
MAXIMUM IN ANY 1 DAY.....GM.....	968	142	318	248	318
MAXIMUM OVER 3 DAYS.....GM.....	968	142	318	253	350
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	105	53	73	74	74
STANDARD DEVIATION.....GM.....	73	34	59	39	44
PERCENTILES: ^{1/}					
5TH.....GM.....	28			28	28
25TH.....GM.....	57			56	43
50TH.....GM.....	86			57	57
75TH.....GM.....	129			85	86
90TH.....GM.....	172			140	129
95TH.....GM.....	240			159	168
99TH.....GM.....	350			248	224
MAXIMUM.....GM.....	968	142	318	248	318
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	26.8	0	.9	0	1.8
PROPORTION OF EATING OCCASIONS.....PCT.....	1.0	0	1.6	0	1.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.5	6.2	7.5	6.2	6.7	5.5	6.1	6.1	4.3	4.8	3.7	3.8
5.7	5.4	6.5	4.9	5.5	4.8	5.1	5.0	3.2	3.5	1.7	3.7
.7	.7	.8	1.0	.9	.6	.9	.9	.5	1.1	2.0	.2
.1	(2)	.1	.3	.2	.1	.1	.1	.6	.2	0	0
1.2	1.2	1.2	1.3	1.3	1.2	1.2	1.3	1.4	1.4	1.7	1.1
.5	.4	.5	.6	.6	.5	.5	.6	.7	.7	.7	.3
3.0	3.0	4.0	4.0	4.0	4.0	5.0	4.0	3.0	4.0	3.0	2.0
42	36	49	39	52	39	54	49	60	48	48	33
28	26	39	27	44	26	40	39	31	33	20	20
14	9	14	11	9	9	14	9		14		
23	19	27	23	23	19	28	28		28		
29	28	38	29	38	35	43	38		38		
55	43	57	43	72	57	75	57		56		
84	57	104	76	103	76	113	95		105		
100	86	140	106	129	86	124	113		132		
123	144	208	151	261	106	173	233		152		
106	97	120	95	125	107	139	125	150	122	119	94
59	73	80	58	87	66	105	105	72	86	66	56
129	88	153	102	164	100	136	120	115	82	84	99
52	37	74	55	102	50	58	57	38	37	20	0
95	86	152	66	78	66	88	107	105	134	0	0
28	0	76	8	28	0	57	30	20	17	0	0
344	509	484	484	562	340	968	968	454	454	284	227
547	509	680	484	842	480	968	968	454	455	284	227
103	90	120	89	119	99	129	110	129	102	86	90
56	54	68	47	82	63	92	80	59	63	46	54
43	28	42	28	28	28	43	28		36		
57	57	70	57	57	56	84	57		57		
86	85	112	85	113	85	113	86		86		
113	112	142	113	159	113	170	113		113		
170	159	227	159	227	170	227	171		140		
227	172	258	210	258	247	284	240		170		
327	350	340	242	454	318	363	376		454		
327	432	340	248	454	340	968	968	454	454	284	227
0	.9	1.8	.8	4.5	2.8	6.7	3.0	0	1.7	0	1.9
0	.6	1.4	.7	1.3	.8	1.8	.5	0	1.5	0	4.6

TABLE 7.18.--LIVER
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.8	1.5	3.5	2.9	2.8
ON ONLY 1 OF 3 DAYS.....PCT.....	3.6	1.4	3.3	2.8	2.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.2	.2	.1	.1	.1
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.1	1.1	1.0	1.0
STANDARD DEVIATION.....DO.....	.3	.3	.2	.2	.2
MAXIMUM.....DO.....	3.0	2.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	43	29	16	24	26
STANDARD DEVIATION.....GM.....	28	20	10	15	14
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9			6	9
25TH.....GM.....	25			14	19
50TH.....GM.....	37			19	28
75TH.....GM.....	56			35	37
90TH.....GM.....	75			37	38
95TH.....GM.....	90			56	47
99TH.....GM.....	149			75	75
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	123	67	49	71	76
STANDARD DEVIATION.....GM.....	80	36	30	44	38
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	102	112	28	56	72
STANDARD DEVIATION.....GM.....	49	0	0	0	20
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	112	0	0	0	0
STANDARD DEVIATION.....GM.....	0	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	907	128	126	224	224
MAXIMUM OVER 3 DAYS.....GM.....	907	224	126	224	224
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	118	77	46	71	75
STANDARD DEVIATION.....GM.....	76	37	27	43	37
PERCENTILES: <u>1/</u>					
5TH.....GM.....	27			18	28
25TH.....GM.....	70			43	56
50TH.....GM.....	112			56	75
75TH.....GM.....	150			86	100
90TH.....GM.....	224			112	113
95TH.....GM.....	227			168	113
99TH.....GM.....	360			224	224
MAXIMUM.....GM.....	907	128	113	224	224
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	42.9	0	1.1	1.7	0
PROPORTION OF EATING OCCASIONS.....PCT.....	2.8	0	2.9	3.4	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.8	2.3	3.6	2.1	3.3	2.9	4.6	4.8	5.8	5.8	4.2	5.2
2.8	2.3	3.5	2.1	3.3	2.9	4.4	4.5	5.4	4.7	4.2	4.2
0	0	.1	0	(2)	(2)	.2	.4	.4	1.0	0	1.0
0	0	0	0	0	0	0	(2)	0	0	0	0
1.1	1.0	1.0	1.0	1.0	1.0	1.1	1.1	1.1	1.3	1.0	1.3
.2	.0	.2	.0	.1	.1	.3	.3	.3	.5	.0	.4
2.0	1.0	2.0	1.0	2.0	2.0	2.0	3.0	2.0	3.0	1.0	2.0
35	30	44	34	56	42	49	43	44	49	44	41
21	17	27	19	33	37	25	25	23	30	23	27
8	7	9		12	9	18	13	9	9		9
19	19	28		33	28	33	28	28	28		21
28	28	37		48	37	47	37	37	37		37
47	37	56		75	50	57	56	56	75		56
57	56	76		93	75	76	75	75	76		66
68	56	93		113	75	93	82	93	105		75
112	93	131		149	302	149	149	112	149		151
105	91	131	103	167	126	144	121	128	126	131	116
63	50	83	57	100	111	71	66	66	80	68	88
0	0	57	0	98	156	116	106	82	123	0	76
0	0	0	0	0	0	53	54	50	38	0	21
0	0	0	0	0	0	0	112	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
336	280	392	336	448	907	492	448	336	448	280	454
336	280	392	336	448	907	504	504	336	448	280	454
99	91	128	103	166	125	138	118	120	115	131	98
61	50	82	57	99	110	70	66	61	57	68	52
28	22	28		35	20	43	25	26	36		43
56	56	57		98	84	86	84	84	70		57
84	84	112		140	112	125	112	112	112		85
112	112	168		224	150	168	150	168	140		112
168	168	227		280	198	224	224	196	224		168
204	168	280		340	224	280	227	224	224		227
336	280	392		448	907	448	360	280	324		227
336	280	392	336	448	907	492	448	280	324	280	227
.7	0	.7	1.3	2.9	6.7	9.4	14.1	2.2	0	1.0	1.0
1.1	0	1.4	4.3	2.2	4.2	3.9	3.8	3.1	0	4.5	1.6

TABLE 7.19.--FISH AND SHELLFISH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	24.5	2.3	17.3	20.4	22.5
ON ONLY 1 OF 3 DAYS.....PCT.....	20.5	2.2	14.2	18.2	19.9
ON ONLY 2 OF 3 DAYS.....PCT.....	3.6	.2	2.5	2.1	2.2
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	.6	.2	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.1	1.3	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.3	.7	.5	.4
MAXIMUM.....DO.....	6.0	2.0	5.0	5.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	48	15	22	27	32
STANDARD DEVIATION.....GM.....	42	9	18	19	24
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	8		3	4	6
25TH.....GM.....	20		9	13	19
50TH.....GM.....	37		19	25	28
75TH.....GM.....	57		28	37	38
90TH.....GM.....	94		47	49	57
95TH.....GM.....	128		56	57	75
99TH.....GM.....	215		93	96	131
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	124	48	55	75	86
STANDARD DEVIATION.....GM.....	106	26	40	53	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	120	7	56	65	83
STANDARD DEVIATION.....GM.....	83	0	38	37	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	111	0	54	59	56
STANDARD DEVIATION.....GM.....	53	0	10	12	15
MAXIMUM IN ANY 1 DAY.....GM.....	2,350	91	228	432	524
MAXIMUM OVER 3 DAYS.....GM.....	2,350	91	324	432	580
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	117	42	52	70	81
STANDARD DEVIATION.....GM.....	98	28	38	51	58
PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		8	12	19
25TH.....GM.....	57		28	36	40
50TH.....GM.....	85		43	57	72
75TH.....GM.....	152		58	85	112
90TH.....GM.....	227		112	113	160
95TH.....GM.....	284		125	170	170
99TH.....GM.....	456		168	240	288
MAXIMUM.....GM.....	1,410	91	228	432	448
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	736.0	.8	8.3	23.1	34.5
PROPORTION OF EATING OCCASIONS.....PCT.....	6.5	6.1	3.6	5.7	7.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
22.0	22.9	21.1	24.2	25.1	25.7	26.9	28.0	27.5	24.7	20.7	19.6
19.1	20.4	18.6	20.7	20.5	20.9	21.9	23.2	21.7	20.6	17.9	17.1
2.5	2.1	2.3	3.2	4.3	4.2	4.4	4.4	5.0	3.8	2.0	2.1
.3	.3	.2	.4	.3	.6	.6	.3	.7	.3	.9	.3
1.2	1.2	1.2	1.2	1.3	1.3	1.3	1.2	1.3	1.2	1.2	1.2
.5	.4	.5	.5	.5	.6	.6	.5	.5	.5	.5	.4
4.0	4.0	4.0	3.0	4.0	6.0	6.0	4.0	4.0	4.0	3.0	3.0
40	33	46	45	62	44	62	49	62	50	51	45
31	25	45	42	58	35	49	40	47	41	28	28
9	8	9	9	11	8	13	7	12	9	18	7
19	19	27	19	27	20	28	23	33	25	28	24
30	28	37	33	47	37	51	38	49	38	47	38
47	38	57	57	80	56	76	61	76	61	71	57
76	61	84	85	128	85	123	93	121	93	89	76
96	82	113	112	165	112	151	122	151	119	112	97
154	151	297	217	224	180	245	224	253	215	114	141
107	90	129	121	157	110	157	127	160	126	144	121
81	62	117	112	156	84	116	105	106	86	77	74
100	94	109	105	153	102	148	119	138	130	103	105
59	63	109	87	96	60	95	72	108	96	49	42
73	49	80	91	134	122	152	102	110	149	87	101
29	26	32	48	33	42	66	29	32	50	15	38
840	567	1,344	932	2,350	758	996	830	768	624	432	450
840	582	1,429	1,158	2,350	758	1,087	853	1,152	900	432	450
101	86	117	111	149	104	147	119	145	123	124	112
78	62	115	102	125	74	116	98	109	87	68	69
28	19	20	24	28	20	28	20	35	24	36	20
56	45	57	56	64	57	80	57	75	61	80	61
84	79	85	85	113	85	113	85	113	103	106	112
113	112	142	130	196	135	180	152	180	168	170	151
170	168	200	225	284	184	258	227	270	227	227	196
255	206	252	270	362	227	360	280	392	304	227	225
425	288	454	568	643	394	577	480	480	448	336	360
840	567	1,344	864	1,410	758	996	830	768	624	336	450
58.8	35.3	21.7	34.7	59.8	78.2	143.4	173.7	14.4	31.6	6.2	11.7
10.8	6.2	6.2	8.0	4.8	4.5	8.5	7.1	3.7	6.0	4.5	5.1

TABLE 7.20.--FINFISH OTHER THAN CANNED, DRIED, AND RAW
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	14.5	1.7	10.7	13.2	13.9
ON ONLY 1 OF 3 DAYS.....PCT.....	13.3	1.7	9.2	12.2	13.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	0	1.5	1.0	.7
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.1	1.0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.4	.3	.3
MAXIMUM.....DO.....	5.0	1.0	3.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	54	15	26	31	36
STANDARD DEVIATION.....GM.....	43	9	17	20	22
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	17		9	9	10
25TH.....GM.....	28		14	19	19
50TH.....GM.....	38		19	28	29
75TH.....GM.....	67		37	38	45
90TH.....GM.....	96		47	56	57
95TH.....GM.....	132		56	70	76
99TH.....GM.....	221		93	142	131
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	148	44	70	89	105
STANDARD DEVIATION.....GM.....	113	26	40	55	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	151	0	63	74	94
STANDARD DEVIATION.....GM.....	91	0	38	45	52
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	106	0	0	0	0
STANDARD DEVIATION.....GM.....	48	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	2,350	91	228	432	448
MAXIMUM OVER 3 DAYS.....GM.....	2,350	91	324	432	560
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	145	44	68	85	101
STANDARD DEVIATION.....GM.....	106	26	39	55	59
PERCENTILES: ^{1/}					
5TH.....GM.....	43		21	28	28
25TH.....GM.....	84		38	56	57
50TH.....GM.....	113		57	84	85
75TH.....GM.....	170		85	112	113
90TH.....GM.....	255		113	144	170
95TH.....GM.....	340		144	170	210
99TH.....GM.....	565		168	340	336
MAXIMUM.....GM.....	1,410	91	228	432	448
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	681.0	.8	8.3	23.1	32.6
PROPORTION OF EATING OCCASIONS.....PCT.....	11.1	8.7	6.5	9.3	11.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
15.1	13.8	12.5	14.7	14.0	13.0	16.5	16.2	18.4	16.6	12.5	13.1
13.9	13.1	12.0	13.5	12.4	12.1	14.9	15.0	15.7	14.7	10.7	11.8
1.1	.6	.5	1.1	1.5	1.0	1.5	1.2	2.3	1.9	1.3	1.2
.1	0	0	.1	.1	0	.1	(2)	.3	0	.5	.2
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.1	1.3	1.1
.3	.3	.3	.4	.4	.3	.4	.3	.5	.4	.5	.4
3.0	3.0	3.0	3.0	3.0	5.0	4.0	3.0	3.0	3.0	3.0	3.0
43	36	52	52	72	47	68	55	70	57	58	49
31	24	49	46	61	32	48	39	51	44	27	26
13	12	19	19	19	17	19	19	19	19	19	19
27	19	28	28	37	28	37	28	37	28	38	28
37	28	38	38	57	38	57	47	57	47	57	44
56	38	57	63	90	57	80	70	76	75	76	57
85	66	88	85	132	80	131	93	135	101	98	76
108	84	117	136	169	112	152	128	151	139	113	101
149	144	299	288	267	180	227	221	284	253	144	150
121	103	154	145	198	128	189	154	178	148	168	139
86	59	148	116	171	75	132	106	104	90	77	73
116	125	111	152	190	154	175	147	200	174	101	90
48	64	26	130	96	73	76	78	130	114	51	30
95	0	0	75	57	0	179	85	105	0	86	141
43	0	0	0	0	0	32	0	18	0	21	0
840	454	1,344	932	2,350	565	996	830	768	624	432	450
840	582	1,344	1,158	2,350	663	996	853	1,152	900	432	450
117	102	145	139	191	129	181	150	175	150	137	130
82	61	138	114	136	73	125	102	117	94	69	69
28	28	56	54	57	48	56	55	56	56	56	56
68	57	85	84	112	84	105	85	100	85	85	84
90	85	112	112	168	112	168	120	140	113	113	113
140	113	170	170	227	168	227	170	227	180	170	170
224	170	227	252	340	224	336	252	340	280	227	224
284	227	304	304	432	270	432	336	432	340	280	227
425	324	897	649	649	396	680	589	577	454	336	450
840	454	1,344	864	1,410	540	996	830	768	624	336	450
58.8	28.8	21.7	32.9	52.9	67.7	134.3	160.7	13.0	29.9	6.2	9.5
16.7	9.0	11.4	13.5	8.4	8.8	14.6	12.8	5.3	9.1	7.3	6.5

TABLE 7.21.--CANNED TUNA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.2	.6	6.2	6.2	7.6
ON ONLY 1 OF 3 DAYS.....PCT.....	6.2	.4	5.3	5.4	6.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	.2	.6	.6	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.3	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.2	1.3	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.4	.5	.6	.4	.5
MAXIMUM.....DO.....	4.0	2.0	4.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	28	17	13	18	22
STANDARD DEVIATION.....GM.....	22	8	9	12	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	7		2	3	3
25TH.....GM.....	13		7	7	9
50TH.....GM.....	27		13	13	18
75TH.....GM.....	34		13	27	27
90TH.....GM.....	56		27	33	42
95TH.....GM.....	66		28	43	54
99TH.....GM.....	107		43	57	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	73	65	32	48	52
STANDARD DEVIATION.....GM.....	46	11	19	32	36
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	76	7	30	53	60
STANDARD DEVIATION.....GM.....	52	0	15	21	41
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	80	0	34	28	58
STANDARD DEVIATION.....GM.....	41	0	7	0	16
MAXIMUM IN ANY 1 DAY.....GM.....	480	80	113	170	284
MAXIMUM OVER 3 DAYS.....GM.....	960	80	128	170	340
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	73	39	31	47	54
STANDARD DEVIATION.....GM.....	47	30	18	31	39
PERCENTILES: ^{1/}					
5TH.....GM.....	19		7	10	10
25TH.....GM.....	40		19	20	28
50TH.....GM.....	61		28	40	40
75TH.....GM.....	85		40	61	61
90TH.....GM.....	120		57	80	85
95TH.....GM.....	169		57	99	113
99TH.....GM.....	213		85	170	200
MAXIMUM.....GM.....	480	80	113	170	284
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	16.5	0	0	0	.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	0	0	.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.7	6.9	6.3	7.4	8.0	9.4	6.4	8.7	4.5	5.5	5.8	3.6
5.0	6.4	5.7	6.0	7.1	7.8	5.5	7.4	4.2	5.0	5.8	2.9
.6	.5	.6	1.2	.9	1.3	.8	1.2	.4	.5	0	.7
.1	.1	.1	.1	(2)	.3	.1	.1	0	.1	0	0
1.1	1.1	1.1	1.2	1.2	1.2	1.2	1.2	1.1	1.1	1.0	1.2
.4	.4	.4	.5	.4	.5	.4	.4	.3	.3	.0	.4
3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0	3.0	1.0	2.0
26	23	27	29	35	31	31	28	28	26	36	35
19	14	16	22	35	23	20	19	15	15	21	28
9	7	5	7	9	7	10	7	12	7		
13	13	13	13	18	14	19	13	19	18		
23	19	27	20	27	27	27	27	27	20		
31	28	33	38	40	38	38	33	32	33		
53	47	53	59	66	57	57	53	53	47		
53	53	60	80	95	71	67	61	57	57		
154	66	67	113	184	113	92	99	61	66		
72	64	77	71	90	75	83	71	79	74	107	86
40	38	44	54	60	46	45	40	42	40	62	58
65	59	56	77	106	84	73	77	61	53	0	98
56	27	28	29	112	29	37	38	22	25	0	58
43	63	60	63	184	92	100	102	0	93	0	0
27	3	0	25	0	40	28	16	0	0	0	0
320	170	200	340	480	312	262	320	184	198	198	223
462	217	200	340	960	553	454	640	184	278	198	397
69	62	71	70	91	76	80	71	75	71	107	89
45	37	42	47	70	41	43	39	39	38	62	59
20	19	15	20	20	19	20	20	35	10		
40	40	40	40	40	40	57	40	40	40		
57	57	61	57	80	80	80	61	66	61		
80	80	80	80	113	96	92	85	92	85		
120	113	120	113	170	120	160	113	160	113		
160	160	170	160	198	169	169	142	169	160		
198	170	200	340	480	184	198	184	184	198		
320	170	200	340	480	213	262	320	184	198	198	223
0	1.7	0	0	3.9	4.4	0	4.4	0	0	0	1.3
0	1.0	0	0	1.1	.7	0	.6	0	0	0	3.0

TABLE 7.22.--SHRIMP
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.5	0	1.0	1.0	1.1
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	0	1.0	1.0	1.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.1	0	0	0	0
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	0	1.0	1.0	1.0
STANDARD DEVIATION.....DO.....	.3	0	.0	.0	.2
MAXIMUM.....DO.....	3.0	0	1.0	1.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	24	0	7	10	14
STANDARD DEVIATION.....GM.....	30	0	10	11	12
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3				
25TH.....GM.....	8				
50TH.....GM.....	14				
75TH.....GM.....	27				
90TH.....GM.....	47				
95TH.....GM.....	76				
99TH.....GM.....	151				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	66	0	22	31	42
STANDARD DEVIATION.....GM.....	76	0	31	34	37
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	89	0	0	0	0
STANDARD DEVIATION.....GM.....	102	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	30	0	0	0	0
STANDARD DEVIATION.....GM.....	0	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	907	0	135	135	135
MAXIMUM OVER 3 DAYS.....GM.....	971	0	135	135	135
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	67	0	22	31	41
STANDARD DEVIATION.....GM.....	82	0	31	34	37
PERCENTILES: <u>1/</u>					
5TH.....GM.....	8				
25TH.....GM.....	22				
50TH.....GM.....	40				
75TH.....GM.....	70				
90TH.....GM.....	135				
95TH.....GM.....	227				
99TH.....GM.....	454				
MAXIMUM.....GM.....	907	0	135	135	135
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	11.9	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.0	2.1	2.9	2.2	2.8	3.3	3.1	3.4	2.7	1.4	.7	1.5
1.0	2.1	2.6	2.2	2.7	3.0	2.9	3.2	2.5	1.4	.7	1.5
0	(2)	.2	0	.1	.3	.2	.1	.2	0	0	0
0	0	.1	0	0	0	0	0	0	0	0	0
1.0	1.0	1.1	1.0	1.1	1.1	1.1	1.1	1.2	1.0	1.0	1.1
.0	.1	.4	.0	.2	.3	.3	.2	.5	.0	.0	.3
1.0	2.0	3.0	1.0	2.0	2.0	2.0	2.0	3.0	1.0	1.0	2.0
21	15	32	23	27	22	29	22	36	23	24	14
16	15	32	24	33	25	40	29	40	16	23	8
				2	3	3	3				
				11	8	8	5				
				16	16	19	11				
				32	27	34	23				
				57	45	50	46				
				81	76	76	97				
				218	151	197	151				
63	45	96	68	71	63	75	63	105	69	72	42
49	46	101	71	69	72	79	82	126	47	70	25
0	30	49	0	178	56	143	72	64	0	0	0
0	0	14	0	115	44	161	73	0	0	0	0
0	0	30	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
227	227	454	340	453	454	907	454	454	135	170	97
227	227	454	340	654	454	971	486	454	135	170	97
63	44	87	68	78	62	82	62	93	69	72	38
49	45	94	71	79	69	111	79	116	47	70	17
				6	6	8	8				
				32	24	24	16				
				48	43	57	32				
				101	68	85	68				
				170	135	135	135				
				227	169	227	227				
				453	454	561	340				
227	227	454	340	453	454	907	454	454	135	170	68
0	2.7	0	.9	0	1.2	5.2	1.0	0	0	0	.9
0	5.9	0	2.7	0	.6	3.2	.4	0	0	0	5.6

TABLE 7.23.--FRANKFURTERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.8	2.9	28.6	29.7	27.8
ON ONLY 1 OF 3 DAYS.....PCT.....	15.4	2.1	22.5	23.7	24.1
ON ONLY 2 OF 3 DAYS.....PCT.....	2.1	.3	5.2	5.2	3.1
ON ALL 3 OF 3 DAYS.....PCT.....	.3	.5	.9	.9	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.5	1.3	1.3	1.2
STANDARD DEVIATION.....DO.....	.5	.8	.6	.5	.5
MAXIMUM.....DO.....	7.0	3.0	3.0	4.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	29	16	24	24	25
STANDARD DEVIATION.....GM.....	19	14	17	16	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		7	9	14
25TH.....GM.....	15		15	15	15
50TH.....GM.....	29		15	15	19
75TH.....GM.....	29		29	29	29
90TH.....GM.....	52		44	44	44
95TH.....GM.....	59		59	48	57
99TH.....GM.....	103		95	89	88
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	76	28	55	58	63
STANDARD DEVIATION.....GM.....	41	14	32	31	29
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	77	35	60	57	69
STANDARD DEVIATION.....GM.....	37	8	29	25	31
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	78	40	57	60	63
STANDARD DEVIATION.....GM.....	45	15	26	37	30
MAXIMUM IN ANY 1 DAY.....GM.....	660	57	280	280	308
MAXIMUM OVER 3 DAYS.....GM.....	836	157	352	567	484
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	75	33	55	56	62
STANDARD DEVIATION.....GM.....	40	15	29	29	30
PERCENTILES: <u>1/</u>					
5TH.....GM.....	43		22	28	35
25TH.....GM.....	44		44	44	44
50TH.....GM.....	57		44	44	44
75TH.....GM.....	88		57	57	88
90TH.....GM.....	132		88	88	88
95TH.....GM.....	142		113	112	113
99TH.....GM.....	220		176	170	170
MAXIMUM.....GM.....	454	57	280	280	280
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	13.3	0	0	0	2.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
25.9	24.9	20.8	18.9	15.9	18.7	13.9	13.8	13.2	9.0	9.7	11.4
22.4	21.9	17.7	16.6	13.9	15.9	12.4	12.7	12.0	8.0	8.3	9.4
3.1	2.6	3.1	2.0	1.8	2.6	1.2	1.0	.9	1.0	1.2	1.7
.4	.3	.1	.2	.3	.2	.2	.1	.3	.1	.2	.3
1.2	1.1	1.2	1.2	1.2	1.2	1.1	1.1	1.1	1.2	1.2	1.3
.5	.4	.4	.5	.4	.4	.4	.3	.4	.4	.6	.8
4.0	3.0	4.0	4.0	3.0	4.0	4.0	3.0	3.0	3.0	4.0	7.0
29	25	36	26	40	28	36	26	33	28	35	27
19	16	21	16	23	19	24	15	19	15	35	16
15	12	15	15	15	15	15	15	15	15	12	15
15	15	19	15	29	15	22	15	28	15	19	15
29	19	29	23	29	29	29	29	29	29	29	29
29	29	44	29	47	29	44	29	37	29	37	29
48	44	59	44	59	47	59	44	44	56	66	44
59	59	76	59	88	59	76	57	76	59	76	59
113	76	113	113	117	103	132	88	113	67	235	103
77	65	93	68	106	71	95	72	88	75	82	69
42	36	49	29	56	33	46	34	35	38	44	27
73	70	94	78	104	82	91	76	95	70	73	59
35	25	38	38	41	39	39	31	62	25	30	27
88	65	73	56	77	74	149	73	81	59	235	95
40	22	0	11	19	31	60	19	30	0	0	7
454	444	352	227	660	284	454	264	255	179	308	176
572	444	352	340	660	425	836	340	340	264	704	308
75	65	90	67	102	73	95	71	88	71	84	65
41	35	42	30	48	36	48	32	43	31	47	27
44	43	44	44	44	43	44	44	44	44	43	44
44	44	44	44	85	44	57	44	44	44	44	44
57	44	88	56	88	66	88	57	88	66	88	44
88	88	113	88	132	88	113	88	88	88	88	88
132	88	140	88	176	113	132	112	132	99	170	88
132	132	176	132	176	132	176	132	176	132	176	112
220	176	220	170	227	176	264	176	255	176	227	142
454	444	264	227	440	284	454	227	255	179	227	142
3.4	.8	0	0	0	0	.8	3.3	0	2.6	0	0
.5	.1	0	0	0	0	.1	.3	0	1.4	0	0

TABLE 7.24.--BOLOGNA
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	15.9	2.4	20.4	23.8	22.1
ON ONLY 1 OF 3 DAYS.....PCT.....	11.7	1.7	14.2	18.3	15.3
ON ONLY 2 OF 3 DAYS.....PCT.....	3.2	.5	4.4	4.1	5.3
ON ALL 3 OF 3 DAYS.....PCT.....	1.0	.2	1.8	1.4	1.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.4	1.4	1.4	1.3	1.4
STANDARD DEVIATION.....DO.....	.6	.6	.7	.6	.6
MAXIMUM.....DO.....	6.0	3.0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	21	19	15	16	18
STANDARD DEVIATION.....GM.....	16	22	11	11	12
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	7		5	5	9
25TH.....GM.....	9		9	9	9
50TH.....GM.....	19		9	9	19
75TH.....GM.....	28		19	19	19
90TH.....GM.....	38		28	28	37
95TH.....GM.....	56		37	38	38
99TH.....GM.....	76		75	65	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	49	27	32	37	42
STANDARD DEVIATION.....GM.....	32	18	16	20	23
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	48	82	37	37	39
STANDARD DEVIATION.....GM.....	26	40	23	20	19
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	47	13	27	36	34
STANDARD DEVIATION.....GM.....	24	0	11	16	12
MAXIMUM IN ANY 1 DAY.....GM.....	312	112	140	142	170
MAXIMUM OVER 3 DAYS.....GM.....	448	224	227	280	227
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	47	41	32	36	40
STANDARD DEVIATION.....GM.....	29	37	18	20	21
PERCENTILES: ^{1/}					
5TH.....GM.....	14		14	14	14
25TH.....GM.....	28		28	28	28
50TH.....GM.....	28		28	28	28
75TH.....GM.....	57		28	43	56
90TH.....GM.....	85		57	57	57
95TH.....GM.....	112		57	84	84
99TH.....GM.....	168		112	112	113
MAXIMUM.....GM.....	284	112	113	140	140
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	14.5	0	.9	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	.3	0	0

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.7	21.3	20.2	15.9	19.1	15.0	16.5	11.5	12.8	8.3	13.3	7.0
14.8	15.9	15.3	12.5	12.7	11.7	11.3	9.2	8.9	5.9	11.1	4.8
4.0	4.3	4.0	2.8	4.6	2.7	3.7	1.9	2.9	2.1	1.5	1.5
.9	1.1	.9	.6	1.8	.6	1.5	.4	.9	.3	.7	.6
1.3	1.3	1.3	1.3	1.5	1.3	1.4	1.3	1.4	1.4	1.2	1.4
.6	.6	.6	.6	.8	.5	.7	.6	.6	.6	.5	.7
4.0	5.0	6.0	4.0	5.0	3.0	4.0	4.0	3.0	3.0	3.0	4.0
20	18	26	19	29	19	27	18	22	19	25	20
14	13	19	14	21	14	19	13	13	11	19	13
7	7	9	9	9	5	9	5	9	9	9	9
9	9	19	9	19	9	19	9	9	9	9	9
19	19	19	19	19	19	19	19	19	19	19	19
28	19	37	19	37	19	37	19	28	19	38	28
38	37	56	37	56	38	56	37	38	37	38	38
47	47	69	47	75	47	65	42	47	38	47	47
76	75	113	75	104	66	76	66	75	56	104	57
49	42	62	46	66	44	62	43	50	45	66	49
32	24	37	28	40	27	39	25	21	23	56	34
47	41	59	44	56	47	58	46	50	42	62	38
23	19	34	21	29	25	29	25	26	19	28	17
38	43	60	56	63	43	51	46	36	31	39	35
15	21	34	27	26	17	26	15	9	11	9	10
284	227	258	255	282	170	255	227	141	113	312	170
284	255	448	336	392	340	420	340	224	170	312	170
46	41	60	45	59	44	57	43	48	42	60	42
28	23	36	27	34	26	34	24	23	22	40	27
14	14	28	21	28	14	28	14	21	14	28	14
28	28	28	28	28	28	28	28	28	28	28	28
28	28	56	28	56	28	56	28	56	28	56	28
57	56	57	56	57	56	57	57	57	56	85	56
85	64	112	84	112	84	112	57	84	70	113	84
113	85	113	112	113	112	113	85	85	84	113	85
113	113	170	113	170	140	170	113	113	113	227	170
284	227	227	255	280	170	255	227	113	113	227	170
3.3	1.1	0	3.0	.8	0	1.2	2.3	0	1.9	0	0
.6	.2	0	1.0	.1	0	.1	.2	0	1.0	0	0

TABLE 7.25.--SALAMI

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.4	.2	1.9	3.0	4.3
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	.2	1.5	2.4	3.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.3	.5	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.3	1.0	1.3	1.3	1.3
STANDARD DEVIATION.....DO.....	.6	.0	.6	.5	.7
MAXIMUM.....DO.....	5.0	1.0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	25	9	15	18	21
STANDARD DEVIATION.....GM.....	20	0	9	14	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	7			3	9
25TH.....GM.....	9			9	9
50TH.....GM.....	19			9	15
75TH.....GM.....	37			28	28
90TH.....GM.....	47			38	38
95TH.....GM.....	65			38	47
99TH.....GM.....	95			65	93
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	60	28	40	43	46
STANDARD DEVIATION.....GM.....	42	0	22	31	26
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	60	0	22	39	53
STANDARD DEVIATION.....GM.....	35	0	6	16	33
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	54	0	38	56	62
STANDARD DEVIATION.....GM.....	31	0	0	9	32
MAXIMUM IN ANY 1 DAY.....GM.....	284	28	112	113	140
MAXIMUM OVER 3 DAYS.....GM.....	420	28	113	196	280
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	58	28	35	43	47
STANDARD DEVIATION.....GM.....	39	0	20	28	22
PERCENTILES: ^{1/}					
5TH.....GM.....	20			10	25
25TH.....GM.....	28			28	28
50TH.....GM.....	56			28	42
75TH.....GM.....	57			57	57
90TH.....GM.....	113			84	84
95TH.....GM.....	113			113	85
99TH.....GM.....	224			113	113
MAXIMUM.....GM.....	284	28	112	113	113
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	4.8	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.6	4.3	5.7	4.5	6.8	3.8	7.7	3.3	3.5	1.9	2.5	2.4
3.0	3.1	4.8	4.1	5.0	3.3	5.5	2.8	2.4	1.5	2.1	1.8
.5	1.0	.7	.2	1.6	.4	1.5	.4	.7	.3	.2	.4
.2	.2	.2	.2	.2	(2)	.7	.1	.3	.1	.2	.3
1.3	1.3	1.3	1.2	1.3	1.1	1.4	1.2	1.5	1.3	1.3	1.4
.6	.6	.6	.5	.6	.4	.8	.4	.8	.6	.6	.7
3.0	3.0	3.0	3.0	4.0	3.0	5.0	3.0	4.0	3.0	3.0	3.0
26	23	30	22	29	20	31	18	31	16	23	16
19	16	21	14	19	14	26	13	25	10	21	11
9	5	7	7	9	5	9	5				
13	9	19	9	19	9	14	9				
19	19	19	19	19	19	25	14				
28	28	38	28	38	22	38	19				
56	38	56	38	56	38	75	37				
76	47	66	47	75	56	76	43				
93	75	112	75	84	76	132	57				
69	55	77	58	74	54	68	45	77	34	51	37
53	35	47	32	46	36	49	28	59	15	38	31
54	56	58	55	58	51	77	53	76	52	28	38
36	31	36	17	26	29	44	23	52	10	0	7
48	28	88	51	65	47	59	41	32	28	76	28
11	0	19	20	29	0	36	16	7	0	0	0
280	227	170	142	252	227	284	170	243	84	113	112
280	340	336	225	336	227	420	227	340	113	227	112
59	52	71	57	67	53	65	46	65	38	54	35
44	33	44	32	41	35	43	27	54	15	34	24
28	25	14	25	28	14	20	14				
28	28	28	28	28	28	28	28				
56	43	57	56	56	56	57	40				
57	57	112	57	85	57	85	57				
112	85	142	112	113	85	113	84				
113	140	168	140	140	112	140	112				
280	142	170	142	227	227	224	168				
280	170	170	142	252	227	284	170	243	84	113	112
0	0	0	0	1.7	0	1.2	0	0	1.8	0	0
0	0	0	0	.5	0	.2	0	0	4.2	0	0

TABLE 7.26.--PORK BREAKFAST SAUSAGE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	13.5	1.4	11.9	13.4	12.8
ON ONLY 1 OF 3 DAYS.....PCT.....	10.6	1.1	8.8	10.4	10.4
ON ONLY 2 OF 3 DAYS.....PCT.....	2.3	.3	2.8	2.8	2.1
ON ALL 3 OF 3 DAYS.....PCT.....	.6	0	.3	.2	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.2	1.3	1.3	1.2
STANDARD DEVIATION.....DO.....	.6	.4	.5	.5	.5
MAXIMUM.....DO.....	5.0	2.0	3.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	24	19	15	16	18
STANDARD DEVIATION.....GM.....	21	16	14	14	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	4		4	4	4
25TH.....GM.....	9		7	9	9
50TH.....GM.....	18		9	9	13
75TH.....GM.....	28		18	19	22
90TH.....GM.....	47		28	37	38
95TH.....GM.....	64		38	47	57
99TH.....GM.....	113		72	76	75
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	59	50	33	42	47
STANDARD DEVIATION.....GM.....	49	52	20	35	37
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	55	40	37	32	40
STANDARD DEVIATION.....GM.....	42	15	32	19	24
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	46	0	33	66	50
STANDARD DEVIATION.....GM.....	27	0	13	18	20
MAXIMUM IN ANY 1 DAY.....GM.....	452	160	162	227	227
MAXIMUM OVER 3 DAYS.....GM.....	510	160	324	255	255
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	55	46	34	39	44
STANDARD DEVIATION.....GM.....	45	42	26	31	34
PERCENTILES: <u>1/</u>					
5TH.....GM.....	13		10	13	13
25TH.....GM.....	26		14	22	26
50TH.....GM.....	52		27	27	28
75TH.....GM.....	57		43	54	57
90TH.....GM.....	113		57	85	85
95TH.....GM.....	142		85	85	113
99TH.....GM.....	227		113	170	170
MAXIMUM.....GM.....	452	160	162	227	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	37.0	0	0	0	3.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	0	1.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
16.3	13.2	14.3	13.4	16.1	11.0	17.4	12.0	16.8	10.1	15.4	11.4
13.7	11.0	11.6	11.4	12.1	8.9	13.3	9.4	12.5	8.8	12.5	7.6
2.0	1.7	1.9	1.4	3.3	1.7	3.1	2.1	3.0	1.0	2.3	2.2
.5	.5	.8	.5	.8	.3	1.0	.5	1.2	.3	.6	1.5
1.2	1.2	1.3	1.2	1.3	1.2	1.3	1.3	1.4	1.2	1.3	1.5
.5	.5	.6	.5	.6	.5	.6	.6	.6	.4	.6	.8
3.0	3.0	4.0	3.0	4.0	5.0	5.0	5.0	4.0	3.0	4.0	4.0
20	19	27	22	30	22	30	22	26	17	27	22
17	16	25	16	26	18	25	19	22	13	20	22
4	4	4	4	4	4	9	4	9	4	4	4
9	9	9	9	13	9	13	9	9	9	9	9
17	17	19	18	19	18	19	18	19	13	20	18
27	22	38	30	38	28	38	28	36	19	38	28
38	38	57	45	57	42	66	43	57	38	57	38
47	52	80	57	76	54	76	57	76	53	66	57
76	85	113	76	142	85	124	95	113	57	91	170
51	48	67	59	75	58	71	57	67	44	68	48
43	33	62	47	64	45	55	43	59	33	48	36
49	51	60	52	69	46	75	49	57	48	70	37
37	39	49	20	49	36	48	38	29	23	44	15
45	39	56	40	46	46	52	41	41	42	29	49
9	19	34	16	17	15	33	26	29	15	1	41
340	255	340	227	452	364	373	284	340	170	264	227
454	369	454	227	454	454	509	454	340	197	274	510
50	46	63	55	68	53	67	52	59	44	61	44
41	33	57	41	56	41	50	40	44	32	43	30
13	13	13	13	13	13	14	13	13	13	13	13
26	26	26	26	27	26	27	26	27	26	27	27
39	39	54	52	54	45	54	43	54	28	54	27
57	56	81	57	85	57	85	57	81	56	85	57
85	85	113	113	135	113	113	113	113	85	113	85
113	113	170	113	170	126	170	113	142	113	150	113
227	170	312	227	227	227	227	227	227	170	170	170
340	227	340	227	452	364	373	284	227	170	170	170
2.6	.9	0	0	3.7	1.4	13.1	4.9	0	0	.9	6.4
.6	.2	0	0	.4	.2	1.1	.5	0	0	.8	3.8

TABLE 7.27.--SAUSAGE OTHER THAN BOLOGNA, FRANKFURTERS, SALAMI, AND PORK BREAKFAST SAUSAGE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.2	.8	4.8	4.0	3.9
ON ONLY 1 OF 3 DAYS.....PCT.....	4.4	.8	4.2	3.6	3.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	0	.5	.3	.3
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	.1	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.0	1.2	1.1	1.2
STANDARD DEVIATION.....DO.....	.6	.0	.4	.4	.5
MAXIMUM.....DO.....	5.0	1.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	32	12	17	20	22
STANDARD DEVIATION.....GM.....	30	5	11	13	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		2	5	4
25TH.....GM.....	14		9	9	9
50TH.....GM.....	25		16	18	19
75TH.....GM.....	38		24	25	28
90TH.....GM.....	68		33	38	38
95TH.....GM.....	76		38	47	66
99TH.....GM.....	151		50	50	93
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	87	35	49	59	63
STANDARD DEVIATION.....GM.....	76	15	33	40	52
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	0	38	30	46
STANDARD DEVIATION.....GM.....	47	0	15	10	31
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	73	0	16	18	15
STANDARD DEVIATION.....GM.....	65	0	0	3	(2)
MAXIMUM IN ANY 1 DAY.....GM.....	907	57	151	150	280
MAXIMUM OVER 3 DAYS.....GM.....	907	57	151	150	280
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	79	35	44	52	55
STANDARD DEVIATION.....GM.....	70	15	31	38	49
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		8	14	11
25TH.....GM.....	28		19	25	22
50TH.....GM.....	57		34	35	34
75TH.....GM.....	113		64	71	71
90TH.....GM.....	168		85	114	113
95TH.....GM.....	227		113	140	140
99TH.....GM.....	340		151	150	280
MAXIMUM.....GM.....	907	57	151	150	280
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	60.0	0	1.0	6.6	5.1
PROPORTION OF EATING OCCASIONS.....PCT.....	2.5	0	1.7	8.5	6.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.5	3.9	4.6	4.0	5.9	4.6	7.7	5.6	6.0	3.8	3.5	3.4
4.6	3.5	3.7	3.6	5.0	4.3	6.3	4.8	5.2	3.0	2.5	2.9
.8	.3	.9	.3	.7	.2	1.0	.6	.6	.7	.9	.5
.1	.1	.1	0	.2	.1	.4	.2	.3	.1	.2	0
1.3	1.1	1.3	1.2	1.2	1.1	1.3	1.2	1.2	1.2	1.4	1.2
.6	.4	.6	.4	.6	.4	.7	.6	.5	.5	.6	.5
4.0	3.0	3.0	2.0	4.0	3.0	5.0	4.0	3.0	3.0	3.0	3.0
28	24	30	27	49	31	38	30	26	26	50	23
21	18	23	25	42	38	31	22	19	16	52	15
4	5	5	5	5	5	7	5	5	5		
11	9	17	9	19	14	19	15	11	16		
19	19	25	19	38	23	28	25	19	19		
38	31	38	32	67	38	56	38	38	38		
56	47	73	57	95	60	76	57	57	47		
75	75	85	101	140	76	101	76	66	57		
113	76	95	127	175	274	151	113	76	76		
75	67	79	74	134	78	100	84	66	72	137	68
60	50	59	65	125	59	81	58	46	49	165	49
70	69	68	82	104	62	89	54	90	49	64	37
31	38	45	65	50	59	54	30	20	15	23	7
15	28	57	0	91	204	71	59	34	57	150	0
0	0	0	0	44	107	34	41	11	0	0	0
340	227	284	340	907	454	454	340	170	227	454	168
340	227	284	380	907	822	454	425	227	227	454	168
67	64	71	69	119	84	87	74	66	62	108	58
53	46	52	57	107	76	68	54	43	42	127	45
11	11	17	11	15	14	16	11	17	16		
28	28	32	28	57	28	42	28	32	28		
57	56	57	57	85	57	64	57	56	57		
87	85	85	76	152	113	113	113	85	76		
142	113	142	114	227	180	170	152	114	113		
168	142	198	227	284	227	170	168	168	170		
227	227	284	304	454	454	340	227	170	227		
340	227	284	304	907	454	454	340	170	227	454	168
11.0	0	0	1.0	10.6	8.3	7.2	9.1	0	0	0	0
7.6	0	0	1.5	3.7	3.1	1.4	1.9	0	0	0	0

TABLE 7.28.--BEEF STEW WITH VEGETABLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.2	.7	5.1	4.6	5.5
ON ONLY 1 OF 3 DAYS.....PCT.....	4.7	.3	5.1	4.2	5.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	.2	0	.4	.4
ON ALL 3 OF 3 DAYS.....PCT.....	.1	.2	0	0	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.9	1.0	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.9	.1	.3	.3
MAXIMUM.....DO.....	4.0	3.0	2.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	133	109	60	75	90
STANDARD DEVIATION.....GM.....	92	80	45	43	49
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	41		20	20	41
25TH.....GM.....	82		41	41	50
50TH.....GM.....	123		41	82	82
75TH.....GM.....	163		82	82	102
90TH.....GM.....	245		82	163	163
95TH.....GM.....	327		163	163	204
99TH.....GM.....	490	225	327	190	245
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	357	122	181	210	247
STANDARD DEVIATION.....GM.....	199	0	136	115	125
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	382	122	0	191	247
STANDARD DEVIATION.....GM.....	200	0	0	85	66
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	451	225	0	0	223
STANDARD DEVIATION.....GM.....	221	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,960	245	980	613	735
MAXIMUM OVER 3 DAYS.....GM.....	2,940	674	980	613	735
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	355	173	178	207	247
STANDARD DEVIATION.....GM.....	189	55	112	111	119
PERCENTILES: <u>1/</u>					
5TH.....GM.....	123		61	61	123
25TH.....GM.....	245		123	123	151
50TH.....GM.....	306		123	245	245
75TH.....GM.....	490		245	245	245
90TH.....GM.....	490		245	326	368
95TH.....GM.....	735		490	490	490
99TH.....GM.....	981		490	490	613
MAXIMUM.....GM.....	1,470	245	490	613	735
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	28.0	0	0	0	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.3	0	0	0	1.4

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.0	6.7	5.5	5.4	4.9	4.4	5.7	5.0	6.9	6.0	6.5	4.0
4.7	6.4	5.0	5.1	4.5	4.2	5.0	4.7	5.8	4.9	5.4	2.9
.3	.3	.5	.2	.4	.3	.6	.3	.8	.9	.6	1.1
0	0	0	.1	(2)	0	.1	(2)	.3	.2	.5	.1
1.1	1.0	1.1	1.1	1.1	1.1	1.2	1.1	1.2	1.3	1.4	1.3
.2	.2	.3	.3	.4	.3	.4	.4	.5	.6	.7	.6
2.0	2.0	2.0	3.0	3.0	3.0	3.0	4.0	3.0	4.0	3.0	3.0
110	104	162	114	179	121	168	128	163	140	194	141
69	50	89	65	126	64	107	75	94	111	143	82
31	41	41	41	51	41	50	41	82	41		
82	82	82	82	82	82	82	82	102	82		
82	82	163	82	163	123	163	123	163	113		
123	123	204	163	245	163	163	163	163	163		
163	163	327	163	290	163	327	204	327	245		
245	163	327	245	368	245	327	245	408	327		
327	245	490	408	898	327	490	327	490	612		
312	297	454	325	471	342	444	355	397	324	443	320
176	135	252	177	243	162	233	157	127	171	311	103
338	299	398	339	580	353	450	321	434	361	532	314
172	93	100	119	366	153	134	94	158	204	91	103
0	0	0	245	490	0	466	523	450	490	490	490
0	0	0	0	0	0	312	346	41	141	0	0
980	735	1,470	1,225	1,960	980	1,470	1,470	735	980	1,470	490
1,225	980	1,470	1,225	2,695	1,470	2,940	2,695	1,470	1,838	1,470	1,470
314	297	440	315	471	336	439	344	405	327	426	316
179	132	234	163	219	158	216	151	137	168	248	114
113	123	123	123	123	123	151	123	245	93		
245	245	245	245	245	245	245	245	245	245		
245	245	490	245	490	245	490	368	490	245		
368	368	490	368	490	490	490	490	490	490		
490	490	736	490	736	490	735	490	490	551		
735	490	980	490	980	490	736	490	490	735		
736	735	1,470	1,224	980	736	1,225	736	735	735		
980	735	1,470	1,225	980	980	1,470	980	735	735	1,470	490
.8	.9	1.0	0	4.5	1.4	7.7	6.6	0	3.6	0	0
.7	.6	1.1	0	2.1	.6	2.4	1.7	0	2.6	0	0

TABLE 7.29.--CHILI CON CARNE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.5	.7	4.2	4.6	6.7
ON ONLY 1 OF 3 DAYS.....PCT.....	5.0	.7	3.6	4.3	6.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.5	.3	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.1	(2)	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.0	1.2	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.4	.4	.2
MAXIMUM.....DO.....	5.0	1.0	3.0	4.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	116	30	63	71	81
STANDARD DEVIATION.....GM.....	90	10	39	48	54
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11			5	10
25TH.....GM.....	64			43	43
50TH.....GM.....	85			64	85
75TH.....GM.....	170			85	85
90TH.....GM.....	213			170	170
95TH.....GM.....	255			170	170
99TH.....GM.....	510			170	340
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	308	91	156	197	228
STANDARD DEVIATION.....GM.....	200	31	82	132	135
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	362	0	189	223	303
STANDARD DEVIATION.....GM.....	222	0	38	32	134
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	426	0	175	4	0
STANDARD DEVIATION.....GM.....	228	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,530	128	383	510	510
MAXIMUM OVER 3 DAYS.....GM.....	2,550	128	526	510	1,020
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	308	91	164	187	232
STANDARD DEVIATION.....GM.....	198	31	82	130	139
PERCENTILES: <u>1/</u>					
5TH.....GM.....	32			5	24
25TH.....GM.....	171			84	128
50TH.....GM.....	255			191	255
75TH.....GM.....	383			255	255
90TH.....GM.....	510			383	510
95TH.....GM.....	765			510	510
99TH.....GM.....	1,020			510	510
MAXIMUM.....GM.....	1,530	128	383	510	510
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	42.0	0	1.0	0	1.0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.8	0	2.0	0	.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.0	6.4	7.4	5.2	6.6	5.5	6.3	5.3	5.8	1.8	4.1	1.3
6.7	5.9	7.0	4.9	5.8	5.0	5.6	4.9	5.1	1.8	4.1	1.1
.3	.5	.4	.2	.8	.4	.6	.4	.7	0	0	.2
0	0	0	.1	.1	.1	.1	(2)	0	0	0	0
1.0	1.1	1.1	1.1	1.2	1.1	1.2	1.1	1.1	1.0	1.0	1.4
.2	.3	.3	.5	.5	.4	.5	.4	.3	.2	.0	.5
2.0	3.0	3.0	5.0	4.0	3.0	4.0	4.0	2.0	2.0	1.0	2.0
106	99	129	92	156	112	140	110	123	94	98	126
72	60	98	64	133	94	103	65	78	61	48	67
11	16	11	5	11	11	21	11	11			
59	64	64	43	85	53	85	85	85			
85	85	106	85	128	85	128	85	85			
128	128	170	128	170	170	170	133	170			
213	170	255	170	301	170	255	170	213			
255	170	340	170	425	255	340	234	340			
340	298	510	340	596	510	552	340	340			
303	279	362	255	388	284	366	305	324	283	295	310
197	155	256	169	254	177	217	165	193	184	143	130
338	246	427	310	460	413	390	299	366	0	0	382
155	148	271	124	282	261	227	138	100	0	0	0
0	0	0	170	850	433	521	267	0	0	0	0
0	0	0	0	0	10	106	37	0	0	0	0
1,020	765	1,530	765	1,275	1,275	1,275	1,020	1,020	893	510	510
1,275	1,020	1,785	1,020	2,550	2,040	2,040	1,020	1,020	893	510	765
305	270	365	244	395	294	363	294	332	271	295	272
196	157	250	168	245	199	208	154	180	141	143	130
32	32	32	16	32	16	64	32	32			
158	128	191	128	255	158	255	224	255			
255	255	319	255	383	255	383	255	255			
383	383	510	319	510	383	510	383	383			
510	510	510	510	766	510	637	510	510			
765	510	766	510	766	510	765	510	637			
766	637	1,021	765	1,020	1,021	1,020	765	1,020			
1,020	765	1,530	765	1,020	1,275	1,020	1,020	1,020	510	510	510
1.9	1.0	1.3	2.7	8.6	4.7	9.8	5.9	0	4.2	0	0
1.2	.6	1.2	3.1	2.8	1.4	2.7	1.4	0	12.5	0	0

TABLE 7.30.--MEAT LOAF
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1:	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.8	2.7	9.3	8.5	10.6
ON ONLY 1 OF 3 DAYS.....PCT.....	7.8	2.2	8.7	7.8	10.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	.5	.5	.5	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	.2	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.2	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.4	.3	.4	.3
MAXIMUM.....DO.....	4.0	2.0	2.0	3.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	48	18	26	40	35
STANDARD DEVIATION.....GM.....	38	13	24	48	27
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		7	9	9
25TH.....GM.....	28		14	19	19
50TH.....GM.....	38		19	28	28
75TH.....GM.....	57		29	38	38
90TH.....GM.....	88		38	66	57
95TH.....GM.....	117		56	119	88
99TH.....GM.....	188		150	300	150
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	45	72	98	98
STANDARD DEVIATION.....GM.....	101	34	67	84	76
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	111	44	89	85	97
STANDARD DEVIATION.....GM.....	62	15	35	50	54
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	142	0	0	300	0
STANDARD DEVIATION.....GM.....	104	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,056	113	528	675	528
MAXIMUM OVER 3 DAYS.....GM.....	1,406	113	528	900	528
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	125	45	72	105	96
STANDARD DEVIATION.....GM.....	93	30	66	92	75
PERCENTILES: <u>1/</u>					
5TH.....GM.....	42		22	28	28
25TH.....GM.....	57		43	56	56
50TH.....GM.....	113		57	85	85
75TH.....GM.....	169		85	113	113
90TH.....GM.....	225		113	225	170
95TH.....GM.....	284		113	338	225
99TH.....GM.....	450		450	450	450
MAXIMUM.....GM.....	1,056	113	528	675	528
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	137.6	0	2.4	4.0	9.0
PROPORTION OF EATING OCCASIONS.....PCT.....	3.6	0	2.3	2.4	4.3

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
11.0	9.2	10.3	8.9	8.6	8.2	8.8	8.9	8.9	6.8	7.0	7.8
10.4	8.7	9.3	8.0	7.8	7.3	7.5	7.7	6.9	5.7	6.2	6.3
.5	.5	1.0	.8	.8	.8	1.2	1.2	1.8	1.0	.8	1.5
(2)	0	.1	.1	0	.1	.1	.1	.2	.1	0	0
1.1	1.1	1.1	1.1	1.2	1.2	1.2	1.2	1.3	1.2	1.1	1.3
.3	.3	.4	.4	.4	.4	.4	.4	.5	.5	.4	.5
3.0	2.0	3.0	3.0	3.0	4.0	3.0	3.0	3.0	3.0	2.0	3.0
44	41	59	48	63	47	59	47	54	40	49	45
33	34	37	31	44	31	38	39	33	25	49	46
14	14	19	14	15	14	17	14	19	7		7
27	19	38	28	38	28	29	28	29	19		21
38	29	47	38	57	38	56	38	44	38		28
57	44	75	66	76	57	76	57	71	57		57
76	75	113	76	113	85	112	76	113	76		103
85	110	150	101	138	113	132	110	132	76		150
175	225	176	176	227	149	191	192	150	113		235
127	118	164	139	180	127	162	129	133	102	147	113
99	97	101	93	117	84	106	102	70	61	150	128
86	105	135	91	135	113	124	98	126	115	80	115
30	76	61	37	108	46	62	47	58	23	66	71
131	0	189	57	0	120	132	137	132	57	0	0
0	0	0	0	0	23	27	156	0	0	0	0
792	792	528	528	1,056	450	675	1,056	352	281	675	704
792	792	567	528	1,276	450	732	1,406	450	338	675	704
120	113	157	125	162	121	145	119	127	98	129	106
93	81	93	86	115	75	98	96	60	51	123	86
43	43	43	43	43	43	43	43	43	28		22
71	57	112	82	86	74	85	66	85	57		57
86	86	129	113	114	113	113	88	113	113		85
149	113	176	169	225	169	176	132	170	113		113
225	225	264	225	264	225	264	225	224	170		220
227	330	340	227	352	227	338	227	225	170		352
524	450	495	528	676	397	495	454	225	281		450
792	450	528	528	1,056	450	675	1,056	352	281	675	450
13.8	9.8	1.2	7.8	13.4	19.5	19.8	25.9	3.7	7.3	0	0
5.5	4.5	.7	5.2	3.4	3.9	3.8	3.4	2.9	5.0	0	0

TABLE 7.31.--TUNA SALAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.4	.4	2.6	4.1	3.7
ON ONLY 1 OF 3 DAYS.....PCT.....	3.9	.4	2.4	3.6	3.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.2	.5	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.0	1.1	1.2	1.2
STANDARD DEVIATION.....DO.....	.4	.0	.4	.4	.5
MAXIMUM.....DO.....	4.0	1.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	39	11	20	26	27
STANDARD DEVIATION.....GM.....	30	6	18	20	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9			4	8
25TH.....GM.....	17			9	17
50TH.....GM.....	34			17	26
75TH.....GM.....	51			34	34
90TH.....GM.....	68			51	51
95TH.....GM.....	103			68	68
99TH.....GM.....	137			85	85
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	106	33	56	75	78
STANDARD DEVIATION.....GM.....	82	19	53	60	47
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	87	0	62	54	55
STANDARD DEVIATION.....GM.....	53	0	13	25	27
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	79	0	20	18	41
STANDARD DEVIATION.....GM.....	45	0	0	0	30
MAXIMUM IN ANY 1 DAY.....GM.....	666	51	205	256	205
MAXIMUM OVER 3 DAYS.....GM.....	666	51	205	256	256
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	99	33	54	68	68
STANDARD DEVIATION.....GM.....	75	19	49	56	42
PERCENTILES: <u>1/</u>					
5TH.....GM.....	25			12	13
25TH.....GM.....	51			26	51
50TH.....GM.....	85			51	51
75TH.....GM.....	103			103	103
90TH.....GM.....	205			154	103
95TH.....GM.....	205			205	154
99TH.....GM.....	410			256	205
MAXIMUM.....GM.....	666	51	205	256	205
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	22.2	0	1.5	0	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.1	0	4.7	0	1.8

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.8	4.0	4.2	5.3	4.4	5.5	3.6	6.0	3.4	4.2	1.9	1.8
3.2	3.6	3.9	4.6	3.9	4.7	3.2	5.3	2.9	3.6	1.9	1.8
.5	.4	.2	.5	.4	.6	.4	.5	.5	.6	0	0
.1	0	.2	.2	.1	.2	0	.2	0	.1	0	0
1.2	1.1	1.2	1.2	1.2	1.2	1.1	1.2	1.1	1.2	1.0	1.0
.5	.3	.5	.5	.4	.5	.3	.5	.4	.4	.0	.0
3.0	2.0	3.0	4.0	3.0	3.0	2.0	3.0	2.0	3.0	1.0	1.0
42	35	43	38	40	37	48	40	34	39	41	27
33	27	37	34	26	28	39	31	19	27	36	21
9	9	9	9	13	9	13	9		8		
17	17	17	17	23	17	23	17		17		
28	34	34	28	34	34	34	34		34		
68	42	68	51	51	51	61	51		51		
103	68	92	68	68	68	137	71		91		
111	85	137	103	91	92	137	111		103		
137	137	171	137	137	137	137	137		103		
109	93	121	98	112	102	135	110	91	110	124	80
87	67	101	76	72	78	109	83	52	76	109	63
107	98	61	90	92	78	116	89	82	76	0	0
57	59	57	63	47	45	71	51	25	39	0	0
64	0	103	128	85	60	0	101	0	102	0	0
38	0	57	60	0	40	0	25	0	0	0	0
410	308	513	410	410	410	666	410	205	308	308	205
410	410	513	615	410	410	666	410	205	308	308	205
103	93	111	97	102	93	130	104	89	102	124	80
77	65	92	74	65	71	101	73	47	70	109	63
26	26	26	25	26	26	26	25		25		
51	51	51	51	51	51	57	51		51		
85	103	85	78	103	68	103	103		103		
103	103	137	103	103	103	154	115		113		
205	205	205	205	205	205	205	205		205		
205	205	308	205	205	205	410	205		284		
410	308	512	410	308	410	410	410		308		
410	308	513	410	410	410	666	410	205	308	308	205
0	0	1.3	1.9	0	2.8	5.0	6.8	1.5	0	0	0
0	0	1.9	2.0	0	.8	2.5	1.4	3.6	0	0	0

TABLE 7.32.--POT PIES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.1	.4	2.3	2.0	2.2
ON ONLY 1 OF 3 DAYS.....PCT.....	2.0	.4	2.1	1.9	2.2
ON ONLY 2 OF 3 DAYS.....PCT.....	.1	0	.1	.1	(2)
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.2	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	1.0	1.0	1.3	1.1	1.0
STANDARD DEVIATION.....DO.....	.2	.0	.7	.2	.1
MAXIMUM.....DO.....	4.0	1.0	4.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	91	38	75	68	67
STANDARD DEVIATION.....GM.....	49	22	57	31	23
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	38				
25TH.....GM.....	76				
50TH.....GM.....	76				
75TH.....GM.....	95				
90TH.....GM.....	160				
95TH.....GM.....	167				
99TH.....GM.....	250				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	260	115	178	186	197
STANDARD DEVIATION.....GM.....	126	65	96	64	59
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	296	0	227	234	227
STANDARD DEVIATION.....GM.....	110	0	0	6	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	232	0	232	0	0
STANDARD DEVIATION.....GM.....	42	0	42	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	960	188	454	360	360
MAXIMUM OVER 3 DAYS.....GM.....	1,200	188	850	480	454
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	261	115	180	192	198
STANDARD DEVIATION.....GM.....	127	65	79	62	58
PERCENTILES: ^{1/}					
5TH.....GM.....	113				
25TH.....GM.....	227				
50TH.....GM.....	227				
75TH.....GM.....	250				
90TH.....GM.....	454				
95TH.....GM.....	480				
99TH.....GM.....	750				
MAXIMUM.....GM.....	960	188	454	360	360
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	5.6	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	0	0

^{1/} SEE TABLE NOTES IN APPENDIX.

^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.1	2.7	2.4	1.8	2.6	2.1	1.6	1.7	2.2	3.0	2.0	1.6
2.9	2.6	2.2	1.8	2.5	2.1	1.5	1.7	2.0	2.9	1.7	1.6
.2	.1	.2	.1	.1	(2)	.1	(2)	.2	(2)	.2	0
0	0	0	0	0	0	0	0	0	0	0	0
1.1	1.0	1.1	1.0	1.0	1.0	1.1	1.0	1.1	1.0	1.1	1.0
.3	.2	.3	.2	.2	.1	.2	.2	.3	.1	.3	.0
2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	1.0
90	85	119	81	107	88	114	86	94	75	116	88
39	40	49	31	53	34	82	43	37	30	61	28
40	38			47	53	38	33		38		
76	72			76	76	76	76		76		
76	76			76	76	76	76		76		
95	83			151	83	151	83		76		
151	151			167	151	200	151		100		
160	160			227	160	303	160		100		
240	240			240	160	400	240		250		
259	240	346	231	312	259	303	251	266	217	334	263
110	97	148	81	150	100	177	120	104	60	191	83
234	288	233	240	340	227	544	288	227	375	227	0
6	66	6	0	0	0	71	47	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0
720	625	720	500	750	750	960	720	480	454	750	500
720	720	720	500	750	750	1,200	720	480	750	750	500
252	244	327	231	310	258	326	253	260	222	311	263
100	98	142	77	147	98	204	117	96	66	175	83
120	114			142	158	113	99		114		
227	227			227	227	227	227		227		
227	227			227	227	227	227		227		
240	240			454	250	454	284		227		
360	375			480	454	500	375		300		
454	480			568	480	901	480		300		
709	625			721	480	960	720		454		
720	625	720	500	750	750	960	720	480	454	750	500
0	1.0	0	.9	0	0	1.2	.8	0	1.7	0	0
0	1.6	0	3.2	0	0	1.5	.6	0	3.2	0	0

TABLE 7.33.--FROZEN PLATE MEALS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.4	.4	1.6	1.2	1.0
ON ONLY 1 OF 3 DAYS.....PCT.....	1.2	.4	1.6	1.0	.9
ON ONLY 2 OF 3 DAYS.....PCT.....	.1	0	.1	.1	0
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	.1	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.1	1.0	1.0	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.2	.5	.4
MAXIMUM.....DO.....	3.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	122	19	96	117	111
STANDARD DEVIATION.....GM.....	57	(2)	34	72	52
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	57				
25TH.....GM.....	93				
50TH.....GM.....	104				
75TH.....GM.....	142				
90TH.....GM.....	196				
95TH.....GM.....	238				
99TH.....GM.....	335				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	324	56	271	295	308
STANDARD DEVIATION.....GM.....	110	(2)	70	67	105
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	307	0	312	142	0
STANDARD DEVIATION.....GM.....	60	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	300	0	0	369	301
STANDARD DEVIATION.....GM.....	50	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	907	57	340	539	510
MAXIMUM OVER 3 DAYS.....GM.....	1,106	57	624	1,106	903
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	319	56	274	296	307
STANDARD DEVIATION.....GM.....	103	(2)	68	80	99
PERCENTILES: <u>1/</u>					
5TH.....GM.....	142				
25TH.....GM.....	279				
50TH.....GM.....	312				
75TH.....GM.....	340				
90TH.....GM.....	454				
95TH.....GM.....	510				
99TH.....GM.....	680				
MAXIMUM.....GM.....	907	57	340	539	510
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	2.7	0	0	0	.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	0	0	3.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.3	1.1	1.1	1.5	1.5	1.5	1.2	1.4	2.5	1.7	1.5	2.1
1.3	1.0	.9	1.4	1.3	1.3	1.0	1.3	2.0	1.7	1.4	1.7
(2)	.1	.2	.2	.1	.2	.1	.1	.3	0	0	0
0	0	0	0	.1	0	.1	(2)	.3	0	.1	.4
1.0	1.1	1.1	1.1	1.2	1.1	1.2	1.1	1.3	1.0	1.2	1.4
.2	.3	.3	.3	.5	.3	.6	.4	.7	.0	.6	.8
2.0	2.0	2.0	2.0	3.0	2.0	3.0	3.0	3.0	1.0	3.0	3.0
106	102	134	107	134	119	151	121	135	105	110	133
37	47	71	38	64	46	72	50	60	16	46	82
				80	57	78	47				
				93	95	102	95				
				104	104	113	109				
				151	149	196	149				
				217	186	265	180				
				301	210	302	217				
				340	248	343	333				
308	280	340	298	340	321	386	339	328	315	286	282
97	108	146	97	83	97	163	115	92	49	8	56
312	276	399	248	343	309	325	300	287	0	0	0
0	88	60	42	21	43	46	40	8	0	0	0
0	0	0	0	328	0	318	256	266	0	256	299
0	0	0	0	16	0	19	83	45	0	0	16
539	539	680	482	539	539	907	680	510	397	312	340
624	680	922	567	1,021	744	1,028	999	1,020	397	769	936
309	279	354	288	338	318	364	330	306	315	279	289
94	105	144	91	72	91	138	113	100	49	31	45
				241	142	248	113				
				312	284	284	280				
				312	312	312	312				
				347	340	397	369				
				454	447	532	454				
				482	538	680	540				
				538	539	907	680				
539	539	680	482	539	539	907	680	510	397	312	340
0	.7	0	0	0	0	1.3	0	0	0	0	0
0	2.6	0	0	0	0	1.9	0	0	0	0	0

TABLE 7.34.--HAMBURGER AND CHEESEBURGER SANDWICHES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.0	0	7.1	9.4	10.8
ON ONLY 1 OF 3 DAYS.....PCT.....	8.5	0	6.3	8.5	9.8
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	0	.7	.8	.9
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	0	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	0	.4	.3	.4
MAXIMUM.....DO.....	6.0	0	3.0	2.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	65	0	38	43	51
STANDARD DEVIATION.....GM.....	44	0	23	27	36
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	32		9	16	28
25TH.....GM.....	37		19	32	32
50TH.....GM.....	52		32	32	37
75TH.....GM.....	69		51	52	56
90TH.....GM.....	113		65	65	77
95TH.....GM.....	154		95	103	113
99TH.....GM.....	244		120	162	248
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	160	0	98	118	133
STANDARD DEVIATION.....GM.....	75	0	53	64	60
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	174	0	120	130	157
STANDARD DEVIATION.....GM.....	74	0	36	57	73
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	195	0	0	0	310
STANDARD DEVIATION.....GM.....	77	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	770	0	336	485	558
MAXIMUM OVER 3 DAYS.....GM.....	1,386	0	359	558	930
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	161	0	99	116	138
STANDARD DEVIATION.....GM.....	74	0	49	58	65
PERCENTILES: <u>1/</u>					
5TH.....GM.....	97		42	49	84
25TH.....GM.....	97		73	97	97
50TH.....GM.....	157		97	97	111
75TH.....GM.....	186		111	113	168
90TH.....GM.....	266		157	170	211
95TH.....GM.....	308		186	222	266
99TH.....GM.....	444		336	336	372
MAXIMUM.....GM.....	616	0	336	485	558
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	2.3	0	1.0	1.2	0
PROPORTION OF EATING OCCASIONS.....PCT.....	(2)	0	1.3	.7	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
13.1	12.9	19.0	15.1	18.0	11.6	10.4	5.2	3.6	3.2	2.1	1.2
11.7	12.1	14.3	12.7	13.6	9.6	8.9	4.8	3.2	3.1	2.1	1.2
1.2	.6	3.7	1.6	3.4	1.9	1.0	.3	.4	(2)	0	0
.2	.2	1.0	.7	1.0	.1	.5	.1	0	0	0	0
1.1	1.1	1.4	1.3	1.3	1.2	1.2	1.1	1.2	1.0	1.0	1.0
.4	.4	.7	.7	.6	.4	.5	.4	.4	.1	.0	.0
4.0	4.0	5.0	6.0	4.0	3.0	5.0	3.0	2.0	2.0	1.0	1.0
58	50	82	60	84	65	71	53	54	46	53	45
36	24	59	40	56	35	43	31	22	18	12	10
32	30	32	32	32	32	32	28		28		
32	32	51	32	52	37	51	32		32		
52	38	61	52	62	56	61	51		32		
65	61	99	65	105	69	89	61		61		
105	76	157	97	154	111	124	89		74		
124	89	198	126	210	150	177	103		89		
174	154	333	207	266	186	238	177		89		
155	138	180	147	188	165	181	143	142	137	160	136
73	50	86	62	89	64	88	55	41	53	35	31
169	136	195	162	199	161	158	178	152	111	0	0
94	51	87	65	70	60	63	94	20	0	0	0
143	134	218	146	220	191	189	127	0	0	0	0
21	45	96	83	75	52	55	28	0	0	0	0
732	340	616	532	770	497	532	454	266	266	200	186
1,098	652	1,386	1,029	1,330	798	873	871	340	266	200	186
154	136	182	143	191	162	175	145	138	136	160	136
73	49	90	53	80	62	82	61	38	53	35	31
97	85	97	97	97	97	97	85		97		
97	97	111	97	154	111	111	97		97		
154	111	157	142	183	157	157	154		97		
183	157	194	183	222	186	186	170		183		
266	194	308	194	266	266	266	222		186		
314	227	366	266	340	266	366	266		266		
399	266	532	308	532	336	532	340		266		
523	340	608	334	616	485	532	454	266	266	200	186
0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0

TABLE 7.35.--SOUPS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	31.3	12.4	47.0	39.7	31.2
ON ONLY 1 OF 3 DAYS.....PCT.....	22.1	6.5	29.6	27.5	24.3
ON ONLY 2 OF 3 DAYS.....PCT.....	7.1	4.8	13.1	9.6	5.5
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	1.1	4.3	2.7	1.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.7	1.5	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.8	.8	.7	.6
MAXIMUM.....DO.....	10.0	4.0	7.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	139	80	96	101	110
STANDARD DEVIATION.....GM.....	100	69	80	66	70
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	40	10	20	31	31
25TH.....GM.....	80	20	41	60	76
50TH.....GM.....	107	63	80	82	82
75TH.....GM.....	163	122	121	123	160
90TH.....GM.....	245	160	163	189	204
95TH.....GM.....	327	188	242	242	243
99TH.....GM.....	509	395	327	320	345
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	306	113	188	224	259
STANDARD DEVIATION.....GM.....	178	90	102	117	136
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	307	173	196	219	274
STANDARD DEVIATION.....GM.....	150	80	110	89	121
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	310	177	216	215	224
STANDARD DEVIATION.....GM.....	145	105	142	95	68
MAXIMUM IN ANY 1 DAY.....GM.....	2,450	735	1,800	827	1,080
MAXIMUM OVER 3 DAYS.....GM.....	4,595	1,185	3,180	2,040	1,455
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	290	140	186	211	249
STANDARD DEVIATION.....GM.....	155	97	103	100	125
PERCENTILES: <u>1/</u>					
5TH.....GM.....	114	40	60	81	92
25TH.....GM.....	240	61	120	123	180
50TH.....GM.....	245	123	180	240	240
75TH.....GM.....	360	188	245	245	245
90TH.....GM.....	490	245	245	250	480
95TH.....GM.....	600	250	368	375	490
99TH.....GM.....	736	368	601	601	721
MAXIMUM.....GM.....	1,960	735	735	735	858
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	191.4	0	8.8	6.9	2.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.1	0	1.2	.7	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
27.8	29.4	24.1	24.0	27.0	29.9	32.2	31.4	37.5	36.9	34.6	42.1
22.0	22.7	18.3	19.0	20.8	22.5	21.6	21.4	23.6	23.7	19.7	25.7
4.9	5.5	4.5	3.9	5.2	5.8	7.8	7.8	10.4	9.4	9.9	11.7
.9	1.3	.8	1.2	1.0	1.6	2.8	2.1	3.6	3.8	5.0	4.8
1.3	1.3	1.3	1.3	1.3	1.4	1.5	1.5	1.5	1.6	1.7	1.6
.6	.6	.6	.6	.6	.7	.8	.8	.8	.8	1.0	.8
10.0	5.0	4.0	5.0	4.0	6.0	6.0	9.0	5.0	5.0	6.0	5.0
126	124	156	127	152	131	166	141	172	150	194	145
99	79	120	85	104	96	112	95	117	98	162	97
41	41	40	27	41	40	50	40	60	41	41	41
80	80	80	80	82	80	82	80	82	82	82	80
82	90	120	83	123	85	151	120	160	123	160	123
162	160	179	163	203	163	208	163	213	194	242	163
240	204	323	240	280	243	323	245	327	250	395	250
250	245	404	306	327	323	400	323	405	322	512	320
483	485	613	487	571	551	561	492	605	478	817	490
298	297	336	310	357	289	361	306	359	316	355	286
148	150	181	183	207	156	215	159	229	164	305	133
316	295	405	276	359	311	355	295	356	312	353	288
160	143	199	115	149	150	168	130	148	149	151	126
305	260	442	329	384	337	329	323	326	288	411	301
293	96	180	94	151	138	117	134	150	107	222	109
1,940	1,470	1,470	1,225	2,000	1,470	1,960	1,470	1,816	1,715	2,450	980
4,595	1,573	2,373	1,508	2,002	1,960	2,573	2,293	2,695	2,817	3,341	2,200
294	282	358	290	347	282	334	287	333	284	334	270
158	131	198	153	181	139	183	142	160	129	175	109
120	120	120	61	120	91	123	120	123	120	123	120
240	240	240	240	240	240	240	240	240	240	240	240
245	245	305	245	300	245	245	245	250	245	245	245
360	360	480	368	485	360	480	360	454	360	454	300
490	490	612	490	612	490	500	490	500	490	562	480
500	490	720	540	720	500	659	490	613	495	626	490
751	735	1,001	740	980	731	981	736	930	637	961	613
1,470	735	1,470	980	1,225	1,200	1,960	1,470	980	1,102	1,470	735
8.3	6.8	10.0	9.0	19.9	21.5	28.3	40.3	4.0	12.8	5.1	7.0
1.1	.8	2.3	1.9	1.4	1.0	1.2	1.2	.6	1.3	1.6	1.0

TABLE 8.01.--FLUID MILK, MILK BEVERAGES, AND MILK-BASED INFANT FORMULAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	82.5	89.0	96.9	97.0	98.5
ON ONLY 1 OF 3 DAYS.....PCT.....	11.5	1.8	4.1	4.2	3.8
ON ONLY 2 OF 3 DAYS.....PCT.....	15.8	3.0	9.6	13.3	9.9
ON ALL 3 OF 3 DAYS.....PCT.....	55.2	84.3	83.2	79.5	84.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	4.9	11.8	7.1	5.9	6.2
STANDARD DEVIATION.....DO.....	3.1	4.8	3.5	2.7	2.7
MAXIMUM.....DO.....	39.0	34.0	24.0	17.0	17.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	328	667	424	392	468
STANDARD DEVIATION.....GM.....	270	308	266	215	240
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	31	117	73	83	122
25TH.....GM.....	122	488	240	225	286
50TH.....GM.....	248	683	368	366	447
75TH.....GM.....	470	857	572	528	611
90TH.....GM.....	692	1,030	773	691	793
95TH.....GM.....	833	1,140	904	793	899
99TH.....GM.....	1,221	1,443	1,189	977	1,139
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	246	195	181	269	280
STANDARD DEVIATION.....GM.....	171	155	183	204	186
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	289	331	275	286	333
STANDARD DEVIATION.....GM.....	189	329	161	149	161
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	418	695	470	441	514
STANDARD DEVIATION.....GM.....	279	289	256	201	223
MAXIMUM IN ANY 1 DAY.....GM.....	4,410	2,083	1,960	1,960	2,318
MAXIMUM OVER 3 DAYS.....GM.....	10,053	5,856	5,856	4,087	4,911
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	203	170	179	198	227
STANDARD DEVIATION.....GM.....	134	71	80	83	89
PERCENTILES: ^{1/}					
5TH.....GM.....	15	61	61	81	122
25TH.....GM.....	122	123	122	123	183
50TH.....GM.....	244	171	183	184	244
75TH.....GM.....	245	244	244	244	244
90TH.....GM.....	366	248	245	245	281
95TH.....GM.....	488	248	245	306	375
99TH.....GM.....	552	310	488	488	490
MAXIMUM.....GM.....	1,960	496	976	1,069	1,220
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	466.1	4.5	21.8	31.2	40.9
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	.1	.3	.3	.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
97.4	95.1	93.2	88.0	78.8	77.4	75.2	72.4	81.5	79.9	83.1	83.6
5.5	5.6	8.1	12.1	14.5	16.6	13.7	15.0	10.3	10.9	11.2	9.0
12.3	15.7	15.1	20.1	18.6	19.4	16.6	16.3	14.0	15.0	13.5	13.3
79.6	73.9	70.1	55.7	45.7	41.3	44.9	41.1	57.2	54.1	58.4	61.3
5.9	5.4	5.3	4.4	4.2	3.9	4.4	4.1	4.8	4.3	4.8	4.5
2.7	2.5	2.7	2.5	2.8	2.8	3.1	2.9	3.3	2.7	3.0	2.7
15.0	15.0	23.0	17.0	26.0	29.0	21.0	26.0	29.0	23.0	14.0	39.0
517	433	558	359	355	242	258	195	263	207	269	237
301	238	375	246	306	205	243	174	225	166	249	180
122	82	81	81	30	20	18	15	20	20	29	21
305	245	287	165	126	82	82	81	92	81	95	107
488	407	488	325	259	184	183	158	224	163	183	204
672	573	737	488	488	327	351	264	359	285	376	326
895	734	1,051	693	772	500	565	429	553	434	576	474
1,079	857	1,224	832	955	650	732	531	733	542	755	570
1,466	1,140	1,879	1,151	1,423	952	1,189	746	977	732	1,222	819
347	287	343	303	298	234	243	210	228	201	209	186
214	183	220	188	196	147	173	151	133	127	108	137
408	349	470	335	347	269	269	218	230	226	251	255
205	180	304	158	232	158	179	143	153	125	195	136
583	501	661	464	486	337	341	260	324	250	331	277
288	219	363	242	327	225	272	196	237	179	265	187
3,172	2,982	4,157	2,187	4,410	2,235	2,928	2,713	2,806	1,490	2,196	1,496
6,710	5,420	10,053	5,359	7,381	4,715	6,019	6,159	7,015	3,484	4,392	4,452
265	242	314	244	254	185	178	142	165	143	167	157
125	103	164	113	181	131	153	120	127	111	125	109
122	122	122	61	15	11	10	5	5	5	5	10
244	206	244	244	122	61	31	31	31	31	61	61
244	244	245	244	244	244	183	122	176	123	183	123
250	245	368	245	366	245	245	244	244	244	244	244
488	366	490	366	488	306	366	245	305	245	305	245
488	488	500	488	490	469	488	306	368	305	366	305
733	500	977	521	976	490	611	490	488	488	490	488
1,464	1,776	1,960	980	1,960	1,500	1,464	980	1,000	976	976	624
46.7	27.7	41.4	23.9	61.5	32.6	51.3	43.0	7.3	16.5	3.9	11.8
.4	.3	.6	.4	.5	.2	.3	.2	.2	.3	.2	.3

TABLE 8.02.--FLUID MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	78.9	50.8	95.3	95.2	96.9
ON ONLY 1 OF 3 DAYS.....PCT.....	12.4	3.3	4.7	5.7	5.3
ON ONLY 2 OF 3 DAYS.....PCT.....	16.1	2.8	11.3	14.0	12.2
ON ALL 3 OF 3 DAYS.....PCT.....	50.3	44.7	79.4	75.4	79.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	4.6	9.5	6.9	5.7	5.8
STANDARD DEVIATION.....DO.....	3.0	4.5	3.5	2.7	2.7
MAXIMUM.....DO.....	36.0	26.0	24.0	17.0	16.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	309	558	409	372	434
STANDARD DEVIATION.....GM.....	259	338	265	212	236
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	31	61	61	81	82
25TH.....GM.....	122	306	214	204	245
50TH.....GM.....	244	539	366	346	407
75TH.....GM.....	434	777	561	498	570
90TH.....GM.....	651	976	763	651	733
95TH.....GM.....	794	1,061	885	753	876
99TH.....GM.....	1,160	1,466	1,184	940	1,058
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	243	247	175	271	284
STANDARD DEVIATION.....GM.....	172	280	174	201	181
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	284	219	269	277	323
STANDARD DEVIATION.....GM.....	186	231	157	149	155
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	403	618	463	428	490
STANDARD DEVIATION.....GM.....	270	311	254	197	219
MAXIMUM IN ANY 1 DAY.....GM.....	4,410	2,083	1,960	1,960	2,318
MAXIMUM OVER 3 DAYS.....GM.....	9,053	5,856	5,856	4,087	4,911
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	201	176	179	195	224
STANDARD DEVIATION.....GM.....	133	78	79	82	89
PERCENTILES: <u>1/</u>					
5TH.....GM.....	15	61	61	61	122
25TH.....GM.....	122	122	122	122	183
50TH.....GM.....	244	183	183	184	244
75TH.....GM.....	244	244	244	244	244
90TH.....GM.....	366	245	245	245	245
95TH.....GM.....	488	245	245	305	368
99TH.....GM.....	550	337	429	488	488
MAXIMUM.....GM.....	1,960	488	976	1,069	1,220
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	406.1	3.4	20.9	27.9	39.3
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	.1	.3	.3	.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
96.2	93.0	91.0	85.2	76.6	74.1	71.5	68.5	76.6	74.1	77.5	79.0
7.6	7.6	9.3	14.7	15.2	16.5	14.2	15.7	11.1	12.0	11.1	10.0
14.7	18.1	15.0	20.5	18.6	19.1	16.3	15.7	14.7	14.7	14.1	14.0
73.9	67.4	66.7	50.0	42.7	38.4	40.9	37.1	50.8	47.3	52.3	55.1
5.5	5.0	5.1	4.2	4.0	3.8	4.2	4.0	4.4	4.1	4.6	4.3
2.6	2.5	2.7	2.5	2.8	2.7	3.0	2.9	2.8	2.6	2.9	2.6
14.0	14.0	23.0	17.0	26.0	29.0	21.0	26.0	17.0	23.0	14.0	36.0
472	396	523	333	340	231	249	186	250	198	262	223
293	232	365	243	301	199	235	171	213	159	246	164
81	81	81	51	29	20	20	15	25	20	24	20
245	224	245	163	122	81	82	66	86	81	92	92
427	366	449	285	244	166	171	142	203	163	166	183
630	530	711	447	488	325	327	245	336	265	369	325
854	711	978	651	735	489	534	407	510	417	570	452
1,017	814	1,198	814	916	592	693	528	691	528	753	564
1,423	1,098	1,872	1,139	1,423	976	1,060	740	936	732	1,222	733
329	284	338	288	292	230	250	203	234	188	197	169
206	173	233	185	189	148	182	144	153	123	108	125
390	341	459	321	342	263	269	212	232	220	260	231
199	173	301	155	226	156	182	140	163	123	193	131
551	476	629	452	475	325	335	254	315	249	327	271
281	215	356	242	326	221	265	194	227	169	265	167
3,172	1,952	4,157	1,960	4,410	1,952	2,928	2,713	2,806	1,220	2,196	996
6,710	4,880	9,053	5,124	7,381	4,404	6,019	6,159	7,015	2,724	4,392	2,952
260	238	310	241	252	181	177	140	170	144	172	154
122	102	161	114	180	130	151	118	125	110	123	106
122	122	122	61	15	10	10	5	10	5	10	10
244	183	244	244	122	61	31	31	61	31	61	61
244	244	244	244	244	244	183	122	183	123	183	123
245	244	368	245	366	244	245	244	244	244	244	244
488	366	488	366	488	306	366	245	305	245	305	245
488	488	490	488	490	427	488	305	366	305	366	275
732	490	976	549	976	490	610	490	488	488	488	488
1,464	1,776	1,960	980	1,960	1,220	1,464	980	976	976	976	490
38.7	26.9	36.5	20.7	53.7	26.2	43.5	35.2	5.3	12.1	3.9	11.8
.4	.3	.6	.4	.4	.2	.3	.2	.1	.2	.2	.3

TABLE 8.03.--FLUID WHOLE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	64.6	37.4	76.0	79.1	86.3
ON ONLY 1 OF 3 DAYS.....PCT.....	13.6	3.1	9.6	9.5	9.2
ON ONLY 2 OF 3 DAYS.....PCT.....	13.9	3.2	10.1	13.2	13.9
ON ALL 3 OF 3 DAYS.....PCT.....	37.1	31.1	56.3	56.4	63.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	4.3	9.1	6.2	5.2	5.2
STANDARD DEVIATION.....DO.....	3.0	4.7	3.7	2.8	2.8
MAXIMUM.....DO.....	36.0	24.0	24.0	16.0	16.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	282	542	375	341	391
STANDARD DEVIATION.....GM.....	250	350	266	213	239
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	20	51	41	44	81
25TH.....GM.....	81	244	164	163	203
50TH.....GM.....	214	529	325	316	366
75TH.....GM.....	407	777	529	488	529
90TH.....GM.....	611	956	733	650	732
95TH.....GM.....	752	1,138	864	733	854
99TH.....GM.....	1,100	1,465	1,058	915	1,056
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	243	225	211	288	289
STANDARD DEVIATION.....GM.....	187	214	184	241	198
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	284	232	290	272	318
STANDARD DEVIATION.....GM.....	196	217	169	147	156
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	390	629	460	421	473
STANDARD DEVIATION.....GM.....	267	316	252	195	220
MAXIMUM IN ANY 1 DAY.....GM.....	3,294	1,952	1,952	1,830	2,318
MAXIMUM OVER 3 DAYS.....GM.....	7,930	5,856	5,856	4,087	4,911
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	198	178	180	197	226
STANDARD DEVIATION.....GM.....	135	79	80	83	90
PERCENTILES: ^{1/}					
5TH.....GM.....	15	46	61	61	122
25TH.....GM.....	122	122	122	122	183
50TH.....GM.....	244	183	183	183	244
75TH.....GM.....	244	244	244	244	244
90TH.....GM.....	366	244	244	244	244
95TH.....GM.....	488	244	244	305	427
99TH.....GM.....	529	366	488	488	488
MAXIMUM.....GM.....	1,952	488	976	1,069	1,220
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	313.9	2.6	17.9	21.6	32.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	.2	.4	.3	.4

^{1/} SEE TABLE NOTES IN APPENDIX.

^{2/} MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
87.9	82.7	80.1	71.7	65.0	57.8	57.7	51.9	59.2	54.6	66.9	63.1
10.7	10.9	12.7	16.7	16.9	16.1	14.3	15.3	12.0	11.7	10.7	10.4
17.0	18.1	16.8	16.9	16.5	15.0	13.4	11.8	10.6	10.7	12.9	12.3
60.1	53.7	50.6	38.1	31.6	26.7	30.0	24.8	36.7	32.3	43.3	40.3
4.8	4.5	4.5	3.8	3.7	3.7	4.0	3.7	4.3	3.9	4.6	4.2
2.7	2.5	2.7	2.5	2.8	2.8	3.1	2.9	2.9	2.7	3.0	2.8
14.0	14.0	23.0	14.0	26.0	29.0	21.0	26.0	17.0	23.0	14.0	36.0
416	355	450	303	305	208	221	162	235	173	257	212
287	228	340	238	288	192	215	162	225	153	254	169
81	81	81	41	20	10	12	10	15	12	20	20
203	163	203	122	97	81	81	46	81	70	81	81
356	325	402	244	224	163	163	117	163	122	168	164
569	488	611	407	407	285	300	225	325	244	325	305
773	651	855	611	671	454	488	366	528	407	570	452
936	773	1,057	744	874	570	650	488	692	488	772	569
1,343	1,017	1,709	1,138	1,423	976	976	712	977	718	1,464	733
322	290	354	272	288	228	251	190	204	176	187	159
189	183	229	173	209	157	217	146	148	125	111	122
363	329	442	311	344	251	263	203	218	219	269	222
193	169	336	157	230	158	196	151	179	128	201	132
521	454	584	438	455	311	306	235	315	224	329	273
280	214	333	242	327	224	249	192	243	171	278	174
3,172	1,952	3,294	1,952	3,172	1,952	2,928	2,713	2,806	1,220	2,196	996
6,710	4,880	7,930	5,124	7,381	4,209	6,019	6,159	7,015	2,684	4,392	2,952
258	238	303	239	244	171	166	129	165	133	169	153
120	100	155	115	181	130	152	119	131	114	128	109
122	122	122	31	15	10	8	5	8	5	5	10
244	183	244	198	122	31	31	29	31	31	61	61
244	244	244	244	244	183	122	93	153	122	153	122
244	244	366	244	305	244	244	244	244	244	244	244
488	366	488	366	488	305	366	244	305	244	305	244
488	488	488	488	488	366	488	305	366	305	366	305
732	488	976	549	976	488	610	488	488	488	488	488
1,464	1,769	1,464	976	1,952	1,220	1,464	976	976	976	976	488
34.0	22.6	20.5	12.3	47.5	19.9	33.0	27.9	.9	12.1	3.9	4.6
.4	.3	.4	.3	.5	.2	.3	.2	(2)	.3	.2	.2

TABLE 8.04.--FLUID LOWFAT MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.3	14.0	24.2	21.7	21.5
ON ONLY 1 OF 3 DAYS.....PCT.....	4.2	1.8	3.4	2.4	2.7
ON ONLY 2 OF 3 DAYS.....PCT.....	4.5	1.1	3.1	5.3	5.2
ON ALL 3 OF 3 DAYS.....PCT.....	8.6	11.0	17.7	14.1	13.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	3.9	8.8	6.3	5.2	4.6
STANDARD DEVIATION.....DO.....	2.6	4.3	3.7	2.7	2.4
MAXIMUM.....DO.....	26.0	26.0	21.0	13.0	12.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	279	513	365	321	328
STANDARD DEVIATION.....GM.....	237	332	274	203	200
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	41	31	41	61	62
25TH.....GM.....	91	245	163	163	163
50TH.....GM.....	223	489	327	286	306
75TH.....GM.....	388	736	490	449	459
90TH.....GM.....	581	980	674	613	572
95TH.....GM.....	735	1,030	858	695	673
99TH.....GM.....	1,062	1,836	1,472	858	959
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	263	338	215	253	320
STANDARD DEVIATION.....GM.....	187	329	176	142	204
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	311	478	307	322	333
STANDARD DEVIATION.....GM.....	211	163	223	220	188
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	409	600	448	400	413
STANDARD DEVIATION.....GM.....	251	313	265	188	187
MAXIMUM IN ANY 1 DAY.....GM.....	4,410	2,083	1,960	1,960	1,348
MAXIMUM OVER 3 DAYS.....GM.....	6,125	5,513	5,145	3,063	3,430
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	217	175	173	185	213
STANDARD DEVIATION.....GM.....	130	76	78	79	85
PERCENTILES: <u>1/</u>					
5TH.....GM.....	25	61	61	61	92
25TH.....GM.....	123	123	123	123	123
50TH.....GM.....	245	184	153	184	245
75TH.....GM.....	245	245	245	245	245
90TH.....GM.....	368	245	245	245	245
95TH.....GM.....	490	245	245	306	368
99TH.....GM.....	613	276	368	490	490
MAXIMUM.....GM.....	1,960	368	735	613	551
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	68.5	.8	3.0	6.3	6.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	.1	.2	.3	.4

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
19.4	19.6	19.7	17.6	16.1	17.3	15.9	15.3	16.9	17.5	10.6	14.2
3.7	4.1	2.9	4.4	4.2	5.7	4.4	5.0	3.5	4.3	1.9	3.3
3.9	5.0	4.7	4.7	4.7	4.9	4.8	4.3	4.2	4.2	2.2	2.2
11.8	10.6	12.1	8.5	7.1	6.8	6.6	5.9	9.2	8.9	6.5	8.6
4.4	4.0	4.5	3.7	3.4	3.1	3.4	3.2	3.7	3.6	3.7	3.8
2.3	2.2	2.3	2.3	2.3	2.2	2.5	2.4	2.4	2.4	2.4	2.1
11.0	11.0	13.0	16.0	13.0	13.0	17.0	17.0	12.0	14.0	9.0	10.0
384	311	502	305	322	223	249	181	229	207	230	199
262	204	354	209	274	186	250	166	161	162	194	153
82	54	82	61	41	31	31	20	41	27	41	30
174	163	245	163	123	82	82	61	102	82	82	82
327	265	408	245	245	163	163	123	201	163	136	163
510	449	613	408	470	327	327	245	277	306	368	286
735	593	1,041	571	695	490	541	408	449	439	530	412
878	695	1,225	654	857	602	736	530	573	490	673	490
1,144	939	1,878	1,061	1,224	899	1,328	762	775	826	736	660
375	339	465	354	330	238	245	205	240	184	196	202
245	241	367	217	215	146	153	116	101	110	59	120
414	336	529	359	408	295	281	225	247	235	141	184
226	199	279	144	291	168	194	142	93	167	77	96
500	428	641	437	480	363	408	299	315	300	322	270
255	184	360	210	296	199	299	195	166	157	194	153
2,573	1,776	2,940	1,960	4,410	1,409	2,029	1,470	1,103	1,118	919	980
5,513	3,185	6,125	4,043	6,003	2,940	5,189	3,430	2,328	2,618	2,205	2,328
264	236	336	247	284	214	218	169	186	174	187	159
120	109	183	108	179	125	145	111	103	97	83	101
123	92	123	77	31	15	15	10	15	15	31	10
245	184	245	245	245	123	123	61	123	123	123	92
245	245	245	245	245	245	245	184	245	184	184	123
245	245	490	245	368	245	245	245	245	245	245	245
490	368	490	368	490	368	368	245	277	245	245	245
490	490	735	490	490	490	490	368	368	306	260	245
735	490	980	520	980	552	735	490	490	368	490	490
980	1,776	1,960	980	1,960	919	980	613	490	490	490	490
4.7	4.3	8.5	4.2	4.3	5.2	9.6	3.8	4.4	0	0	2.6
.3	.3	.7	.4	.2	.2	.4	.1	.6	0	0	.5

TABLE 8.05.--FLUID SKIM MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.6	2.3	4.2	4.0	3.9
ON ONLY 1 OF 3 DAYS.....PCT.....	1.7	0	.6	.7	.7
ON ONLY 2 OF 3 DAYS.....PCT.....	1.4	.9	.8	.8	.8
ON ALL 3 OF 3 DAYS.....PCT.....	2.6	1.4	2.8	2.5	2.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	3.6	8.0	6.3	5.4	4.4
STANDARD DEVIATION.....DO.....	2.6	5.1	3.3	2.9	2.4
MAXIMUM.....DO.....	21.0	21.0	13.0	13.0	9.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	226	380	374	336	309
STANDARD DEVIATION.....GM.....	201	230	245	206	189
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20			41	41
25TH.....GM.....	82			163	163
50TH.....GM.....	163			327	286
75TH.....GM.....	327			464	449
90TH.....GM.....	490			592	531
95TH.....GM.....	623			653	694
99TH.....GM.....	899			1,061	817
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	250	0	213	241	220
STANDARD DEVIATION.....GM.....	194	0	108	102	90
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	278	260	297	399	378
STANDARD DEVIATION.....GM.....	169	222	150	153	152
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	341	516	496	431	393
STANDARD DEVIATION.....GM.....	228	164	208	189	174
MAXIMUM IN ANY 1 DAY.....GM.....	2,940	827	1,103	1,225	980
MAXIMUM OVER 3 DAYS.....GM.....	4,410	2,435	2,818	3,185	2,450
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	190	143	179	186	209
STANDARD DEVIATION.....GM.....	123	57	75	76	82
PERCENTILES: <u>1/</u>					
5TH.....GM.....	15			61	61
25TH.....GM.....	123			123	123
50TH.....GM.....	245			184	245
75TH.....GM.....	245			245	245
90TH.....GM.....	306			245	245
95TH.....GM.....	368			245	368
99TH.....GM.....	490			490	490
MAXIMUM.....GM.....	980	260	735	490	613
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	20.7	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.5	4.3	4.3	5.3	3.4	6.2	5.5	8.2	6.3	9.1	2.6	7.4
.8	1.2	1.4	1.7	1.2	2.2	2.2	2.3	1.3	2.6	0	1.8
.8	1.1	.9	1.5	.8	1.5	1.3	2.3	1.5	2.0	.6	1.6
1.9	2.0	2.1	2.1	1.4	2.4	2.1	3.6	3.5	4.5	2.0	4.0
4.1	3.9	3.7	3.1	3.5	2.9	3.1	3.5	3.8	3.4	4.3	3.7
2.4	2.3	2.2	2.1	2.3	2.2	2.5	2.6	2.6	2.2	1.8	2.2
9.0	9.0	10.0	10.0	9.0	12.0	13.0	14.0	9.0	10.0	8.0	9.0
391	304	374	244	304	188	190	179	213	174	225	191
312	207	235	209	293	165	181	154	182	137	144	144
61	41	44	41	20	20	41	20	10	20		20
123	123	123	82	85	82	82	71	61	82		82
368	286	408	184	204	163	123	123	163	134		143
552	449	572	327	408	245	245	245	286	245		255
837	531	694	510	736	398	408	408	490	378		408
981	694	775	735	899	530	530	490	612	408		449
1,448	919	873	981	1,186	715	837	735	877	572		653
246	311	476	262	330	240	256	225	227	188	0	163
70	222	371	149	307	135	187	174	255	123	0	72
473	389	519	274	305	259	261	238	250	218	236	216
255	198	391	120	132	109	150	144	139	110	14	111
565	448	533	409	530	297	313	259	286	251	244	273
308	183	141	233	328	203	225	180	201	145	158	141
1,715	1,348	1,470	1,470	2,940	1,470	1,470	1,235	980	746	735	735
4,349	3,185	2,622	2,940	4,410	4,165	3,001	2,450	2,634	2,205	1,470	1,960
288	234	300	238	263	193	186	152	169	155	156	154
156	108	159	108	155	119	122	115	110	92	105	86
123	123	123	61	31	15	15	10	10	15		15
245	184	245	214	184	123	123	31	61	61		93
245	245	245	245	245	245	184	123	184	123		123
368	245	368	245	368	245	245	245	245	245		245
490	368	490	368	490	306	306	245	276	245		245
490	490	490	490	490	368	368	306	306	245		245
980	490	980	490	736	490	490	490	490	368		274
980	980	980	980	980	919	980	980	551	490	490	306
0	0	7.6	4.3	1.8	0	.9	1.6	0	0	0	4.5
0	0	3.4	1.8	.4	0	.1	.1	0	0	0	1.7

TABLE 8.06.--RECONSTITUTED DRY MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FCOD:					
AT LEAST CNCE IN 3 DAYS.....PCT.....	1.9	1.9	2.0	2.3	2.1
ON ONLY 1 OF 3 DAYS.....PCT.....	.6	0	.4	.4	.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	.4	.7	.9	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.8	1.5	1.0	.9	1.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER.....	3.4	9.7	5.0	4.0	3.8
STANDARD DEVIATION.....DO.....	2.5	5.0	3.2	2.6	2.1
MAXIMUM.....DO.....	22.0	22.0	15.0	9.0	9.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	206	515	262	234	223
STANDARD DEVIATION.....GM.....	198	302	247	185	149
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	14				
25TH.....GM.....	68				
50TH.....GM.....	163				
75TH.....GM.....	265				
90TH.....GM.....	449				
95TH.....GM.....	600				
99TH.....GM.....	981				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	191	0	213	197	166
STANDARD DEVIATION.....GM.....	162	0	205	114	62
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	258	364	269	271	348
STANDARD DEVIATION.....GM.....	147	30	181	143	183
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	322	583	389	363	283
STANDARD DEVIATION.....GM.....	233	301	282	203	138
MAXIMUM IN ANY 1 DAY.....GM.....	1,960	1,164	1,348	1,225	919
MAXIMUM OVER 3 DAYS.....GM.....	3,675	3,063	3,308	2,205	2,083
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	183	160	158	177	179
STANDARD DEVIATION.....GM.....	128	70	84	94	75
PERCENTILES: ^{1/}					
5TH.....GM.....	15				
25TH.....GM.....	123				
50TH.....GM.....	184				
75TH.....GM.....	245				
90TH.....GM.....	306				
95TH.....GM.....	368				
99TH.....GM.....	735				
MAXIMUM.....GM.....	980	245	368	490	490
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER.....	4.7	0	1.0	1.4	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	1.0	.9	0

^{1/} SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.3	1.4	1.2	1.2	1.5	1.6	1.7	2.3	3.5	3.5	2.3	3.3
.4	.5	.2	.3	.5	.5	.5	.7	.8	1.1	.4	1.3
.6	.5	.3	.3	.3	.6	.4	.6	.5	.8	.6	.4
.3	.4	.7	.6	.6	.5	.7	1.0	2.1	1.6	1.2	1.5
2.9	3.2	4.1	3.1	2.9	2.9	3.3	3.1	4.2	3.2	2.8	2.9
2.0	2.4	1.8	1.6	1.8	2.4	2.3	2.1	3.2	1.9	1.3	1.8
9.0	10.0	6.0	7.0	8.0	12.0	9.0	10.0	12.0	7.0	6.0	8.0
192	206	358	261	277	154	220	163	263	169	165	128
159	156	255	244	307	129	245	136	200	150	57	114
				3	5	3	10		10		
				61	61	56	41		41		
				184	113	163	123		143		
				408	245	245	245		245		
				898	327	490	368		432		
				980	416	817	425		459		
				1,225	571	1,199	602		613		
310	243	329	295	192	161	221	134	121	226	261	219
274	229	116	99	179	97	165	111	113	188	121	167
243	292	360	281	320	219	264	229	469	183	206	151
128	142	114	127	146	124	126	114	282	86	57	92
388	359	466	389	497	274	344	256	338	257	205	182
193	143	268	296	355	145	315	140	174	174	29	134
858	854	1,103	1,715	1,960	735	1,960	827	980	980	368	551
2,083	1,899	3,001	2,940	3,675	1,715	3,597	1,807	2,205	1,838	735	1,409
201	192	263	251	289	162	198	157	187	159	177	132
140	91	150	170	222	105	160	101	106	112	56	82
				31	8	10	15		5		
				123	93	61	61		61		
				245	152	245	164		123		
				337	245	245	245		245		
				490	274	306	245		245		
				736	306	490	306		306		
				980	490	980	490		490		
735	732	735	980	980	490	980	490	490	490	306	245
0	0	0	0	0	0	2.3	0	0	0	0	0
0	0	0	0	0	0	.8	0	0	0	0	0

TABLE 8.07.--CHOCOLATE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.5	.7	4.6	9.5	15.5
ON ONLY 1 OF 3 DAYS.....PCT.....	3.4	.7	3.5	6.2	8.6
ON ONLY 2 OF 3 DAYS.....PCT.....	1.4	0	.7	2.3	4.4
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	.4	.9	2.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.6	1.6	1.7
STANDARD DEVIATION.....DO.....	1.0	.0	1.3	1.0	1.0
MAXIMUM.....DO.....	10.0	1.0	9.0	6.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	159	53	98	119	152
STANDARD DEVIATION.....GM.....	128	18	64	83	99
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	63			42	63
25TH.....GM.....	83			63	83
50TH.....GM.....	125			83	104
75TH.....GM.....	167			146	167
90TH.....GM.....	333			250	250
95TH.....GM.....	365			333	333
99TH.....GM.....	583			417	500
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	305	158	226	236	267
STANDARD DEVIATION.....GM.....	160	55	121	104	86
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	310	0	216	248	293
STANDARD DEVIATION.....GM.....	146	0	50	124	116
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	346	0	230	266	296
STANDARD DEVIATION.....GM.....	209	0	80	85	104
MAXIMUM IN ANY 1 DAY.....GM.....	2,500	250	563	1,000	1,000
MAXIMUM OVER 3 DAYS.....GM.....	5,500	250	1,000	1,500	1,750
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	290	158	186	222	260
STANDARD DEVIATION.....GM.....	139	55	83	76	84
PERCENTILES: <u>1/</u>					
5TH.....GM.....	125			125	125
25TH.....GM.....	250			188	250
50TH.....GM.....	250			250	250
75TH.....GM.....	250			250	250
90TH.....GM.....	500			250	375
95TH.....GM.....	500			313	500
99TH.....GM.....	1,000			500	500
MAXIMUM.....GM.....	1,500	250	500	625	750
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	16.8	0	0	.8	1.6
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	0	.3	.3

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
18.2	16.9	12.6	11.7	4.0	3.1	1.6	.9	.9	.3	.5	.4
10.2	8.7	7.5	6.9	3.0	2.5	1.0	.7	.4	.2	.2	.4
5.3	6.0	3.6	3.6	.6	.4	.4	.1	.3	.1	0	0
2.7	2.3	1.4	1.2	.4	.2	.1	.1	.2	0	.3	0
1.8	1.8	1.6	1.6	1.5	1.4	1.6	1.3	1.9	1.3	2.2	1.0
1.2	1.1	.9	.8	.9	.9	1.0	.6	.9	.4	1.0	.0
9.0	10.0	5.0	6.0	5.0	6.0	7.0	3.0	3.0	2.0	3.0	1.0
182	165	191	152	168	134	188	114	217	105	185	83
154	122	139	96	137	97	236	53	142	37	81	0
83	83	83	83	63	63	63	56				
83	83	83	83	83	83	83	83				
167	167	167	125	156	83	125	83				
250	171	250	167	167	167	167	167				
333	333	333	250	333	250	333	167				
500	333	417	333	333	333	500	250				
958	500	646	458	833	500	1,833	250				
321	291	390	295	369	314	320	290	379	250	250	250
165	135	219	109	209	184	161	103	311	0	0	0
329	296	389	308	338	314	346	216	299	250	0	0
155	109	253	117	108	134	135	49	61	0	0	0
393	338	333	301	425	335	656	250	437	0	250	0
242	198	137	128	234	123	577	0	0	0	0	0
1,750	2,250	2,000	750	1,500	1,500	2,500	500	1,000	250	250	250
3,750	4,688	4,000	2,250	2,500	1,875	5,500	750	1,313	500	750	250
306	281	352	282	343	290	356	268	348	250	250	250
156	114	185	97	166	150	202	90	176	0	0	0
125	188	250	125	188	188	188	125				
250	250	250	250	250	250	250	250				
250	250	250	250	250	250	250	250				
250	250	500	250	500	250	500	250				
500	500	500	500	500	500	500	375				
500	500	626	500	500	500	1,000	500				
1,000	750	1,000	500	1,000	751	1,000	500				
1,125	1,000	1,250	500	1,000	1,500	1,000	500	1,000	250	250	250
5.5	0	0	3.1	0	2.2	0	1.6	2.0	0	0	0
.8	0	0	1.1	0	1.0	0	1.9	10.2	0	0	0

TABLE 8.08.--COCOA AND HOT CHOCOLATE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.8	0	4.1	5.8	8.7
ON ONLY 1 OF 3 DAYS.....PCT.....	3.0	0	3.0	4.1	5.4
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	1.0	.9	2.1
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	.2	.7	1.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.7	0	1.5	1.6	1.6
STANDARD DEVIATION.....DO.....	1.0	0	.9	1.1	.9
MAXIMUM.....DO.....	9.0	0	5.0	8.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	142	0	98	121	129
STANDARD DEVIATION.....GM.....	104	0	88	98	85
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	53			42	42
25TH.....GM.....	83			63	83
50TH.....GM.....	85			83	83
75TH.....GM.....	169			167	167
90TH.....GM.....	256			250	250
95TH.....GM.....	333			256	292
99TH.....GM.....	562			666	363
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	262	0	186	237	236
STANDARD DEVIATION.....GM.....	105	0	85	116	145
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	282	0	223	236	265
STANDARD DEVIATION.....GM.....	107	0	87	76	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	305	0	432	306	266
STANDARD DEVIATION.....GM.....	139	0	62	135	63
MAXIMUM IN ANY 1 DAY.....GM.....	1,563	0	750	1,250	1,563
MAXIMUM OVER 3 DAYS.....GM.....	3,250	0	1,500	2,000	1,563
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	255	0	196	228	236
STANDARD DEVIATION.....GM.....	90	0	78	67	61
PERCENTILES: <u>1/</u>					
5TH.....GM.....	128			125	125
25TH.....GM.....	250			188	219
50TH.....GM.....	250			250	250
75TH.....GM.....	256			250	250
90TH.....GM.....	313			256	256
95TH.....GM.....	500			275	256
99TH.....GM.....	513			500	500
MAXIMUM.....GM.....	1,250	0	500	500	576
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	6.9	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
7.8	8.0	7.5	7.1	4.0	5.5	2.8	3.5	2.3	2.9	2.2	4.8
4.3	4.9	5.1	4.5	2.6	3.2	1.7	2.4	.9	1.7	1.2	2.8
2.1	2.0	1.6	2.2	.8	1.4	.6	.8	.9	.4	.6	1.2
1.4	1.1	.8	.5	.6	.9	.5	.4	.6	.7	.4	.9
1.8	1.6	1.6	1.6	1.7	1.7	1.9	1.6	2.3	1.7	1.6	1.7
1.0	1.0	.8	1.2	1.1	.9	1.4	1.0	1.4	1.0	.8	.8
7.0	8.0	5.0	9.0	5.0	6.0	8.0	6.0	5.0	4.0	3.0	4.0
153	139	149	132	157	144	162	133	182	142	147	147
103	95	96	106	146	93	137	84	117	90	64	90
63	63	63	63	63	63	63	63		64		
83	83	83	83	83	83	83	83		83		
128	85	104	83	92	92	92	85		85		
169	167	171	167	167	171	206	169		167		
256	250	333	171	254	254	275	256		333		
339	271	333	252	363	313	513	333		335		
445	513	421	667	1,082	563	708	412		341		
275	267	301	252	272	267	251	271	246	243	312	275
97	88	152	55	100	93	77	107	17	33	146	98
269	266	360	249	317	284	298	302	284	295	250	252
86	118	125	60	140	121	93	105	116	114	0	3
302	295	276	442	382	272	370	268	320	257	250	293
128	109	91	196	242	99	171	96	108	72	0	94
1,000	1,024	1,000	1,000	1,500	1,500	1,125	768	549	512	500	576
2,500	2,500	1,375	2,250	3,250	1,706	2,125	1,250	1,250	1,024	750	1,500
259	253	282	247	283	259	260	250	240	245	271	266
83	73	128	55	125	105	85	86	26	65	89	75
160	188	125	172	188	160	156	125		128		
250	250	250	250	250	250	250	250		212		
250	250	250	250	250	250	250	250		250		
256	250	256	250	256	256	256	256		256		
375	275	500	256	375	375	375	275		256		
500	375	500	275	500	469	500	500		384		
500	500	1,000	500	751	550	512	512		500		
512	768	1,000	500	1,125	1,250	512	549	275	500	500	576
1.8	.8	.9	0	1.8	1.7	0	0	0	0	0	0
.6	.3	.6	0	.7	.3	0	0	0	0	0	0

TABLE 8.09.--NATURAL AND PROCESSED CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	40.0	6.1	35.9	37.0	35.3
ON ONLY 1 OF 3 DAYS.....PCT.....	25.0	4.9	23.3	23.7	23.1
ON ONLY 2 OF 3 DAYS.....PCT.....	10.8	.8	9.1	10.5	9.5
ON ALL 3 OF 3 DAYS.....PCT.....	4.2	.4	3.4	2.8	2.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.3	1.6	1.6	1.6
STANDARD DEVIATION.....DO.....	.9	.6	.9	.8	.9
MAXIMUM.....DO.....	8.0	3.0	8.0	5.0	7.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	11	17	16	18
STANDARD DEVIATION.....GM.....	19	11	13	11	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		4	3	4
25TH.....GM.....	9		9	9	9
50TH.....GM.....	19		9	13	13
75TH.....GM.....	28		19	19	23
90TH.....GM.....	46		35	29	37
95TH.....GM.....	57		45	37	47
99TH.....GM.....	95		65	49	66
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	42	22	31	34	36
STANDARD DEVIATION.....GM.....	29	22	19	19	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	46	26	38	34	38
STANDARD DEVIATION.....GM.....	29	3	20	17	20
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	49	40	36	34	42
STANDARD DEVIATION.....GM.....	29	9	16	16	26
MAXIMUM IN ANY 1 DAY.....GM.....	700	113	142	198	227
MAXIMUM OVER 3 DAYS.....GM.....	765	142	282	312	453
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	41	25	31	31	35
STANDARD DEVIATION.....GM.....	28	21	19	17	23
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		7	7	7
25TH.....GM.....	28		19	28	28
50TH.....GM.....	28		28	28	28
75TH.....GM.....	56		28	28	38
90TH.....GM.....	58		57	57	57
95TH.....GM.....	85		61	57	71
99TH.....GM.....	140		85	85	140
MAXIMUM.....GM.....	700	113	113	140	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	756.5	1.3	5.1	19.3	40.3
PROPORTION OF EATING OCCASIONS.....PCT.....	3.0	3.3	.8	2.0	4.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
31.2	34.9	39.0	39.8	47.0	45.5	43.1	42.6	34.5	35.8	32.8	31.9
22.8	24.5	25.4	26.4	27.5	27.5	25.2	27.4	21.2	22.1	17.9	19.0
6.6	8.7	9.4	10.0	13.3	12.8	12.0	11.5	8.8	10.0	9.0	9.3
1.8	1.8	4.2	3.4	6.2	5.2	6.0	3.7	4.6	3.7	5.9	3.5
1.4	1.5	1.6	1.6	1.8	1.7	1.7	1.6	1.6	1.6	1.7	1.6
.8	.8	1.0	.9	1.1	1.0	1.0	.9	.9	.8	.9	.8
5.0	6.0	6.0	6.0	8.0	8.0	7.0	7.0	6.0	6.0	5.0	4.0
18	18	25	19	29	23	26	20	24	20	25	19
14	14	22	16	25	20	24	15	21	15	21	13
5	5	7	5	7	5	6	5	5	5	5	5
9	9	9	9	9	9	9	9	9	9	9	9
15	13	19	19	19	19	19	19	19	19	19	19
21	20	28	26	37	28	33	28	29	28	34	23
37	37	47	38	57	47	56	38	47	38	56	38
47	42	66	52	76	66	66	52	57	47	61	52
73	77	112	84	117	95	112	75	122	76	136	57
40	38	49	37	53	41	48	40	44	40	42	39
21	25	34	24	37	29	33	26	28	30	30	25
45	40	51	43	56	47	53	42	50	38	61	35
23	22	29	21	39	25	37	22	29	17	43	15
44	42	57	47	60	52	53	43	49	43	33	38
21	23	36	23	32	28	34	21	33	18	12	15
227	284	395	227	339	310	700	252	255	228	306	170
280	312	529	349	765	423	765	414	510	255	408	210
39	35	46	37	49	40	46	37	44	37	43	36
22	23	30	23	33	27	37	23	28	22	36	21
14	7	14	10	14	10	14	14	14	14	5	14
28	28	28	28	28	28	28	28	28	28	28	28
28	28	28	28	43	28	28	28	28	28	28	28
56	56	57	56	57	57	57	56	57	56	57	56
57	57	84	57	85	61	84	57	84	57	85	57
84	81	112	84	113	85	112	84	91	84	113	58
113	113	170	113	170	128	168	116	170	113	170	113
168	284	227	227	315	280	700	244	170	170	306	170
33.7	50.7	29.4	46.4	94.4	115.7	108.3	148.8	19.9	22.8	11.0	9.5
3.6	4.5	3.3	5.0	2.9	2.8	3.0	3.1	3.1	2.3	3.6	1.9

TABLE 8.10.--CHEDDAR CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.4	.7	4.6	3.5	2.9
ON ONLY 1 OF 3 DAYS.....PCT.....	4.1	.7	3.0	2.8	2.5
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	1.4	.3	.5
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.2	.3	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.0	1.5	1.3	1.2
STANDARD DEVIATION.....DO.....	.7	.0	.7	.8	.4
MAXIMUM.....DO.....	6.0	1.0	3.0	5.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	20	4	16	15	15
STANDARD DEVIATION.....GM.....	17	1	12	9	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5			5	5
25TH.....GM.....	9			9	9
50TH.....GM.....	19			9	9
75TH.....GM.....	28			19	19
90TH.....GM.....	38			28	28
95TH.....GM.....	57			33	38
99TH.....GM.....	85			56	56
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	46	12	34	36	36
STANDARD DEVIATION.....GM.....	34	3	23	21	22
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	0	38	33	47
STANDARD DEVIATION.....GM.....	28	0	21	9	20
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	49	0	28	32	0
STANDARD DEVIATION.....GM.....	25	0	(2)	10	0
MAXIMUM IN ANY 1 DAY.....GM.....	306	14	113	113	113
MAXIMUM OVER 3 DAYS.....GM.....	510	14	142	168	169
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	45	12	33	34	38
STANDARD DEVIATION.....GM.....	31	3	23	18	22
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14			14	14
25TH.....GM.....	28			28	28
50TH.....GM.....	28			28	28
75TH.....GM.....	57			28	56
90TH.....GM.....	85			57	57
95TH.....GM.....	112			77	77
99TH.....GM.....	170			85	113
MAXIMUM.....GM.....	306	14	113	113	113
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	52.2	0	0	0	2.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.9	0	0	0	3.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.1	3.6	3.9	4.5	5.6	7.0	6.7	5.9	7.6	5.5	7.6	4.8
2.8	3.0	3.2	3.6	4.6	5.1	5.3	4.4	5.8	3.6	3.8	3.6
.3	.5	.7	.6	.7	1.5	1.1	1.2	1.1	1.3	1.5	1.1
0	0	0	.2	.2	.4	.3	.4	.7	.6	2.2	.1
1.1	1.2	1.2	1.3	1.3	1.5	1.3	1.4	1.5	1.5	1.8	1.3
.4	.5	.5	.7	.6	.8	.8	.7	1.1	.7	.9	.5
3.0	3.0	3.0	5.0	4.0	6.0	6.0	5.0	6.0	3.0	3.0	3.0
14	16	23	17	21	22	23	19	27	23	28	18
8	12	17	15	15	18	18	15	25	17	29	13
5	2	9	2	5	5	5	5	9	9		
9	9	9	9	9	9	9	9	17	9		
9	9	19	9	19	19	19	12	19	19		
19	19	28	23	28	28	28	21	28	28		
24	38	47	38	38	47	47	38	61	45		
28	38	66	47	47	57	61	47	76	64		
47	50	76	84	76	85	91	91	170	85		
40	42	65	40	54	45	52	40	56	53	53	48
24	33	50	31	39	32	41	28	24	32	44	33
30	37	41	45	49	56	58	43	55	33	70	32
10	22	27	15	17	28	26	23	33	12	80	17
0	0	0	51	48	48	54	48	79	59	32	64
0	0	0	24	21	22	25	19	46	19	6	0
142	150	227	140	227	254	272	204	170	142	306	136
142	168	227	253	255	310	272	272	510	255	408	192
37	38	56	39	50	45	51	40	54	46	47	42
22	27	46	26	33	31	37	26	26	27	54	31
14	7	28	13	14	14	14	14	21	26		
28	26	28	28	28	28	28	28	28	28		
28	28	28	28	56	28	43	28	56	28		
56	57	57	56	57	57	57	57	57	57		
57	57	113	57	85	84	85	68	85	85		
71	112	170	113	112	112	113	85	113	113		
113	136	227	140	227	170	227	113	113	136		
113	136	227	140	227	227	272	204	113	136	306	136
.9	3.7	2.1	3.5	1.8	7.3	14.6	9.7	1.0	2.1	0	3.0
1.3	3.9	3.2	3.9	.7	1.3	3.3	1.7	.8	1.5	0	4.8

TABLE 8.11.--SWISS CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.2	0	1.0	1.1	1.4
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	0	.4	.9	1.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.5	.2	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	0	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	0	1.6	1.2	1.7
STANDARD DEVIATION.....DO.....	.7	0	.5	.4	1.0
MAXIMUM.....DO.....	5.0	0	2.0	2.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	20	0	18	13	30
STANDARD DEVIATION.....GM.....	17	0	8	6	39
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5				
25TH.....GM.....	9				
50TH.....GM.....	19				
75TH.....GM.....	23				
90TH.....GM.....	38				
95TH.....GM.....	56				
99TH.....GM.....	94				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	45	0	42	31	43
STANDARD DEVIATION.....GM.....	29	0	28	14	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	49	0	32	32	30
STANDARD DEVIATION.....GM.....	31	0	8	7	14
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	59	0	0	0	97
STANDARD DEVIATION.....GM.....	30	0	0	0	50
MAXIMUM IN ANY 1 DAY.....GM.....	227	0	84	57	227
MAXIMUM OVER 3 DAYS.....GM.....	453	0	85	85	453
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	45	0	35	32	53
STANDARD DEVIATION.....GM.....	27	0	19	13	51
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14				
25TH.....GM.....	28				
50TH.....GM.....	28				
75TH.....GM.....	57				
90TH.....GM.....	85				
95TH.....GM.....	112				
99TH.....GM.....	140				
MAXIMUM.....GM.....	227	0	84	57	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	42.6	0	0	0	3.4
PROPORTION OF EATING OCCASIONS.....PCT.....	2.6	0	0	0	7.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.4	1.6	2.4	2.6	5.1	3.8	4.4	4.0	3.3	1.9	2.1	2.5
1.2	1.4	2.1	2.1	3.8	2.9	3.1	3.1	2.1	1.4	1.5	1.9
.1	.2	.2	.4	1.1	.8	1.0	.8	1.0	.5	.6	.6
(2)	.1	.1	.1	.1	.1	.2	.1	.1	0	0	0
1.2	1.2	1.2	1.3	1.4	1.3	1.5	1.3	1.4	1.3	1.3	1.2
.6	.4	.6	.7	.6	.5	.9	.6	.6	.4	.4	.4
4.0	3.0	4.0	4.0	4.0	3.0	5.0	4.0	3.0	2.0	2.0	2.0
18	14	23	17	25	18	23	18	26	18	30	15
13	10	19	14	19	14	21	14	16	10	13	6
				9	5	5	5				
				9	9	9	9				
				19	11	19	14				
				37	19	28	19				
				56	38	38	38				
				66	45	57	49				
				94	85	113	85				
43	40	59	35	56	40	44	42	57	40	71	42
25	29	42	14	37	24	25	27	20	15	26	20
53	28	58	48	61	40	67	38	47	48	71	28
25	10	35	29	34	20	44	16	9	13	0	(2)
54	21	85	48	59	67	37	62	85	0	0	0
0	0	0	10	33	20	13	17	0	0	0	0
113	170	168	113	224	168	227	170	113	84	113	85
170	170	255	170	339	255	340	255	255	113	142	85
43	36	57	38	53	41	46	41	55	43	71	37
26	27	40	15	28	25	24	24	20	17	22	17
				21	14	14	14				
				28	28	28	28				
				56	28	56	28				
				57	57	57	57				
				88	60	70	57				
				113	112	105	85				
				140	113	113	140				
113	170	168	57	142	113	168	170	113	84	113	85
0	1.3	0	1.4	13.7	6.7	1.5	12.7	0	0	0	2.1
0	3.1	0	2.7	4.9	2.6	.5	3.4	0	0	0	6.9

TABLE 8.12.--NATURAL CHEESE OTHER THAN CHEDDAR, SWISS, AND PARMESAN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.3	.5	1.8	2.1	1.7
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	.5	1.6	1.7	1.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	0	.2	.2	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	0	.2	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.0	1.2	1.3	1.2
STANDARD DEVIATION.....DO.....	.8	.0	.4	.6	.5
MAXIMUM.....DO.....	6.0	1.0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	5	14	14	15
STANDARD DEVIATION.....GM.....	20	4	8	7	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5				
25TH.....GM.....	9				
50TH.....GM.....	19				
75TH.....GM.....	28				
90TH.....GM.....	38				
95TH.....GM.....	57				
99TH.....GM.....	92				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	49	16	35	36	36
STANDARD DEVIATION.....GM.....	40	11	17	18	25
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	0	49	27	40
STANDARD DEVIATION.....GM.....	31	0	7	13	18
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	0	0	24	28
STANDARD DEVIATION.....GM.....	30	0	0	6	0
MAXIMUM IN ANY 1 DAY.....GM.....	680	28	57	57	113
MAXIMUM OVER 3 DAYS.....GM.....	737	28	113	85	113
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	46	16	35	32	36
STANDARD DEVIATION.....GM.....	35	11	18	17	22
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14				
25TH.....GM.....	28				
50TH.....GM.....	28				
75TH.....GM.....	57				
90TH.....GM.....	85				
95TH.....GM.....	113				
99TH.....GM.....	140				
MAXIMUM.....GM.....	680	28	57	57	113
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	38.9	0	0	.7	6.4
PROPORTION OF EATING OCCASIONS.....PCT.....	2.2	0	0	1.6	16.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.3	1.8	2.5	1.9	4.0	5.6	3.6	3.8	3.0	3.4	3.3	2.6
1.1	1.6	1.9	1.4	3.0	4.3	2.5	2.6	1.9	2.3	2.0	1.9
.1	.2	.5	.5	.6	.7	.8	.9	.8	.9	.8	.3
(2)	(2)	.1	.1	.4	.6	.3	.3	.4	.2	.5	.5
1.2	1.2	1.4	1.3	1.5	1.4	1.5	1.4	1.5	1.5	1.7	1.5
.5	.5	1.0	.5	.9	.8	.7	.7	.7	.7	1.1	.9
3.0	4.0	6.0	3.0	6.0	5.0	4.0	3.0	3.0	3.0	4.0	4.0
20	19	22	15	27	22	28	19	18	22	28	23
13	16	13	10	20	18	34	12	10	17	19	21
				6	5	9	5		9		
				14	9	9	9		9		
				19	19	19	19		19		
				38	28	28	26		28		
				53	43	56	37		56		
				57	57	75	42		62		
				113	76	227	62		66		
50	48	55	39	60	46	62	45	42	43	45	31
24	46	31	31	33	27	77	29	18	30	23	18
62	49	42	29	66	54	55	37	30	53	75	70
29	17	18	10	46	30	37	20	12	24	22	20
33	38	46	28	50	53	63	34	34	43	47	56
0	0	12	0	16	19	62	12	9	18	0	8
112	284	113	112	226	170	680	170	57	140	103	98
196	284	170	112	339	340	737	224	140	198	181	210
51	47	45	34	55	47	58	39	36	44	49	45
25	40	26	26	31	26	63	23	18	25	25	21
				14	14	14	14		15		
				28	28	28	28		28		
				57	56	38	28		28		
				57	57	57	57		57		
				113	85	113	62		84		
				113	112	140	85		85		
				128	128	284	113		113		
112	284	113	112	142	170	680	113	57	113	103	85
1.6	2.2	0	0	2.3	9.3	2.3	14.1	0	0	0	0
4.9	4.8	0	0	1.0	2.2	.9	3.7	0	0	0	0

TABLE 8.13.--PROCESSED CHEESE, CHEESE FOOD, AND CHEESE SPREAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	22.4	3.9	23.5	24.8	22.6
ON ONLY 1 OF 3 DAYS.....PCT.....	16.2	2.9	16.3	17.1	16.2
ON ONLY 2 OF 3 DAYS.....PCT.....	4.8	.6	5.1	6.4	5.2
ON ALL 3 OF 3 DAYS.....PCT.....	1.4	.4	2.1	1.3	1.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	1.4	1.5	1.4	1.4
STANDARD DEVIATION.....DO.....	.7	.7	.9	.7	.7
MAXIMUM.....DO.....	7.0	3.0	6.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	19	14	16	16	17
STANDARD DEVIATION.....GM.....	15	13	13	10	12
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		5	5	5
25TH.....GM.....	9		9	9	9
50TH.....GM.....	19		9	9	12
75TH.....GM.....	22		19	19	19
90TH.....GM.....	37		30	29	28
95TH.....GM.....	47		38	37	38
99TH.....GM.....	76		65	47	56
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	42	28	32	35	40
STANDARD DEVIATION.....GM.....	28	27	20	20	27
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	26	34	34	35
STANDARD DEVIATION.....GM.....	24	3	16	16	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	40	39	32	37
STANDARD DEVIATION.....GM.....	25	9	19	14	15
MAXIMUM IN ANY 1 DAY.....GM.....	700	113	140	142	168
MAXIMUM OVER 3 DAYS.....GM.....	765	142	282	227	280
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	40	29	32	33	36
STANDARD DEVIATION.....GM.....	26	23	19	17	22
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		9	14	14
25TH.....GM.....	28		21	28	28
50TH.....GM.....	28		28	28	28
75TH.....GM.....	56		28	31	43
90TH.....GM.....	57		57	57	57
95TH.....GM.....	85		78	57	71
99TH.....GM.....	132		85	85	140
MAXIMUM.....GM.....	700	113	112	140	168
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	126.1	0	0	3.7	6.8
PROPORTION OF EATING OCCASIONS.....PCT.....	1.0	0	0	.6	1.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.9	20.6	23.5	21.8	25.7	24.8	23.1	23.7	17.0	18.5	16.4	17.3
14.1	16.8	17.5	15.7	17.0	17.7	16.2	17.8	13.2	13.3	11.6	12.0
2.9	2.9	4.4	4.8	6.4	5.6	5.2	5.0	2.8	3.7	4.0	3.8
.9	.9	1.5	1.3	2.2	1.5	1.7	1.0	1.0	1.5	.9	1.5
1.3	1.3	1.4	1.4	1.5	1.5	1.4	1.4	1.3	1.4	1.4	1.4
.7	.7	.8	.8	.8	.8	.7	.7	.6	.7	.6	.7
4.0	5.0	5.0	5.0	6.0	7.0	5.0	6.0	4.0	3.0	4.0	4.0
18	16	21	18	25	19	22	17	18	16	18	17
13	12	17	14	19	15	20	12	11	11	13	10
7	5	7	5	9	5	7	5	5	5	2	5
9	9	9	9	9	9	9	9	9	9	9	9
14	9	19	15	19	19	19	13	19	9	15	19
19	19	28	19	28	23	28	19	19	19	20	19
37	28	38	37	47	38	38	31	37	29	37	28
43	38	57	47	65	50	56	39	42	38	38	38
71	75	85	76	103	75	85	65	57	56	61	57
41	40	47	39	52	40	49	39	44	35	39	39
19	24	30	22	33	25	40	22	28	23	25	23
47	39	51	41	51	45	47	43	43	39	43	33
23	16	28	19	31	22	26	23	16	13	23	10
42	44	48	49	53	47	50	37	33	33	37	33
19	22	24	17	27	23	37	15	8	10	12	12
140	227	196	210	315	258	700	170	170	228	122	170
245	280	298	241	504	308	765	308	170	228	183	170
40	37	45	38	48	40	46	37	41	36	40	35
21	20	27	21	32	25	36	22	25	19	24	19
14	14	14	14	18	14	15	14	14	14	9	14
28	28	28	28	28	28	28	28	28	28	28	28
28	28	28	28	40	28	28	28	28	28	28	28
56	56	56	56	57	56	57	56	56	45	56	56
57	57	84	57	84	57	76	57	57	57	61	57
84	70	112	61	112	85	85	84	84	57	85	57
113	113	168	113	170	122	142	116	142	113	122	70
140	140	170	210	315	258	700	170	170	158	122	170
7.7	12.3	8.1	10.5	11.6	11.6	14.6	26.7	8.0	.9	3.5	0
1.5	2.1	1.7	2.3	.7	.6	.9	1.2	3.2	.2	2.9	0

TABLE 8.14---COTTAGE CHEESE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.6	2.7	4.9	5.0	4.1
ON ONLY 1 OF 3 DAYS.....PCT.....	5.9	1.6	3.3	4.1	3.7
ON ONLY 2 OF 3 DAYS.....PCT.....	2.1	.5	1.5	.9	.4
ON ALL 3 OF 3 DAYS.....PCT.....	.6	.6	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.7	1.5	1.3	1.1
STANDARD DEVIATION.....DO.....	.8	1.0	.6	.5	.3
MAXIMUM.....DO.....	6.0	4.0	4.0	3.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	55	44	37	40	40
STANDARD DEVIATION.....GM.....	45	45	29	28	26
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	9		5	9	9
25TH.....GM.....	28		19	19	19
50TH.....GM.....	38		38	38	38
75TH.....GM.....	75		42	57	38
90TH.....GM.....	113		75	75	75
95TH.....GM.....	151		94	80	75
99TH.....GM.....	226		177	151	151
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	117	73	85	100	113
STANDARD DEVIATION.....GM.....	75	37	65	64	76
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	123	67	78	101	84
STANDARD DEVIATION.....GM.....	74	14	34	51	42
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	119	98	177	0	0
STANDARD DEVIATION.....GM.....	75	71	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	678	199	339	226	452
MAXIMUM OVER 3 DAYS.....GM.....	1,130	597	532	452	452
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	113	77	77	93	105
STANDARD DEVIATION.....GM.....	71	54	49	56	61
PERCENTILES: <u>1/</u>					
5TH.....GM.....	19		14	28	27
25TH.....GM.....	57		42	57	57
50TH.....GM.....	113		57	85	113
75TH.....GM.....	113		113	113	113
90TH.....GM.....	226		133	226	226
95TH.....GM.....	226		170	226	226
99TH.....GM.....	339		226	226	226
MAXIMUM.....GM.....	452	199	226	226	226
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	29.7	0	.9	0	4.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	1.2	0	4.8

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.2	4.5	3.9	4.6	5.3	8.4	9.9	13.8	11.1	17.3	7.8	16.7
2.5	3.8	3.2	4.0	3.9	5.8	7.0	9.1	6.4	10.7	6.2	8.5
.7	.6	.6	.5	1.3	2.2	2.4	3.5	2.9	5.3	.9	6.1
(2)	.1	.1	.1	.1	.4	.5	1.2	1.8	1.3	.8	2.1
1.2	1.2	1.2	1.2	1.3	1.5	1.4	1.5	1.8	1.6	1.3	1.7
.5	.5	.5	.6	.5	.8	.7	.8	1.2	.8	.6	.9
3.0	3.0	3.0	3.0	3.0	5.0	6.0	5.0	6.0	6.0	3.0	5.0
47	43	48	49	63	59	58	53	66	53	69	59
34	38	34	36	44	49	45	41	63	46	74	50
5	5	9	9	19	14	9	9	9	7		5
29	19	28	25	38	29	38	29	24	19		28
38	38	38	38	47	38	38	38	38	38		38
57	50	75	75	75	75	75	75	75	75		75
94	75	75	94	114	113	113	94	151	132		113
113	113	151	113	151	151	151	132	226	151		151
151	226	151	188	226	226	226	207	264	207		283
106	102	113	131	155	123	125	114	119	96	135	102
59	63	59	102	94	80	74	65	96	66	77	63
135	113	147	119	138	143	139	114	112	126	92	108
62	78	66	42	79	86	80	67	81	73	46	63
76	167	20	104	171	120	120	105	144	105	277	123
0	56	0	9	19	81	68	66	87	64	47	74
339	283	339	565	678	678	678	678	452	509	339	452
452	678	452	565	904	1,130	904	1,017	791	735	904	848
114	106	117	119	145	122	126	106	111	103	160	106
64	67	68	79	85	73	75	63	80	65	98	67
28	14	28	28	42	28	27	27	14	14		14
57	57	85	57	113	75	85	57	57	57		57
113	113	113	113	113	113	113	113	113	113		113
113	113	113	113	226	113	170	113	113	113		113
226	226	226	226	226	226	226	226	226	226		226
226	226	226	226	226	226	226	226	226	226		226
339	226	339	452	452	452	339	226	452	228		226
339	283	339	452	452	452	452	452	452	339	339	339
.8	2.8	0	0	.8	0	0	9.7	0	8.0	0	2.6
1.0	2.4	0	0	.3	0	0	.7	0	1.7	0	.9

TABLE 8.15.--FLUID CREAM
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.0	.1	.7	.6	.5
ON ONLY 1 OF 3 DAYS.....PCT.....	3.3	0	.6	.5	.3
ON ONLY 2 OF 3 DAYS.....PCT.....	1.6	0	.1	(2)	(2)
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	.1	0	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER.....	2.7	18.0	1.2	1.5	1.5
STANDARD DEVIATION.....DO.....	2.2	.0	.4	1.2	.8
MAXIMUM.....DO.....	19.0	18.0	2.0	5.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	23	685	25	24	32
STANDARD DEVIATION.....GM.....	35	0	15	23	30
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	2				
25TH.....GM.....	5				
50TH.....GM.....	12				
75TH.....GM.....	27				
90TH.....GM.....	50				
95TH.....GM.....	75				
99TH.....GM.....	140				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	31	0	65	55	110
STANDARD DEVIATION.....GM.....	36	0	44	42	99
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	32	0	60	60	60
STANDARD DEVIATION.....GM.....	29	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	43	685	0	48	12
STANDARD DEVIATION.....GM.....	54	0	0	39	10
MAXIMUM IN ANY 1 DAY.....GM.....	726	726	121	121	240
MAXIMUM OVER 3 DAYS.....GM.....	2,057	2,057	121	273	240
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	26	114	63	48	65
STANDARD DEVIATION.....GM.....	32	19	37	41	84
PERCENTILES: 1/					
5TH.....GM.....	5				
25TH.....GM.....	10				
50TH.....GM.....	15				
75TH.....GM.....	30				
90TH.....GM.....	60				
95TH.....GM.....	76				
99TH.....GM.....	161				
MAXIMUM.....GM.....	484	121	121	121	240
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER.....	108.2	0	1.0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	11.5	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.1	.9	2.9	2.7	9.0	6.9	11.8	9.9	11.3	10.3	11.5	11.9
.7	.7	1.8	1.6	4.2	3.9	5.6	4.4	5.3	4.2	2.7	4.9
.2	.2	.6	.8	2.2	1.5	2.6	2.1	2.3	2.6	4.1	2.3
.2	.1	.6	.3	2.5	1.5	3.7	3.4	3.6	3.6	4.6	4.7
1.8	1.4	1.7	2.0	2.5	2.3	3.0	2.8	2.4	2.8	3.4	2.7
1.3	.7	1.0	1.6	2.0	2.1	2.6	2.3	1.6	2.1	2.6	1.9
6.0	3.0	5.0	8.0	16.0	17.0	15.0	19.0	8.0	9.0	10.0	10.0
18	16	21	14	21	15	23	21	34	23	60	27
16	16	26	15	35	16	33	26	63	26	76	31
				2	2	2	2	3	2	3	2
				5	5	5	5	7	5	20	5
				12	10	10	13	10	15	40	20
				25	20	30	25	30	30	81	40
				40	35	55	50	82	55	131	55
				60	45	71	70	121	81	161	68
				140	76	130	140	242	101	364	176
41	36	47	29	29	25	34	27	33	32	87	33
34	29	65	39	31	23	41	27	28	42	70	27
48	29	22	26	31	23	34	30	28	33	60	39
37	17	17	21	32	21	27	28	21	28	52	30
14	37	43	29	39	30	42	38	77	38	97	44
4	31	35	12	56	23	48	34	96	33	102	40
121	121	242	180	484	165	484	240	726	242	545	218
240	240	360	180	1,119	334	1,452	625	1,634	489	1,091	529
30	33	36	21	25	19	24	23	42	25	53	30
33	29	46	25	30	16	31	24	57	28	55	34
				5	5	5	5	5	5	5	5
				10	10	10	10	15	10	15	15
				15	15	15	15	15	15	30	15
				30	30	30	30	30	30	66	30
				45	30	45	53	121	60	121	61
				61	60	61	61	182	66	182	121
				151	75	121	121	242	126	187	169
121	121	242	180	242	121	484	182	303	242	242	197
2.0	5.2	1.7	2.9	12.6	9.6	17.1	41.5	2.5	3.1	0	9.2
4.8	18.4	2.3	3.6	1.4	1.1	1.0	2.1	.8	.6	0	2.9

TABLE 8.16.--LIQUID AND FROZEN CREAM SUBSTITUTES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FCOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	1.7	0	0	.2	.2
ON ONLY 1 OF 3 DAYS.....PCT.....	.6	0	0	.1	.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	0	.1	0
ON ALL 3 OF 3 DAYS.....PCT.....	.7	0	0	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	3.1	0	0	1.6	1.8
STANDARD DEVIATION.....DO.....	2.7	0	0	.5	1.0
MAXIMUM.....DO.....	16.0	0	0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	0	0	52	58
STANDARD DEVIATION.....GM.....	28	0	0	17	53
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2				
25TH.....GM.....	5				
50TH.....GM.....	10				
75TH.....GM.....	27				
90TH.....GM.....	47				
95TH.....GM.....	77				
99TH.....GM.....	150				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	34	0	0	91	40
STANDARD DEVIATION.....GM.....	52	0	0	0	22
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	27	0	0	96	0
STANDARD DEVIATION.....GM.....	29	0	0	10	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	33	0	0	0	120
STANDARD DEVIATION.....GM.....	36	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	600	0	0	120	120
MAXIMUM OVER 3 DAYS.....GM.....	600	0	0	211	360
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	21	0	0	95	95
STANDARD DEVIATION.....GM.....	29	0	0	14	39
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5				
25TH.....GM.....	5				
50TH.....GM.....	10				
75TH.....GM.....	29				
90TH.....GM.....	60				
95TH.....GM.....	68				
99TH.....GM.....	120				
MAXIMUM.....GM.....	240	0	0	120	120
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	43.1	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	2.2	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
(2)	.1	.3	.5	1.3	1.4	2.9	2.8	3.7	3.3	3.9	3.5
(2)	(2)	.2	.3	.8	.7	1.0	.9	1.3	1.2	.5	.7
0	0	.1	.1	.2	.2	.7	.7	1.2	.7	1.2	1.1
0	.1	0	.1	.3	.5	1.2	1.3	1.2	1.4	2.2	1.8
1.0	2.2	1.3	3.0	2.3	3.4	3.2	3.3	2.8	2.5	4.2	2.8
.0	1.0	.5	3.7	2.6	3.4	2.3	2.9	2.1	1.4	3.1	1.4
1.0	3.0	2.0	11.0	14.0	16.0	11.0	16.0	9.0	7.0	12.0	6.0
2	71	47	11	10	17	21	21	29	20	29	35
0	57	25	6	6	29	29	26	31	18	28	45
				2	2	2	2		2		
				5	5	5	5		7		
				10	8	11	10		17		
				15	18	25	29		25		
				20	43	40	45		40		
				20	60	75	68		55		
				23	200	200	150		120		
5	15	97	27	24	45	29	30	36	47	11	13
0	0	39	19	16	106	36	33	19	44	6	8
0	0	120	15	17	32	24	17	27	27	39	58
0	0	0	0	9	45	20	15	22	11	37	37
0	120	0	18	15	19	33	34	58	25	37	44
0	0	0	0	3	16	40	33	36	23	30	58
5	120	120	60	60	600	240	240	180	120	90	240
5	360	240	60	68	600	600	450	375	360	269	600
5	99	108	11	13	15	20	19	32	24	21	37
0	42	31	13	10	22	31	24	31	28	20	50
				3	3	5	5		5		
				5	5	5	5		10		
				10	5	11	10		15		
				15	15	15	15		30		
				30	43	45	60		60		
				30	60	60	60		120		
				60	120	180	120		120		
5	120	120	60	60	120	240	166	180	120	80	240
0	.9	0	0	0	.9	11.3	25.5	2.2	0	2.2	0
0	19.7	0	0	0	.3	2.5	3.8	1.9	0	2.5	0

TABLE 8.17.--POWDERED CREAM SUBSTITUTES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.3	0	0	.2	.4
ON ONLY 1 OF 3 DAYS.....PCT.....	1.2	0	0	0	.4
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	0	.2	0
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	3.4	0	0	2.0	1.0
STANDARD DEVIATION.....DO.....	2.6	0	0	.0	.0
MAXIMUM.....DO.....	20.0	0	0	2.0	1.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	4	0	0	4	3
STANDARD DEVIATION.....GM.....	7	0	0	5	3
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1				
25TH.....GM.....	1				
50TH.....GM.....	3				
75TH.....GM.....	5				
90TH.....GM.....	9				
95TH.....GM.....	12				
99TH.....GM.....	27				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	5	0	0	0	8
STANDARD DEVIATION.....GM.....	6	0	0	0	10
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	5	0	0	6	0
STANDARD DEVIATION.....GM.....	6	0	0	7	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	6	0	0	0	0
STANDARD DEVIATION.....GM.....	9	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	106	0	0	24	28
MAXIMUM OVER 3 DAYS.....GM.....	317	0	0	36	28
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	4	0	0	6	8
STANDARD DEVIATION.....GM.....	5	0	0	8	10
PERCENTILES: <u>1/</u>					
5TH.....GM.....	1				
25TH.....GM.....	2				
50TH.....GM.....	2				
75TH.....GM.....	4				
90TH.....GM.....	6				
95TH.....GM.....	12				
99TH.....GM.....	28				
MAXIMUM.....GM.....	94	0	0	24	28
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	67.4	0	0	2.1	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.2	0	0	28.1	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
.2	.2	.6	1.7	4.6	5.0	6.6	7.4	4.8	8.3	5.6	6.6
.2	.1	.4	.6	1.5	1.9	1.8	1.9	.8	1.5	.8	.9
(2)	(2)	0	.4	1.3	1.1	1.6	1.6	1.0	1.9	1.1	1.2
(2)	.1	.1	.6	1.8	2.0	3.3	3.9	3.1	4.9	3.6	4.5
1.5	2.2	1.7	2.6	3.1	3.1	3.4	3.6	3.7	3.6	3.8	4.3
.8	.9	.8	1.8	2.7	2.6	2.4	2.7	2.1	2.9	2.7	2.6
3.0	3.0	3.0	7.0	20.0	12.0	16.0	20.0	9.0	17.0	12.0	10.0
2	2	2	2	4	3	4	4	4	4	4	11
1	1	1	1	5	4	4	5	3	6	3	24
				1	1	1	1	1	1		1
				1	1	1	2	1	1		2
				2	2	3	3	3	2		4
				5	4	6	5	5	5		9
				10	8	10	9	8	9		14
				12	10	14	12	9	12		57
				28	20	22	24	12	36		106
4	2	5	4	4	4	5	6	3	3	3	4
3	0	1	2	4	3	8	6	1	1	2	2
2	2	0	4	7	5	5	5	4	4	3	7
0	0	0	2	10	5	4	5	4	3	1	6
3	3	5	3	6	6	6	6	5	6	5	15
0	1	(2)	1	6	5	5	5	3	7	3	28
10	4	6	12	80	72	57	71	20	98	12	106
10	12	15	16	133	116	113	135	37	108	29	317
3	2	4	3	4	3	4	4	3	4	3	8
2	1	2	1	4	3	4	4	2	6	2	11
				2	1	1	1	2	1		2
				2	2	2	2	2	2		2
				2	2	2	2	2	2		3
				6	4	4	4	4	4		6
				6	6	8	6	6	6		35
				12	8	12	9	6	8		35
				18	18	18	24	12	12		35
10	4	6	6	28	47	28	53	18	94	8	35
0	0	0	0	5.0	27.1	19.3	8.3	0	4.7	0	.8
0	0	0	0	.9	3.2	1.7	.4	0	.9	0	.3

TABLE 8.18.--ICE CREAM AND ICE MILK
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	23.6	3.7	24.0	29.8	30.9
ON ONLY 1 OF 3 DAYS.....PCT.....	15.9	3.3	16.9	19.1	21.1
ON ONLY 2 OF 3 DAYS.....PCT.....	5.6	.3	5.8	8.5	7.2
ON ALL 3 OF 3 DAYS.....PCT.....	2.0	.1	1.3	2.1	2.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	1.1	1.4	1.5	1.5
STANDARD DEVIATION.....DO.....	.8	.4	.7	.7	.7
MAXIMUM.....DO.....	7.0	3.0	4.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	57	17	32	43	50
STANDARD DEVIATION.....GM.....	45	14	23	28	34
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		11	11	17
25TH.....GM.....	22		22	22	22
50TH.....GM.....	44		22	44	44
75TH.....GM.....	67		44	44	66
90TH.....GM.....	111		64	89	89
95TH.....GM.....	133		89	89	122
99TH.....GM.....	222		133	133	177
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	118	47	72	94	107
STANDARD DEVIATION.....GM.....	70	42	35	50	54
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	122	47	69	87	102
STANDARD DEVIATION.....GM.....	68	0	35	41	44
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	129	33	79	85	114
STANDARD DEVIATION.....GM.....	68	0	44	33	47
MAXIMUM IN ANY 1 DAY.....GM.....	1,064	133	233	360	399
MAXIMUM OVER 3 DAYS.....GM.....	1,596	133	466	532	798
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	116		69	86	101
STANDARD DEVIATION.....GM.....	66		36	43	47
PERCENTILES: <u>1/</u>					
5TH.....GM.....	33		17	33	33
25TH.....GM.....	67		44	67	67
50TH.....GM.....	133		67	67	100
75TH.....GM.....	133		67	133	133
90TH.....GM.....	200		133	133	133
95TH.....GM.....	266		133	133	175
99TH.....GM.....	333		133	266	266
MAXIMUM.....GM.....	532	133	200	266	267
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	93.2	0	2.4	1.7	8.4
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	.7	.2	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
31.5	30.1	27.1	23.6	19.9	18.8	26.4	19.4	27.6	21.9	21.3	27.7
20.1	20.2	16.6	16.8	14.7	14.3	17.1	13.7	15.9	14.0	9.7	15.3
8.2	7.7	8.4	5.8	3.9	3.5	6.4	4.3	7.1	5.2	7.8	8.4
3.2	2.2	2.0	1.0	1.3	1.1	2.9	1.4	4.6	2.6	3.9	4.0
1.6	1.5	1.5	1.4	1.4	1.4	1.5	1.4	1.7	1.5	1.9	1.7
.8	.8	.8	.7	.8	.7	.8	.7	.9	.8	1.0	1.0
7.0	5.0	5.0	4.0	7.0	5.0	5.0	5.0	6.0	5.0	6.0	6.0
68	58	82	57	67	50	67	49	64	52	66	53
49	43	70	47	50	35	53	35	49	41	40	42
18	17	22	17	19	15	17	11	15	11	22	11
37	28	44	22	33	22	33	22	23	22	44	22
44	44	67	44	44	44	44	44	44	44	55	44
89	67	89	67	89	67	89	67	89	67	89	67
133	100	155	100	133	89	133	89	133	100	133	111
177	133	200	133	156	122	155	124	155	133	133	133
244	222	355	222	266	177	266	177	266	222	222	222
139	121	149	121	145	112	132	109	114	102	121	89
82	69	77	66	81	58	77	67	66	51	64	46
142	129	168	133	160	119	140	104	109	98	120	99
71	73	93	64	86	53	76	45	49	48	52	48
134	116	216	155	153	116	144	114	140	119	99	114
66	58	120	111	62	52	72	47	55	61	45	54
665	665	1,064	532	532	532	532	798	399	333	266	532
1,131	1,032	1,596	1,264	1,064	931	1,197	798	798	732	665	798
130	116	159	123	142	110	131	104	115	100	103	92
68	60	92	75	79	57	73	57	60	52	54	47
50	34	67	44	50	33	50	33	33	28	33	33
67	67	100	67	82	67	67	67	67	67	67	67
133	133	133	133	133	133	133	100	131	67	131	67
133	133	200	133	141	133	133	133	133	133	133	133
262	200	266	200	266	134	266	133	200	133	133	133
266	266	357	266	266	227	266	200	266	200	200	133
393	266	532	402	399	266	399	266	266	266	266	266
399	399	532	532	532	532	532	532	399	266	266	266
5.5	10.8	3.0	1.8	9.0	14.0	9.1	8.2	7.1	7.9	0	4.3
.5	1.1	.5	.4	.8	1.0	.5	.4	1.4	1.4	0	.9

TABLE 8.19.--NONDAIRY WHIPPED TOPPING
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.4	.2	1.3	1.8	2.0
ON ONLY 1 OF 3 DAYS.....PCT.....	2.0	.2	1.3	1.7	1.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	0	.1	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER.....	1.2	1.0	1.0	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.0	.0	.4	.4
MAXIMUM.....DO.....	4.0	1.0	1.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	5	3	3	4	4
STANDARD DEVIATION.....GM.....	7	0	2	3	4
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	1				
25TH.....GM.....	2				
50TH.....GM.....	3				
75TH.....GM.....	6				
90TH.....GM.....	10				
95TH.....GM.....	13				
99TH.....GM.....	31				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	12	9	9	11	11
STANDARD DEVIATION.....GM.....	12	0	7	8	12
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	15	0	0	9	7
STANDARD DEVIATION.....GM.....	22	0	0	7	5
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	17	0	0	0	0
STANDARD DEVIATION.....GM.....	9	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	160	9	25	38	57
MAXIMUM OVER 3 DAYS.....GM.....	320	9	25	38	57
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	12	9	9	11	10
STANDARD DEVIATION.....GM.....	15	0	7	8	11
PERCENTILES: 1/					
5TH.....GM.....	2				
25TH.....GM.....	5				
50TH.....GM.....	9				
75TH.....GM.....	14				
90TH.....GM.....	20				
95TH.....GM.....	38				
99TH.....GM.....	75				
MAXIMUM.....GM.....	160				
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER.....	168.3	0	1.8	2.5	4.2
PROPORTION OF EATING OCCASIONS.....PCT.....	15.6	0	12.6	7.5	10.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.1	2.6	1.9	2.1	2.0	1.7	2.2	2.9	3.6	4.0	3.7	4.5
1.9	2.5	1.4	1.8	1.8	1.5	1.8	2.5	3.0	2.8	1.6	3.2
.2	.1	.5	.2	.2	.1	.3	.3	.6	1.1	1.7	1.2
0	0	0	.1	0	(2)	.1	(2)	0	.1	.5	0
1.1	1.0	1.3	1.2	1.1	1.1	1.3	1.1	1.3	1.4	1.8	1.3
.3	.2	.6	.5	.3	.3	.7	.4	.5	.6	.8	.7
2.0	2.0	3.0	3.0	2.0	3.0	4.0	3.0	3.0	4.0	4.0	4.0
3	4	7	5	7	3	7	5	6	5	8	3
2	3	7	5	6	3	12	9	8	6	5	2
	1			1	1	1	1		1		
	2			3	2	3	2		2		
	3			6	3	3	3		3		
	5			7	3	6	5		6		
	6			13	7	14	9		10		
	7			18	11	17	16		13		
	13			35	13	38	41		47		
8	11	16	11	18	10	13	12	11	11	17	9
6	8	13	7	17	8	17	13	7	8	12	5
7	7	16	10	18	9	25	21	26	9	12	7
3	2	15	2	9	3	38	34	21	6	5	5
0	0	0	30	0	11	14	19	0	21	15	0
0	0	0	0	0	0	5	5	0	18	1	0
30	38	56	38	106	40	160	160	75	75	38	23
30	38	94	89	106	40	320	320	131	141	50	38
8	10	15	12	17	9	15	14	14	11	13	8
6	8	12	9	13	8	22	19	15	9	8	6
	3			4	2	4	2		2		
	5			9	5	8	5		5		
	9			16	9	9	9		9		
	10			20	9	14	14		15		
	19			38	20	20	25		19		
	21			38	28	38	40		26		
	38			80	40	160	75		38		
	38	42	38	80	40	160	160	75	75	38	23
8.7	8.3	3.2	7.1	11.9	13.0	28.1	42.6	14.3	7.3	10.4	4.9
18.0	14.0	9.4	19.7	13.2	13.4	19.3	18.2	27.7	7.5	29.2	8.3

TABLE 8.20.--SOUR CREAM
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.9	.2	1.1	1.2	1.3
ON ONLY 1 OF 3 DAYS.....PCT.....	2.6	.2	1.1	1.2	1.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.2	0	0	.1	0
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.0	1.0	1.0	1.0
STANDARD DEVIATION.....CO.....	.4	.0	.0	.2	.0
MAXIMUM.....DO.....	4.0	1.0	1.0	2.0	1.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	13	2	3	8	7
STANDARD DEVIATION.....GM.....	16	0	2	9	8
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2				
25TH.....GM.....	4				
50TH.....GM.....	8				
75TH.....GM.....	16				
90TH.....GM.....	29				
95TH.....GM.....	38				
99TH.....GM.....	77				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	33	5	10	24	20
STANDARD DEVIATION.....GM.....	35	0	6	27	24
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	41	0	0	24	0
STANDARD DEVIATION.....GM.....	36	0	0	0	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	72	0	0	0	0
STANDARD DEVIATION.....GM.....	67	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	345	5	24	115	115
MAXIMUM OVER 3 DAYS.....GM.....	690	5	24	115	115
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	35	5	10	24	20
STANDARD DEVIATION.....GM.....	38	0	6	26	24
PERCENTILES: <u>1/</u>					
5TH.....GM.....	6				
25TH.....GM.....	12				
50TH.....GM.....	24				
75TH.....GM.....	36				
90TH.....GM.....	72				
95TH.....GM.....	115				
99TH.....GM.....	230				
MAXIMUM.....GM.....	345	5	24	115	115
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	58.6	0	1.2	3.3	2.1
PROPORTION OF EATING OCCASIONS.....PCT.....	4.8	0	9.7	14.8	8.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.5	1.3	2.1	1.7	3.1	3.9	3.8	4.0	2.7	4.0	1.5	1.9
1.3	1.2	2.1	1.7	2.8	3.6	3.5	3.6	2.5	3.3	1.5	1.7
.1	.1	0	0	.3	.3	.2	.3	.2	.5	0	0
0	0	0	0	0	0	(2)	.1	0	.2	0	.2
1.1	1.1	1.0	1.0	1.1	1.1	1.1	1.1	1.1	1.3	1.0	1.2
.3	.3	.0	.0	.3	.3	.4	.4	.3	.7	.0	.6
2.0	2.0	1.0	1.0	2.0	2.0	4.0	3.0	2.0	4.0	1.0	3.0
18	12	15	13	14	12	15	11	11	18	7	15
15	16	11	9	15	12	18	19	8	22	2	18
				3	3	4	2		3		
				4	4	4	4		4		
				8	8	8	8		8		
				16	14	16	12		19		
				38	27	38	20		54		
				38	38	40	29		57		
				85	77	115	66		115		
50	31	44	38	35	35	41	26	30	35	21	29
42	45	34	26	31	37	51	23	21	30	7	30
50	49	0	0	49	27	48	30	41	74	0	0
20	20	0	0	45	15	35	23	0	56	0	0
0	0	0	0	0	0	40	113	0	48	0	57
0	0	0	0	0	0	0	97	0	15	0	0
154	230	115	115	230	230	345	230	115	230	24	115
154	230	115	115	254	230	345	690	115	345	24	170
50	32	44	38	38	33	41	30	31	42	21	37
39	33	34	26	41	34	47	35	21	42	7	29
				10	7	12	6		10		
				12	12	12	12		12		
				24	24	24	24		24		
				36	36	58	28		58		
				87	58	87	58		115		
				115	115	115	76		115		
				230	151	230	230		230		
154	115	115	115	230	230	345	230	115	230	24	115
3.7	2.3	4.0	1.0	7.0	12.5	4.8	9.5	0	5.4	0	1.8
11.1	7.5	13.7	4.3	5.1	5.3	2.4	3.0	0	5.9	0	8.0

TABLE 8.21.--YOGURT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.2	2.9	3.8	2.9	2.4
ON ONLY 1 OF 3 DAYS.....PCT.....	1.9	.6	2.7	1.9	1.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	1.3	.8	.7	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.4	1.1	.3	.3	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.6	2.4	1.4	1.6	1.3
STANDARD DEVIATION.....DO.....	.9	1.1	.7	.7	.5
MAXIMUM.....DO.....	6.0	5.0	3.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	99	62	55	71	74
STANDARD DEVIATION.....GM.....	76	57	46	54	54
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	16			20	
25TH.....GM.....	41			38	
50TH.....GM.....	82			57	
75TH.....GM.....	143			82	
90TH.....GM.....	204			143	
95TH.....GM.....	245			163	
99TH.....GM.....	378			306	
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	187	139	122	168	150
STANDARD DEVIATION.....GM.....	102	64	82	105	78
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	198	48	149	146	221
STANDARD DEVIATION.....GM.....	94	25	103	59	74
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	198	106	70	101	0
STANDARD DEVIATION.....GM.....	115	71	20	109	0
MAXIMUM IN ANY 1 DAY.....GM.....	735	245	490	490	613
MAXIMUM OVER 3 DAYS.....GM.....	1,593	681	735	918	735
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	182	78	117	137	173
STANDARD DEVIATION.....GM.....	94	66	91	84	82
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31			31	
25TH.....GM.....	123			61	
50TH.....GM.....	227			123	
75TH.....GM.....	245			184	
90TH.....GM.....	245			245	
95TH.....GM.....	245			245	
99TH.....GM.....	490			490	
MAXIMUM.....GM.....	613	245	490	490	368
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	2.4	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.4	1.6	2.0	2.5	3.5	5.0	2.0	4.3	1.6	4.3	0	2.5
1.2	1.3	1.5	1.6	2.1	2.9	1.0	2.5	1.0	2.4	0	1.2
.2	.3	.3	.8	.9	1.4	.6	1.3	.1	1.4	0	.7
0	0	.2	.1	.5	.7	.4	.6	.5	.5	0	.6
1.3	1.3	1.4	1.4	1.6	1.7	1.8	1.6	1.9	1.6	0	1.8
.5	.5	.7	.6	.8	1.0	1.0	.8	1.0	.8	0	.9
3.0	3.0	3.0	3.0	6.0	6.0	5.0	4.0	4.0	6.0	0	3.0
81	88	109	98	112	107	118	94	141	106	0	82
38	59	55	69	83	78	77	73	84	107	0	64
				27	20	27	15		7		
				61	57	76	41		41		
				82	82	82	82		76		
				157	163	163	123		151		
				245	206	227	163		245		
				265	245	245	245		327		
				327	378	490	327		490		
204	196	257	188	190	197	214	173	362	174	0	161
58	110	107	79	72	93	109	92	246	123	0	81
231	259	215	224	230	211	205	187	64	188	0	183
44	77	20	108	90	70	80	97	0	136	0	56
0	0	223	245	240	216	192	189	199	292	0	94
0	0	17	0	93	123	98	104	55	156	0	102
454	490	454	490	613	735	554	490	735	735	0	306
624	735	717	980	1,593	1,516	1,470	1,470	735	1,470	0	674
191	209	234	206	209	189	192	173	219	193	0	136
61	94	74	91	87	83	93	88	124	130	0	91
				61	46	46	29		29		
				123	123	123	114		114		
				227	227	227	184		184		
				245	245	245	245		245		
				245	245	245	245		490		
				368	245	245	245		490		
				490	490	554	490		490		
245	490	454	490	490	613	554	490	490	490	0	245
0	0	0	0	0	0	0	2.4	0	0	0	0
0	0	0	0	0	0	0	.5	0	0	0	0

TABLE 9.01.--WHITE SUGAR
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	43.6	8.8	37.1	42.3	43.0
ON ONLY 1 OF 3 DAYS.....PCT.....	14.3	5.3	16.7	18.9	18.5
ON ONLY 2 OF 3 DAYS.....PCT.....	10.7	1.7	12.0	13.7	13.0
ON ALL 3 OF 3 DAYS.....PCT.....	18.7	1.8	8.4	9.8	11.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	3.3	1.8	2.1	2.2	2.1
STANDARD DEVIATION.....DO.....	2.7	1.1	1.4	1.7	1.3
MAXIMUM.....DO.....	26.0	6.0	11.0	18.0	10.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	10	3	5	6	6
STANDARD DEVIATION.....GM.....	12	3	5	7	6
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1		1	1	1
25TH.....GM.....	3		1	3	3
50TH.....GM.....	6		3	4	4
75TH.....GM.....	13		6	7	8
90TH.....GM.....	21		10	13	14
95TH.....GM.....	30		13	17	19
99TH.....GM.....	55		29	38	33
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	11	6	7	10	9
STANDARD DEVIATION.....GM.....	10	6	7	12	7
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	12	8	8	9	11
STANDARD DEVIATION.....GM.....	11	6	7	6	9
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	16	3	8	12	11
STANDARD DEVIATION.....GM.....	15	2	7	11	8
MAXIMUM IN ANY 1 DAY.....GM.....	405	38	75	150	102
MAXIMUM OVER 3 DAYS.....GM.....	980	42	125	213	150
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	9	5	7	8	9
STANDARD DEVIATION.....GM.....	9	5	6	8	8
PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		2	2	4
25TH.....GM.....	4		4	4	4
50TH.....GM.....	8		4	6	8
75TH.....GM.....	8		8	8	8
90TH.....GM.....	17		13	13	13
95TH.....GM.....	25		13	25	25
99TH.....GM.....	50		28	38	38
MAXIMUM.....GM.....	350	38	50	150	100
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	4284.7	6.9	71.6	193.2	145.1
PROPORTION OF EATING OCCASIONS.....PCT.....	8.0	8.6	8.8	12.1	8.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
44.6	41.4	43.4	39.0	45.8	44.4	49.7	41.1	50.8	43.5	56.0	41.0
19.6	19.1	19.2	17.8	14.5	14.8	11.1	11.6	10.5	11.2	10.9	8.9
13.9	13.4	11.9	11.8	10.7	10.9	10.3	8.2	11.8	9.0	11.8	8.0
11.1	9.0	12.3	9.4	20.6	18.8	28.3	21.3	28.6	23.2	33.3	24.0
2.1	2.1	2.3	2.3	3.4	3.3	4.3	3.8	3.9	3.7	4.1	3.7
1.4	1.3	1.6	1.7	2.8	2.8	3.2	2.9	2.8	2.9	2.7	2.4
9.0	12.0	16.0	12.0	24.0	26.0	21.0	23.0	17.0	23.0	13.0	14.0
8	7	10	8	12	9	13	10	13	9	12	9
9	7	11	9	15	11	14	15	13	9	11	9
1	1	1	1	1	1	1	1	1	1	1	1
3	3	3	3	4	3	4	3	4	3	4	3
6	4	6	5	8	6	8	7	8	6	8	7
9	8	13	8	15	11	17	13	16	11	15	11
17	15	21	17	25	19	28	22	31	18	25	18
24	19	29	26	36	28	41	29	39	25	35	25
42	34	56	38	75	53	67	54	67	44	47	50
12	10	14	11	14	11	11	9	12	9	10	9
10	9	15	14	13	9	8	8	14	9	10	9
12	12	16	13	14	12	12	10	11	8	11	10
10	11	16	11	14	15	11	9	7	6	9	7
14	12	17	15	19	15	19	16	18	13	16	12
15	8	14	12	19	12	15	19	15	10	11	10
225	150	150	188	350	228	240	405	125	100	106	75
425	179	350	304	700	400	363	980	271	171	190	156
11	10	13	10	11	9	9	8	10	7	8	7
10	8	13	8	13	8	8	8	9	6	6	6
4	4	4	4	4	3	3	2	4	2	2	2
4	4	8	4	4	4	4	4	4	4	4	4
8	8	8	8	8	8	8	8	8	4	8	4
13	13	13	13	13	8	8	8	12	8	8	8
25	17	25	21	21	16	17	15	17	13	17	13
25	25	38	25	25	24	25	25	25	17	25	25
50	50	66	50	50	38	38	38	50	38	32	32
100	100	113	100	350	170	200	170	100	69	50	50
186.4	148.4	161.0	127.2	583.8	643.7	857.6	761.0	115.1	150.9	19.3	113.7
9.3	8.0	11.7	9.5	9.7	8.1	8.2	6.9	5.2	5.4	1.6	7.6

TABLE 9.02.--SIRUP

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	10.8	4.0	14.5	18.2	17.9
ON ONLY 1 OF 3 DAYS.....PCT.....	9.1	2.8	12.9	15.4	14.7
ON ONLY 2 OF 3 DAYS.....PCT.....	1.3	.3	1.3	2.2	2.4
ON ALL 3 OF 3 DAYS.....PCT.....	.4	.9	.3	.5	.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	3.2	1.1	1.2	1.2
STANDARD DEVIATION.....DO.....	.6	4.3	.4	.5	.5
MAXIMUM.....DO.....	17.0	17.0	4.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	28	21	15	22	25
STANDARD DEVIATION.....GM.....	22	22	13	17	19
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		2	4	7
25TH.....GM.....	13		7	13	13
50TH.....GM.....	26		13	17	26
75TH.....GM.....	31		26	27	27
90TH.....GM.....	53		27	40	53
95TH.....GM.....	62		40	53	62
99TH.....GM.....	113		55	91	105
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	71	28	41	55	66
STANDARD DEVIATION.....GM.....	49	30	33	34	47
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	67	83	36	47	52
STANDARD DEVIATION.....GM.....	45	37	22	24	31
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	68	46	40	83	61
STANDARD DEVIATION.....GM.....	43	15	34	32	23
MAXIMUM IN ANY 1 DAY.....GM.....	630	178	160	164	320
MAXIMUM OVER 3 DAYS.....GM.....	788	232	261	473	320
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	69	20	40	55	62
STANDARD DEVIATION.....GM.....	48	18	32	32	42
PERCENTILES: <u>1/</u>					
5TH.....GM.....	13		7	13	20
25TH.....GM.....	39		20	38	39
50TH.....GM.....	62		38	40	59
75TH.....GM.....	80		59	79	79
90TH.....GM.....	158		80	82	104
95TH.....GM.....	160		82	104	158
99TH.....GM.....	236		158	160	197
MAXIMUM.....GM.....	630	82	160	160	320
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	282.2	8.8	6.6	23.3	33.0
PROPORTION OF EATING OCCASIONS.....PCT.....	5.7	13.5	3.8	6.2	8.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
18.4	14.6	14.6	12.4	10.5	9.5	10.1	6.7	7.8	6.4	7.8	5.9
14.6	12.4	11.9	10.6	8.2	8.6	8.3	5.9	5.6	5.0	6.0	5.1
2.9	1.7	2.0	1.5	1.3	.8	1.4	.6	1.5	1.1	.7	.7
.8	.4	.7	.2	.4	.1	.4	.1	.7	.4	1.1	.1
1.3	1.2	1.2	1.2	1.2	1.1	1.2	1.1	1.4	1.3	1.4	1.1
.6	.5	.5	.4	.5	.3	.5	.4	.8	.6	.7	.4
4.0	4.0	3.0	3.0	3.0	3.0	5.0	3.0	5.0	3.0	3.0	3.0
34	26	39	28	36	23	32	24	30	22	24	23
31	21	32	21	26	16	23	16	21	16	24	12
7	5	7	7	7	7	9	7	7	5		7
13	13	14	13	20	13	20	13	14	10		13
26	26	27	26	27	20	26	21	27	20		26
40	27	53	27	53	27	35	27	35	27		27
66	53	82	53	55	53	55	53	53	53		40
105	55	106	59	89	53	66	53	60	55		53
160	108	158	107	133	70	133	79	107	66		53
83	66	89	68	91	67	81	67	74	59	58	63
64	43	61	40	59	43	45	43	44	38	64	33
78	68	111	85	83	53	70	62	72	44	42	59
58	39	61	51	42	36	41	33	43	32	22	13
83	71	102	50	93	26	78	36	38	35	45	35
58	51	22	9	39	16	48	25	13	19	33	0
630	315	320	315	473	315	320	223	164	164	320	158
788	513	480	473	492	320	492	320	320	197	349	158
80	67	95	70	89	64	78	65	64	52	52	61
61	43	61	44	55	42	46	41	45	38	55	31
20	14	20	20	20	14	20	13	13	7		13
39	39	59	39	41	39	40	39	38	20		39
79	60	80	62	79	59	79	60	41	40		62
82	80	122	80	118	80	82	80	82	80		80
160	140	164	158	160	120	160	158	158	80		82
164	160	205	160	164	158	164	160	160	158		82
315	164	315	164	320	164	236	164	164	164		158
630	315	320	315	473	315	320	164	164	164	320	158
18.4	25.3	17.0	7.3	31.3	38.2	30.5	33.6	1.0	6.0	0	1.9
3.8	6.8	6.7	3.4	6.3	6.8	5.0	6.3	.8	4.2	0	2.8

TABLE 9.03.--HONEY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	AGE (YEARS) AND SEX OF INDIVIDUAL				
	TOTAL	UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.2	3.7	5.1	4.6	3.2
ON ONLY 1 OF 3 DAYS.....PCT.....	2.2	1.3	2.8	2.9	2.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.9	1.0	1.3	1.3	.3
ON ALL 3 OF 3 DAYS.....PCT.....	1.0	1.4	1.0	.4	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	2.0	3.1	1.8	1.6	1.4
STANDARD DEVIATION.....DO.....	1.3	2.3	1.0	.9	.8
MAXIMUM.....DO.....	14.0	9.0	6.0	6.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	14	32	9	12	11
STANDARD DEVIATION.....GM.....	19	96	11	17	11
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	5		4	5	5
50TH.....GM.....	7		7	7	7
75TH.....GM.....	16		11	14	18
90TH.....GM.....	28		18	21	28
95TH.....GM.....	42		21	59	35
99TH.....GM.....	85		78	85	57
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	27	12	17	25	30
STANDARD DEVIATION.....GM.....	31	7	13	39	30
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	23	20	12	19	18
STANDARD DEVIATION.....GM.....	22	8	8	19	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	25	70	18	32	22
STANDARD DEVIATION.....GM.....	30	147	20	32	13
MAXIMUM IN ANY 1 DAY.....GM.....	678	678	170	254	170
MAXIMUM OVER 3 DAYS.....GM.....	1,314	1,314	233	254	170
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	22	30	15	22	24
STANDARD DEVIATION.....GM.....	25	51	18	31	24
PERCENTILES: <u>1/</u>					
5TH.....GM.....	5		4	7	4
25TH.....GM.....	7		7	7	7
50TH.....GM.....	14		11	14	21
75TH.....GM.....	21		21	21	21
90TH.....GM.....	42		21	42	63
95TH.....GM.....	64		32	85	64
99TH.....GM.....	127		170	170	170
MAXIMUM.....GM.....	339	254	170	254	170
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	171.1	4.5	5.2	7.4	13.5
PROPORTION OF EATING OCCASIONS.....PCT.....	5.5	7.7	5.4	5.7	16.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.7	2.1	2.9	2.3	4.2	4.5	4.6	4.1	8.2	6.2	5.5	6.6
1.0	1.5	1.9	1.6	2.2	2.6	2.2	2.3	2.9	2.8	2.1	2.2
.5	.5	.6	.3	1.1	.9	.9	.9	2.3	1.7	.5	1.7
.1	.1	.4	.5	.9	.9	1.6	.9	3.0	1.8	2.9	2.7
1.6	1.5	1.8	1.7	1.9	1.9	2.3	1.9	2.3	2.1	2.5	2.2
1.0	.7	1.2	1.2	1.0	1.4	1.8	1.3	1.4	1.2	1.2	1.0
5.0	3.0	5.0	6.0	5.0	9.0	14.0	14.0	9.0	7.0	5.0	5.0
22	16	23	13	19	11	18	11	17	10	17	11
29	16	27	11	19	10	20	10	19	8	23	17
				2	2	2	2	2	2		2
				7	5	7	5	7	5		5
				12	7	12	7	12	7		7
				28	14	21	14	21	12		14
				42	24	42	21	42	21		16
				57	32	49	28	71	28		21
				89	49	127	42	85	33		127
63	36	41	25	37	22	32	21	17	18	15	14
101	31	33	22	40	19	27	18	12	11	12	7
32	39	36	25	33	19	23	24	23	14	7	26
26	37	18	19	28	13	19	15	17	9	5	46
37	21	68	26	31	21	30	18	30	16	28	12
25	0	45	10	22	14	27	11	24	10	27	6
339	254	191	85	205	85	254	148	127	42	127	191
339	283	424	125	268	205	381	205	254	99	254	381
41	32	38	23	30	18	24	18	23	14	21	15
67	34	32	16	31	15	25	15	21	10	20	24
				7	7	7	4	7	4		4
				11	7	7	7	7	7		7
				21	14	21	14	14	7		7
				42	21	32	21	21	21		21
				64	42	42	42	42	21		21
				85	42	85	42	64	42		21
				148	85	129	85	85	42		191
339	254	191	85	170	85	191	148	127	42	85	191
2.8	2.5	5.7	5.7	16.8	13.1	41.4	28.3	2.8	10.8	1.7	9.0
5.0	3.7	7.6	9.8	5.5	2.9	8.0	5.2	1.3	4.8	2.4	6.1

TABLE 9.04.--JELLIES

CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	17.0	.7	20.7	28.7	27.8
ON ONLY 1 OF 3 DAYS.....PCT.....	10.8	.7	12.9	18.2	18.9
ON ONLY 2 OF 3 DAYS.....PCT.....	4.0	0	5.6	6.9	6.8
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	0	2.3	3.5	2.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.6	1.6	1.5
STANDARD DEVIATION.....DO.....	.9	.0	.9	.9	.9
MAXIMUM.....DO.....	8.0	1.0	5.0	6.0	8.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	10	5	7	9	9
STANDARD DEVIATION.....GM.....	11	2	7	8	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	4		2	3	4
50TH.....GM.....	6		4	6	6
75TH.....GM.....	13		8	13	13
90TH.....GM.....	25		15	19	19
95TH.....GM.....	31		19	25	25
99TH.....GM.....	50		38	38	50
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	16	15	18	22
STANDARD DEVIATION.....GM.....	22	5	17	17	20
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	0	13	17	18
STANDARD DEVIATION.....GM.....	16	0	9	13	13
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	20	0	13	17	21
STANDARD DEVIATION.....GM.....	16	0	8	12	19
MAXIMUM IN ANY 1 DAY.....GM.....	300	19	114	156	150
MAXIMUM OVER 3 DAYS.....GM.....	375	19	114	250	227
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	20	16	13	16	19
STANDARD DEVIATION.....GM.....	18	5	13	14	17
PERCENTILES: <u>1/</u>					
5TH.....GM.....	6		3	3	6
25TH.....GM.....	6		6	6	6
50TH.....GM.....	19		6	13	19
75TH.....GM.....	19		19	19	19
90TH.....GM.....	38		19	38	38
95TH.....GM.....	56		38	38	38
99TH.....GM.....	75		75	75	75
MAXIMUM.....GM.....	300	19	114	150	150
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	358.3	0	11.5	30.7	31.9
PROPORTION OF EATING OCCASIONS.....PCT.....	3.6	0	3.3	3.9	4.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
26.1	21.3	19.6	15.1	13.8	13.6	15.2	13.3	19.8	15.1	15.5	17.5
18.3	14.6	12.2	10.7	8.5	9.9	9.1	8.4	10.8	8.1	6.3	8.0
5.2	4.7	4.5	3.1	3.0	2.8	4.0	3.3	5.1	4.6	4.1	4.5
2.5	2.0	2.9	1.3	2.2	.9	2.1	1.7	3.9	2.4	5.1	5.0
1.5	1.5	1.6	1.5	1.6	1.4	1.6	1.6	1.7	1.7	2.1	1.9
.8	.7	.8	.8	.9	.7	.9	.8	.9	1.0	1.2	1.0
6.0	5.0	5.0	5.0	7.0	5.0	6.0	5.0	5.0	6.0	7.0	5.0
12	10	13	11	14	9	12	8	11	9	16	10
11	10	13	12	14	9	12	8	9	10	17	11
2	2	2	2	2	2	2	2	2	2	2	2
4	4	6	4	4	3	4	2	4	2	4	4
8	6	10	6	8	6	8	6	7	6	13	6
15	13	19	13	19	13	15	12	13	13	20	10
25	25	25	25	29	19	25	17	25	19	44	25
38	25	38	38	38	25	35	25	31	25	50	38
50	56	57	50	75	43	50	40	41	46	75	52
27	22	27	23	27	19	25	16	22	13	33	17
24	20	27	27	27	18	22	17	20	8	55	10
23	20	27	26	27	20	24	17	18	16	18	15
15	13	17	18	21	15	22	13	12	10	16	17
26	22	25	24	25	20	18	16	19	20	25	16
15	19	18	15	20	14	14	11	10	18	16	14
169	169	300	300	300	150	263	188	113	113	206	99
188	244	305	300	300	225	375	188	150	225	225	206
24	21	25	23	25	19	22	16	19	15	22	16
20	18	23	23	24	16	20	14	16	12	25	14
6	6	6	6	6	6	6	3	6	6	6	6
9	6	13	9	13	6	9	6	13	6	6	6
19	19	19	19	19	18	19	13	14	9	19	13
38	19	38	28	38	19	36	19	19	19	19	19
38	38	38	38	56	38	38	38	38	38	38	38
57	56	56	75	75	38	56	38	38	38	75	38
84	75	114	75	114	75	75	75	99	56	150	75
169	169	300	300	300	150	225	150	113	75	150	99
31.0	29.7	19.0	15.7	27.6	35.1	45.5	37.1	12.0	14.9	3.3	13.3
3.8	4.4	4.3	4.8	3.2	3.5	3.7	2.5	3.2	3.3	1.9	4.1

TABLE 9.05.--JAMS, PRESERVES, AND MARMALADES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	7.1	1.4	6.7	6.8	6.6
ON ONLY 1 OF 3 DAYS.....PCT.....	4.5	1.2	4.9	4.4	4.5
ON ONLY 2 OF 3 DAYS.....PCT.....	1.6	.2	1.1	1.7	1.4
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	.7	.7	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.3	1.5	1.6	1.6
STANDARD DEVIATION.....DO.....	.9	.4	.8	.8	.9
MAXIMUM.....DO.....	6.0	2.0	4.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	12	5	6	9	10
STANDARD DEVIATION.....GM.....	14	4	5	6	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	2	2
25TH.....GM.....	4		2	4	4
50TH.....GM.....	7		7	7	7
75TH.....GM.....	13		7	13	13
90TH.....GM.....	27		13	20	20
95TH.....GM.....	33		13	20	33
99TH.....GM.....	67		27	27	53
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	25	13	16	20	23
STANDARD DEVIATION.....GM.....	27	13	15	14	21
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	22	13	14	18	21
STANDARD DEVIATION.....GM.....	18	0	9	11	15
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	24	0	9	14	20
STANDARD DEVIATION.....GM.....	26	0	3	6	14
MAXIMUM IN ANY 1 DAY.....GM.....	320	40	80	60	160
MAXIMUM OVER 3 DAYS.....GM.....	800	40	80	80	160
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	23	12	13	17	20
STANDARD DEVIATION.....GM.....	23	8	11	12	17
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		3	7	7
25TH.....GM.....	7		7	7	7
50TH.....GM.....	20		7	13	20
75TH.....GM.....	20		20	20	20
90TH.....GM.....	40		20	40	40
95TH.....GM.....	60		40	40	40
99TH.....GM.....	122		40	60	60
MAXIMUM.....GM.....	320	20	80	60	160
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	140.5	0	3.8	7.2	14.3
PROPORTION OF EATING OCCASIONS.....PCT.....	3.3	0	3.7	3.9	7.6

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.8	5.4	6.5	4.8	6.4	4.7	8.3	7.2	11.7	12.6	11.6	12.4
4.2	4.0	3.8	3.7	4.3	3.5	5.1	4.9	4.9	6.9	4.6	5.2
1.3	1.1	2.1	1.0	1.4	.9	1.9	1.4	3.4	3.2	4.2	3.7
.4	.4	.5	.1	.6	.3	1.2	.9	3.4	2.5	2.8	3.5
1.5	1.5	1.6	1.3	1.4	1.4	1.6	1.5	2.1	1.7	1.9	2.0
.8	.8	.8	.5	.7	.7	1.0	.8	1.2	.9	.9	1.0
4.0	4.0	5.0	3.0	3.0	5.0	6.0	6.0	6.0	4.0	5.0	5.0
14	11	17	9	14	9	14	11	20	10	19	9
13	11	13	9	13	8	14	13	35	8	22	7
2	2	4	2	2	2	2	2	2	2	1	2
7	4	7	3	5	4	5	4	7	4	7	4
11	7	13	7	13	7	8	7	13	7	11	7
17	13	27	11	19	13	19	13	20	13	20	13
27	22	33	16	27	13	27	20	40	20	53	20
40	40	53	20	33	20	39	33	53	27	67	22
67	53	67	53	80	47	67	67	267	47	93	31
30	25	40	22	30	23	29	25	21	20	32	14
25	20	33	29	22	20	30	37	15	15	43	9
27	25	35	18	25	17	23	19	23	16	40	14
12	20	23	8	15	10	20	20	14	9	39	7
45	27	23	20	36	18	27	19	42	17	21	16
22	19	9	0	24	13	22	12	58	13	20	7
142	120	160	160	120	160	200	320	320	80	240	40
240	180	200	160	240	160	346	320	800	142	280	93
28	23	32	20	29	19	25	21	29	17	30	14
20	16	26	24	21	17	24	29	30	13	39	9
7	7	7	7	7	7	7	3	7	7	7	7
13	10	20	7	13	7	13	7	10	7	7	7
20	20	20	20	20	20	20	13	20	13	20	13
40	40	40	20	40	20	30	20	40	20	40	20
40	40	80	40	60	40	40	40	60	40	40	20
80	60	80	40	80	40	80	40	80	40	80	40
120	80	160	160	80	80	160	160	160	60	240	40
122	80	160	160	120	160	160	320	160	80	240	40
1.0	6.3	4.1	7.5	13.5	5.9	37.3	15.0	12.3	5.6	.7	6.0
.6	3.7	2.8	8.4	3.7	1.7	5.6	2.0	4.6	1.5	.6	2.4

TABLE 9.06.--PLAIN GELATIN DESSERTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.6	3.8	7.3	7.9	11.5
ON ONLY 1 OF 3 DAYS.....PCT.....	5.4	2.8	5.6	6.4	9.6
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	.9	1.2	1.2	1.6
ON ALL 3 OF 3 DAYS.....PCT.....	.2	.2	.5	.4	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.7	1.4	1.3	1.2
STANDARD DEVIATION.....DO.....	.6	.9	.8	.6	.5
MAXIMUM.....DO.....	6.0	3.0	5.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	67	50	55	61	62
STANDARD DEVIATION.....GM.....	48	44	47	40	39
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20		15	19	20
25TH.....GM.....	40		30	40	40
50TH.....GM.....	51		40	40	40
75TH.....GM.....	80		66	80	80
90TH.....GM.....	120		80	120	100
95TH.....GM.....	160		160	120	160
99TH.....GM.....	240		260	200	200
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	164	137	122	146	157
STANDARD DEVIATION.....GM.....	93	143	72	84	83
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	171	110	132	153	165
STANDARD DEVIATION.....GM.....	82	16	74	50	83
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	170	3	132	146	118
STANDARD DEVIATION.....GM.....	96	0	89	44	31
MAXIMUM IN ANY 1 DAY.....GM.....	960	540	480	600	816
MAXIMUM OVER 3 DAYS.....GM.....	1,440	540	780	720	952
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	159	89	118	144	151
STANDARD DEVIATION.....GM.....	85	62	73	74	78
PERCENTILES: <u>1/</u>					
5TH.....GM.....	60		29	60	60
25TH.....GM.....	120		60	120	120
50TH.....GM.....	120		120	120	120
75TH.....GM.....	240		120	180	240
90TH.....GM.....	240		240	240	240
95TH.....GM.....	240		240	240	240
99TH.....GM.....	480		360	360	480
MAXIMUM.....GM.....	960	240	480	480	480
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	29.5	0	.7	.7	3.3
PROPORTION OF EATING OCCASIONS.....PCT.....	.9	0	.6	.4	1.3

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.5	9.9	7.7	5.5	3.6	4.0	5.7	5.4	8.6	10.8	7.9	9.0
9.1	8.0	6.5	4.7	3.1	3.4	4.5	4.7	5.8	8.5	4.9	6.7
1.4	1.9	1.0	.7	.5	.5	.9	.6	1.9	1.9	1.0	1.9
(2)	.1	.2	.1	0	.1	.2	.1	.9	.4	1.9	.5
1.2	1.3	1.3	1.3	1.2	1.2	1.3	1.2	1.5	1.3	1.7	1.4
.4	.6	.5	.7	.4	.5	.8	.5	.9	.5	.8	.7
3.0	4.0	3.0	6.0	3.0	4.0	6.0	6.0	6.0	3.0	3.0	4.0
65	66	78	76	70	62	80	64	74	61	91	76
46	47	56	63	41	40	61	45	41	41	69	55
20	20	20	20	20	15	20	20	23	10		23
40	40	40	40	40	40	40	40	40	40		40
40	51	80	60	80	40	80	40	77	40		60
80	80	80	80	80	80	80	80	80	80		80
100	120	160	160	120	100	160	110	160	120		160
160	160	183	160	160	160	160	160	160	160		200
306	240	306	440	240	200	360	240	160	240		320
170	162	197	186	180	159	180	161	174	141	154	167
117	92	126	108	96	79	92	84	87	72	113	77
172	163	188	181	194	168	210	153	167	158	183	193
86	95	109	80	64	71	87	67	66	64	49	108
120	240	224	340	0	156	221	244	96	141	174	150
0	0	26	100	0	98	127	81	38	59	60	80
960	480	720	600	720	480	960	600	480	240	480	480
960	960	918	1,320	720	960	1,440	1,200	480	720	720	960
165	155	186	181	180	157	182	157	144	144	164	166
109	79	101	87	90	77	94	74	71	72	84	83
46	60	60	60	60	45	60	60	51	30		60
120	120	120	120	120	120	120	120	91	91		120
120	120	153	180	180	120	161	120	120	120		120
240	240	240	240	240	240	240	240	240	240		240
240	240	240	240	240	240	240	240	240	240		240
240	240	459	360	240	240	240	240	240	240		240
720	360	480	480	720	480	480	240	240	240		480
960	480	480	480	720	480	960	480	240	240	480	480
2.8	1.9	1.8	0	0	8.0	1.5	4.3	0	0	.9	3.6
1.1	.7	1.3	0	0	3.2	.4	.9	0	0	1.3	2.9

TABLE 9.07.--GELATIN DESSERTS WITH FRUIT
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.3	.4	3.4	2.4	4.3
ON ONLY 1 OF 3 DAYS.....PCT.....	2.9	.2	3.2	2.1	4.1
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	.2	.1	.2	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.5	1.1	1.2	1.1
STANDARD DEVIATION.....DO.....	.5	.5	.4	.4	.2
MAXIMUM.....DO.....	5.0	2.0	3.0	3.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	67	20	49	52	55
STANDARD DEVIATION.....GM.....	49	20	42	33	31
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	20				20
25TH.....GM.....	40				40
50TH.....GM.....	40				40
75TH.....GM.....	80				80
90TH.....GM.....	133				80
95TH.....GM.....	160				120
99TH.....GM.....	240				160
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	170	2	128	142	158
STANDARD DEVIATION.....GM.....	101	0	66	89	88
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	191	60	37	136	172
STANDARD DEVIATION.....GM.....	85	0	0	54	37
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	235	0	240	0	0
STANDARD DEVIATION.....GM.....	83	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	720	60	360	480	480
MAXIMUM OVER 3 DAYS.....GM.....	1,200	120	720	480	480
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	167	40	128	135	156
STANDARD DEVIATION.....GM.....	90	27	81	84	82
PERCENTILES: <u>1/</u>					
5TH.....GM.....	60				60
25TH.....GM.....	120				120
50TH.....GM.....	120				120
75TH.....GM.....	240				240
90TH.....GM.....	240				240
95TH.....GM.....	306				240
99TH.....GM.....	480				480
MAXIMUM.....GM.....	612	60	360	480	480
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	22.1	0	.8	1.2	3.0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	2.1	2.6	3.6

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.4	3.5	4.2	3.0	2.0	1.7	3.0	3.8	4.5	5.0	4.4	6.0
5.2	3.2	3.8	2.9	1.7	1.5	2.6	3.1	3.7	3.7	3.7	5.1
.2	.2	.4	.1	.3	.1	.3	.6	.7	1.3	.4	.9
0	.1	.1	0	0	0	.1	.1	.1	.1	.2	0
1.1	1.1	1.1	1.1	1.2	1.1	1.2	1.3	1.3	1.4	1.4	1.3
.3	.4	.4	.3	.5	.3	.5	.6	.5	.6	.9	.6
2.0	3.0	3.0	2.0	3.0	3.0	4.0	5.0	3.0	3.0	5.0	3.0
59	65	69	63	72	57	75	69	78	86	78	65
34	46	45	37	38	32	52	55	46	70	74	47
20	20	20		30	20	23	15	26	20		11
40	40	40		40	40	40	40	40	40		40
40	40	53		80	40	60	40	80	80		40
80	80	80		80	80	80	80	94	120		80
91	160	160		140	120	160	160	160	160		120
140	160	160		160	120	160	180	160	180		200
160	240	240		160	160	240	240	160	360		204
170	177	182	185	204	165	185	159	187	169	179	157
93	119	104	109	110	97	104	92	102	122	83	106
189	156	198	180	143	132	211	184	236	253	150	198
46	51	79	0	58	21	60	90	10	106	22	73
0	190	220	0	0	0	266	242	120	160	400	0
0	53	0	0	0	0	43	86	0	0	0	0
480	720	480	544	480	480	600	720	480	720	480	612
480	720	720	544	480	480	960	1,200	480	1,080	1,200	612
166	172	186	177	175	157	185	161	182	183	168	149
80	103	101	100	90	88	86	84	80	101	73	91
60	60	60		60	60	68	60	79	60		17
120	120	120		120	120	120	120	120	120		120
120	120	153		120	120	180	120	180	120		120
240	240	240		240	240	240	240	240	240		180
240	240	272		240	240	240	240	240	360		240
240	480	480		360	360	272	240	240	360		240
480	480	480		480	480	480	480	480	360		612
480	480	480	544	480	480	480	480	480	612	240	612
3.0	2.3	0	.8	1.3	2.1	4.0	2.3	0	1.3	0	0
2.4	2.7	0	1.8	1.3	2.1	2.2	.7	0	1.1	0	0

TABLE 9.08.--CANDY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	13.0	.8	17.4	20.9	19.9
ON ONLY 1 OF 3 DAYS.....PCT.....	9.4	.8	11.3	15.8	14.7
ON ONLY 2 OF 3 DAYS.....PCT.....	2.8	0	4.6	4.1	3.9
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	1.5	1.0	1.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	1.0	1.6	1.4	1.5
STANDARD DEVIATION.....DO.....	1.0	.0	.9	.9	.9
MAXIMUM.....DO.....	15.0	1.0	6.0	9.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	22	6	15	16	19
STANDARD DEVIATION.....GM.....	24	4	18	15	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3		2	2	2
25TH.....GM.....	9		5	6	7
50TH.....GM.....	14		9	11	13
75TH.....GM.....	28		19	20	24
90TH.....GM.....	47		35	35	43
95TH.....GM.....	66		40	47	58
99TH.....GM.....	116		122	73	102
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	47	18	29	37	41
STANDARD DEVIATION.....GM.....	46	13	27	31	40
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	51	0	33	38	50
STANDARD DEVIATION.....GM.....	45	0	33	27	41
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	50	0	33	36	42
STANDARD DEVIATION.....GM.....	38	0	30	20	26
MAXIMUM IN ANY 1 DAY.....GM.....	907	37	240	227	340
MAXIMUM OVER 3 DAYS.....GM.....	1,105	37	421	340	495
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	43	18	29	33	38
STANDARD DEVIATION.....GM.....	42	13	31	29	34
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		5	6	7
25TH.....GM.....	18		12	14	15
50TH.....GM.....	30		17	28	30
75TH.....GM.....	51		30	43	48
90TH.....GM.....	85		57	60	74
95TH.....GM.....	113		85	85	113
99TH.....GM.....	220		156	170	170
MAXIMUM.....GM.....	907	37	240	227	227
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	311.6	0	8.9	22.9	25.1
PROPORTION OF EATING OCCASIONS.....PCT.....	4.1	0	3.1	4.4	4.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
22.0	23.2	16.9	20.0	11.5	11.7	8.2	9.8	8.1	8.7	6.5	8.1
16.1	15.7	11.8	14.0	7.9	9.0	5.6	7.4	5.9	5.7	4.5	5.9
4.4	5.8	3.5	5.0	2.7	2.2	1.9	1.8	1.3	2.3	1.4	1.7
1.5	1.7	1.6	.9	.9	.5	.7	.6	.9	.7	.6	.5
1.5	1.6	1.6	1.5	1.6	1.4	1.5	1.4	1.9	1.6	1.8	1.4
.9	1.0	1.0	.9	1.3	.9	.8	1.1	2.7	.8	1.8	.8
7.0	9.0	6.0	6.0	12.0	7.0	6.0	13.0	15.0	5.0	9.0	5.0
26	24	28	23	29	20	26	19	23	17	15	15
26	28	32	21	28	18	30	22	23	16	9	13
3	3	5	3	5	2	4	3	5	3		3
9	9	10	10	11	9	9	7	9	7		6
18	16	19	15	19	14	18	13	13	11		11
32	28	33	30	38	26	33	21	27	24		19
57	47	57	48	64	40	57	38	53	38		32
84	68	87	71	89	57	74	57	66	43		36
123	132	159	103	141	87	189	95	110	75		80
54	48	58	57	59	47	52	40	40	42	36	34
52	49	55	55	57	43	51	33	31	48	19	27
65	55	66	49	65	45	61	45	47	34	33	37
47	61	72	31	48	30	57	35	22	21	9	29
59	49	62	43	70	42	56	51	71	23	25	23
34	33	37	34	29	28	49	54	25	14	13	7
425	907	737	340	567	340	340	397	168	298	85	213
510	1,105	907	410	567	405	983	808	330	298	108	241
51	44	53	47	53	41	53	39	35	33	25	31
44	51	49	45	52	37	53	33	26	34	18	30
7	6	11	7	8	7	11	6	7	6		7
24	15	28	21	28	18	25	16	20	14		14
36	30	37	34	39	32	38	30	28	28		25
57	55	63	56	57	48	57	50	48	40		34
113	85	105	104	113	85	102	84	60	60		57
151	113	142	113	142	113	144	113	85	85		97
227	227	284	227	255	170	328	170	168	224		213
330	907	454	340	567	340	340	227	168	298	85	213
31.4	25.0	15.3	19.4	23.4	37.3	28.9	49.1	6.7	8.9	1.8	7.3
4.5	3.1	4.1	4.4	3.1	4.1	4.9	4.9	3.9	3.8	2.8	6.5

TABLE 10.01.--WHITE POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	74.4	18.1	74.5	76.3	80.7
ON ONLY 1 OF 3 DAYS.....PCT.....	34.2	8.7	34.7	37.2	37.3
ON ONLY 2 OF 3 DAYS.....PCT.....	28.4	4.9	30.4	29.4	31.8
ON ALL 3 OF 3 DAYS.....PCT.....	11.8	4.5	9.4	9.6	11.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.9	2.0	1.8	1.8	1.9
STANDARD DEVIATION.....DO.....	1.0	1.2	.9	.8	.9
MAXIMUM.....DO.....	10.0	6.0	6.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	78	47	43	51	62
STANDARD DEVIATION.....GM.....	65	51	42	41	47
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	14	4	6	9	10
25TH.....GM.....	33	14	18	21	28
50TH.....GM.....	62	32	35	39	50
75TH.....GM.....	103	58	55	70	85
90TH.....GM.....	158	132	92	101	125
95TH.....GM.....	202	176	112	130	152
99TH.....GM.....	309	253	168	200	216
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	130	55	69	90	105
STANDARD DEVIATION.....GM.....	98	49	54	69	79
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	135	83	77	93	113
STANDARD DEVIATION.....GM.....	81	61	46	57	66
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	151	95	90	96	112
STANDARD DEVIATION.....GM.....	86	63	78	51	49
MAXIMUM IN ANY 1 DAY.....GM.....	1,560	470	1,164	586	702
MAXIMUM OVER 3 DAYS.....GM.....	2,680	758	2,152	1,260	1,179
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	125	72	70	86	100
STANDARD DEVIATION.....GM.....	90	58	56	62	69
PERCENTILES: <u>1/</u>					
5TH.....GM.....	29	12	14	16	26
25TH.....GM.....	63	28	34	46	57
50TH.....GM.....	105	53	57	70	90
75TH.....GM.....	170	105	93	105	122
90TH.....GM.....	235	122	123	179	210
95TH.....GM.....	280	210	170	210	210
99TH.....GM.....	426	235	250	276	340
MAXIMUM.....GM.....	1,260	368	630	508	588
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	468.8	0	7.6	17.2	29.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.9	0	.5	.7	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
81.8	77.0	81.2	77.2	79.6	70.5	78.1	71.3	76.4	69.4	70.9	66.3
35.9	35.2	34.0	35.9	34.7	36.5	32.1	34.3	32.3	33.2	31.1	29.1
32.5	30.2	30.9	30.5	30.2	25.3	30.5	27.2	28.5	26.7	25.4	23.9
13.4	11.6	16.2	10.9	14.7	8.7	15.5	9.7	15.6	9.5	14.4	13.3
2.0	1.9	2.0	1.9	2.0	1.7	2.0	1.8	2.0	1.8	2.0	1.9
1.0	.9	1.1	1.0	1.0	.9	1.0	.9	1.0	.9	1.1	1.0
6.0	6.0	7.0	6.0	6.0	7.0	7.0	6.0	6.0	6.0	6.0	10.0
81	71	101	73	96	69	99	70	91	73	100	76
63	54	82	60	80	54	78	54	74	52	79	55
15	11	19	10	18	13	19	15	18	16	20	18
35	31	42	31	39	31	45	31	41	35	45	35
66	57	81	60	76	57	77	58	70	62	78	62
111	93	140	96	129	89	132	90	117	94	125	101
159	148	193	140	190	138	202	140	172	140	206	146
198	182	259	188	242	174	250	173	225	162	260	179
297	260	407	288	358	266	340	266	393	267	365	274
135	121	167	123	158	124	154	124	145	133	159	123
102	90	124	93	123	86	109	89	103	84	114	73
139	126	168	127	162	126	163	126	141	130	160	126
84	72	100	69	99	64	91	71	73	64	87	71
148	130	177	147	174	144	179	135	175	137	191	135
70	63	105	86	103	71	98	67	96	63	91	60
985	980	1,560	945	1,260	816	1,134	1,074	850	926	736	854
1,701	1,260	2,143	2,019	2,680	1,294	2,652	1,709	1,865	1,242	1,510	1,129
124	112	149	116	146	119	149	119	140	124	154	119
87	80	112	86	110	80	103	78	91	70	94	68
29	26	29	26	29	29	32	29	39	32	53	29
57	57	68	57	70	60	85	70	85	90	93	83
105	93	114	93	114	98	122	98	118	105	123	105
170	140	210	150	210	158	210	145	184	158	188	140
210	210	276	210	269	210	263	210	245	210	247	210
276	250	368	254	368	250	366	250	310	245	340	244
420	420	508	420	531	420	500	420	490	388	488	368
716	980	1,260	945	1,260	816	1,134	840	732	630	736	490
42.5	35.4	25.3	26.3	67.0	44.7	69.9	61.4	12.3	20.7	4.0	5.4
1.3	1.1	1.1	1.2	1.1	.7	.9	.7	.7	1.0	.5	.4

TABLE 10.02.--BAKED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	18.3	2.1	14.2	13.7	14.3
ON ONLY 1 OF 3 DAYS.....PCT.....	15.6	1.9	12.4	12.6	12.9
ON ONLY 2 OF 3 DAYS.....PCT.....	2.4	.3	1.7	1.0	1.2
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.1	.1	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.1	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.3	.4	.3	.3
MAXIMUM.....DO.....	5.0	2.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	40	22	24	27	31
STANDARD DEVIATION.....GM.....	22	18	15	15	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	16		6	12	16
25TH.....GM.....	31		16	16	23
50TH.....GM.....	31		23	23	31
75TH.....GM.....	47		31	31	31
90TH.....GM.....	62		43	47	47
95TH.....GM.....	93		62	62	61
99TH.....GM.....	124		78	92	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	103	49	62	76	84
STANDARD DEVIATION.....GM.....	43	30	36	37	28
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	105	93	67	72	87
STANDARD DEVIATION.....GM.....	36	0	27	27	34
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	104	0	35	66	100
STANDARD DEVIATION.....GM.....	29	0	0	15	38
MAXIMUM IN ANY 1 DAY.....GM.....	784	98	255	276	280
MAXIMUM OVER 3 DAYS.....GM.....	882	186	255	279	420
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	103	59	62	75	85
STANDARD DEVIATION.....GM.....	42	32	31	36	30
PERCENTILES: <u>1/</u>					
5TH.....GM.....	47		18	35	47
25TH.....GM.....	93		47	47	70
50TH.....GM.....	93		53	70	93
75TH.....GM.....	105		70	93	93
90TH.....GM.....	140		93	98	140
95TH.....GM.....	184		113	140	140
99TH.....GM.....	280		186	210	184
MAXIMUM.....GM.....	784	98	227	276	280
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	10.2	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.1	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
15.6	15.4	14.1	16.3	19.1	17.8	23.4	21.1	21.4	19.1	16.0	16.0
14.4	14.0	12.6	14.5	15.9	15.8	19.5	17.8	16.9	15.3	12.8	11.8
1.1	1.3	1.4	1.6	3.0	1.7	3.4	3.0	3.8	3.3	3.0	3.6
.1	.1	.1	.2	.3	.3	.5	.3	.7	.6	.2	.6
1.1	1.1	1.1	1.1	1.2	1.1	1.2	1.2	1.3	1.2	1.2	1.3
.3	.3	.3	.4	.4	.4	.5	.4	.6	.5	.5	.5
3.0	4.0	3.0	3.0	4.0	4.0	4.0	5.0	4.0	5.0	3.0	3.0
38	36	44	36	48	38	46	39	43	41	44	39
17	22	23	16	27	18	25	20	21	21	22	20
23	16	23	23	23	18	23	23	16	18	23	16
31	31	31	31	31	31	31	31	31	31	31	31
31	31	31	31	35	31	31	31	31	31	31	31
47	35	47	38	62	47	48	47	48	47	47	50
62	54	70	62	93	62	78	62	70	62	93	62
65	62	93	64	97	72	93	78	93	82	93	70
98	98	155	93	140	109	140	117	109	117	109	109
104	97	120	97	120	101	115	99	108	99	103	89
38	38	56	34	53	36	50	36	43	36	23	22
101	103	112	97	124	103	116	101	96	99	121	90
28	71	51	16	40	22	36	32	26	23	26	14
112	105	109	93	133	110	103	98	98	103	93	110
16	4	0	0	46	25	29	21	9	23	0	17
350	784	465	280	454	284	560	465	284	350	186	151
377	882	558	297	746	558	658	558	336	465	326	420
104	97	117	96	121	101	114	99	102	99	107	91
38	50	55	30	51	33	48	36	39	33	24	23
70	47	70	70	70	53	70	53	47	53	93	49
93	75	93	93	93	93	93	93	93	93	93	70
93	93	93	93	93	93	93	93	93	93	93	93
98	93	140	93	140	98	140	98	140	93	140	93
140	140	186	140	186	140	145	140	140	140	140	140
186	140	209	140	210	147	186	140	145	145	140	140
279	280	290	227	280	210	280	280	279	227	186	147
350	784	465	280	454	284	560	465	284	350	186	151
0	1.2	0	0	3.5	1.4	2.9	0	0	0	0	1.2
0	.3	0	0	.4	.1	.2	0	0	0	0	.6

TABLE 10.03.--BOILED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	12.6	3.8	9.9	10.3	11.9
ON ONLY 1 OF 3 DAYS.....PCT.....	10.5	2.9	8.7	9.1	10.6
ON ONLY 2 OF 3 DAYS.....PCT.....	1.9	.9	1.0	1.2	1.3
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.3	1.2	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.5	.5	.3	.4
MAXIMUM.....DO.....	6.0	3.0	3.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	63	37	35	42	47
STANDARD DEVIATION.....GM.....	49	31	53	27	31
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	15		7	7	15
25TH.....GM.....	31		15	30	30
50TH.....GM.....	47		30	41	41
75TH.....GM.....	81		35	60	60
90TH.....GM.....	122		60	77	81
95TH.....GM.....	153		61	85	102
99TH.....GM.....	245		328	129	169
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	156	95	81	113	123
STANDARD DEVIATION.....GM.....	104	95	46	69	71
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	160	83	165	109	141
STANDARD DEVIATION.....GM.....	94	29	203	47	56
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	177	0	11	0	0
STANDARD DEVIATION.....GM.....	93	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,164	368	1,164	508	508
MAXIMUM OVER 3 DAYS.....GM.....	1,701	368	1,522	508	627
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	153	88	87	112	124
STANDARD DEVIATION.....GM.....	94	77	81	66	67
PERCENTILES: <u>1/</u>					
5TH.....GM.....	45		11	22	45
25TH.....GM.....	92		45	61	90
50TH.....GM.....	122		90	92	122
75TH.....GM.....	184		92	127	179
90TH.....GM.....	254		127	184	184
95TH.....GM.....	358		184	194	254
99TH.....GM.....	488		537	276	358
MAXIMUM.....GM.....	920	368	627	508	508
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	31.0	0	.8	2.6	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.6	1.3	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
10.5	10.5	11.0	11.3	9.1	9.2	14.0	14.6	19.4	20.0	25.1	23.0
8.9	9.0	9.3	10.0	8.1	8.2	11.1	12.2	15.4	15.4	16.7	15.3
1.6	1.4	1.4	1.3	.9	1.0	2.4	2.2	3.3	4.0	6.8	5.1
0	.1	.2	0	.1	(2)	.5	.2	.7	.6	1.6	2.6
1.2	1.2	1.2	1.1	1.1	1.1	1.3	1.2	1.3	1.4	1.5	1.5
.4	.4	.5	.3	.4	.3	.6	.5	.7	.6	.8	.8
3.0	3.0	4.0	2.0	3.0	3.0	6.0	5.0	5.0	3.0	6.0	4.0
62	58	79	57	77	57	79	54	78	61	80	68
50	40	61	42	49	35	64	37	74	38	62	48
15	10	9	15	30	15	22	15	30	23	20	30
31	30	41	31	42	31	41	31	41	37	41	41
60	42	61	42	61	42	60	41	60	46	60	49
81	71	90	61	90	81	92	61	81	81	111	90
121	119	149	119	143	119	163	100	123	119	163	123
142	153	203	123	163	123	209	122	245	123	207	143
215	179	298	259	244	171	345	184	469	203	325	285
155	148	200	151	211	155	184	140	164	154	159	146
99	102	136	93	124	91	123	93	92	83	109	91
176	155	207	160	177	150	205	137	188	122	185	151
121	55	116	102	83	55	127	61	98	53	98	80
0	152	257	0	276	171	208	121	340	162	170	122
0	42	25	0	103	0	74	43	118	65	68	42
985	582	895	776	732	552	1,089	1,074	732	488	736	854
1,701	582	1,196	1,074	1,074	644	1,611	1,074	1,406	988	1,094	976
157	146	198	150	202	152	185	135	176	135	157	133
97	87	112	97	112	84	113	77	104	66	92	66
45	45	45	46	90	45	61	45	90	46	61	47
92	90	122	90	122	92	113	90	93	92	95	92
122	122	179	122	179	122	179	122	179	122	127	122
184	183	244	184	244	184	244	179	184	179	184	142
269	244	358	244	358	254	358	194	358	194	244	184
366	358	366	358	448	358	368	276	368	269	269	244
488	488	626	538	552	448	553	460	552	368	488	368
716	582	776	776	732	552	920	537	732	426	736	488
1.1	4.0	1.6	1.7	3.1	.8	3.6	10.4	1.2	0	0	0
.4	1.5	.9	.9	.8	.1	.4	.8	.4	0	0	0

TABLE 10.04.--CREAMED, SCALLOPED, AND AU GRATIN POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.4	2.1	5.6	4.7	5.1
ON ONLY 1 OF 3 DAYS.....PCT.....	4.8	1.6	4.8	4.1	4.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.6	.1	.7	.4	.6
ON ALL 3 OF 3 DAYS.....PCT.....	.1	.3	.1	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.6	1.2	1.1	1.2
STANDARD DEVIATION.....DO.....	.4	1.3	.5	.4	.4
MAXIMUM.....DO.....	5.0	5.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	69	44	35	46	55
STANDARD DEVIATION.....GM.....	46	75	33	31	37
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	10		5	5	5
25TH.....GM.....	39		10	20	26
50TH.....GM.....	61		20	41	41
75TH.....GM.....	82		41	78	78
90TH.....GM.....	124		82	82	82
95TH.....GM.....	163		98	82	118
99TH.....GM.....	235		157	163	196
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	189	32	80	127	152
STANDARD DEVIATION.....GM.....	121	32	59	90	107
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	179	29	120	113	125
STANDARD DEVIATION.....GM.....	85	0	74	45	51
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	113	209	98	52	68
STANDARD DEVIATION.....GM.....	77	29	0	7	10
MAXIMUM IN ANY 1 DAY.....GM.....	980	470	245	490	588
MAXIMUM OVER 3 DAYS.....GM.....	980	705	470	490	588
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	179	80	87	119	141
STANDARD DEVIATION.....GM.....	112	70	65	84	98
PERCENTILES: <u>1/</u>					
5TH.....GM.....	29		15	15	15
25TH.....GM.....	118		30	61	61
50TH.....GM.....	123		61	118	118
75TH.....GM.....	245		118	123	235
90TH.....GM.....	245		235	235	245
95TH.....GM.....	372		235	245	245
99TH.....GM.....	490		245	353	587
MAXIMUM.....GM.....	980	235	245	490	588
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	35.4	0	.7	2.0	3.2
PROPORTION OF EATING OCCASIONS.....PCT.....	1.5	0	.9	2.1	2.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.0	5.2	6.9	4.1	5.5	4.4	5.7	5.4	8.6	5.9	5.2	6.4
5.7	4.3	6.0	3.5	5.2	3.9	5.0	4.7	8.2	5.3	5.0	5.5
.2	.9	1.0	.5	.3	.5	.7	.6	.4	.7	0	.4
.1	0	0	.1	(2)	(2)	0	(2)	.1	0	.2	.4
1.1	1.2	1.2	1.2	1.1	1.1	1.2	1.2	1.1	1.2	1.2	1.3
.4	.4	.4	.5	.3	.4	.4	.4	.3	.5	.5	.6
4.0	2.0	3.0	3.0	3.0	3.0	3.0	4.0	3.0	3.0	3.0	3.0
63	67	89	60	87	69	80	67	68	63	79	62
48	49	52	54	54	41	48	42	43	36	38	39
15	18	20	9	20	14	20	15	10	20		5
39	39	55	39	41	39	41	39	39	39		39
41	44	78	41	78	78	78	59	61	59		64
82	82	118	78	123	82	82	82	82	82		82
82	127	163	123	163	123	157	123	141	102		118
163	163	163	137	201	163	163	157	163	123		163
313	245	235	313	245	163	245	235	186	212		163
169	176	236	135	254	184	222	180	195	165	230	184
93	122	117	75	159	103	130	109	122	72	114	106
228	158	233	216	199	192	191	173	192	192	0	147
139	93	101	126	74	56	77	75	48	71	0	107
296	0	0	196	93	54	0	137	117	0	117	44
19	0	0	0	0	0	0	0	0	0	0	16
588	980	790	470	735	490	735	705	558	470	490	490
940	980	875	940	735	588	735	735	558	637	490	490
174	169	228	154	240	182	209	171	183	161	194	143
98	115	119	91	142	95	119	96	107	76	97	106
31	29	59	28	61	41	59	44	29	29		15
118	118	123	118	123	118	118	118	118	118		44
123	123	235	123	235	235	235	123	123	123		118
235	245	245	235	245	245	245	235	235	235		235
245	245	369	245	490	245	368	245	306	245		245
368	353	470	245	490	368	490	368	372	245		353
490	490	490	470	706	490	490	470	558	470		490
490	980	705	470	735	490	735	490	558	470	490	490
3.0	3.6	2.8	1.5	1.0	7.7	2.0	2.6	3.6	.8	.9	0
2.2	2.7	2.5	2.2	.4	2.9	.6	.6	3.3	.7	2.7	0

TABLE 10.05.--MASHED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	26.5	11.5	29.0	30.1	32.8
ON ONLY 1 OF 3 DAYS.....PCT.....	21.4	8.0	23.0	24.9	25.9
ON ONLY 2 OF 3 DAYS.....PCT.....	4.5	2.1	5.1	4.4	6.1
ON ALL 3 OF 3 DAYS.....PCT.....	.6	1.4	.9	.8	.9
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.6	1.3	1.2	1.3
STANDARD DEVIATION.....DO.....	.5	1.0	.6	.5	.5
MAXIMUM.....DO.....	9.0	6.0	4.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	67	41	37	46	56
STANDARD DEVIATION.....GM.....	51	42	29	38	40
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	18	4	7	8	13
25TH.....GM.....	35	18	18	18	35
50TH.....GM.....	70	35	35	35	35
75TH.....GM.....	70	48	41	70	70
90TH.....GM.....	140	70	70	70	105
95TH.....GM.....	140	144	96	123	140
99TH.....GM.....	280	210	140	175	210
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	163	73	91	112	133
STANDARD DEVIATION.....GM.....	100	55	69	75	80
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	167	98	86	112	135
STANDARD DEVIATION.....GM.....	96	66	46	59	66
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	176	100	96	150	152
STANDARD DEVIATION.....GM.....	104	61	49	91	63
MAXIMUM IN ANY 1 DAY.....GM.....	1,260	368	630	420	525
MAXIMUM OVER 3 DAYS.....GM.....	2,310	630	630	1,260	840
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	160	79	87	112	130
STANDARD DEVIATION.....GM.....	98	52	62	70	74
PERCENTILES: <u>1/</u>					
5TH.....GM.....	39	25	13	25	26
25TH.....GM.....	105	39	53	53	105
50TH.....GM.....	105	53	92	105	105
75TH.....GM.....	210	105	105	146	210
90TH.....GM.....	210	151	158	210	210
95TH.....GM.....	368	210	210	210	210
99TH.....GM.....	420	221	210	368	420
MAXIMUM.....GM.....	1,260	221	630	420	525
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	68.0	0	1.0	1.0	4.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.3	.2	.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
35.9	31.6	32.3	30.1	24.9	23.2	28.2	22.5	26.0	21.9	26.2	22.5
28.2	25.9	25.8	25.4	20.5	19.3	21.9	18.5	19.7	16.8	19.5	17.7
6.9	5.2	6.3	4.4	4.1	3.4	5.5	3.6	5.7	4.3	4.5	3.6
.8	.6	.2	.4	.3	.5	.9	.4	.5	.8	2.2	1.2
1.3	1.2	1.3	1.2	1.2	1.2	1.3	1.2	1.3	1.3	1.4	1.3
.5	.5	.5	.5	.5	.5	.6	.5	.6	.6	.9	.9
4.0	4.0	4.0	4.0	6.0	4.0	4.0	6.0	6.0	5.0	4.0	9.0
71	58	86	63	84	63	84	60	69	62	86	61
46	42	62	45	61	44	60	45	45	45	63	49
18	13	22	18	27	18	27	13	18	18	35	13
35	35	53	35	53	35	35	35	35	35	35	35
70	47	70	62	70	53	70	48	70	53	70	53
70	70	105	70	105	70	105	70	70	70	105	70
140	105	140	105	140	105	140	105	140	140	175	116
148	140	210	140	210	140	210	140	140	140	245	175
210	210	280	263	350	245	280	216	280	280	292	228
175	145	211	160	209	160	200	148	166	140	186	133
102	86	127	91	110	89	109	88	93	74	98	64
168	146	215	172	219	153	197	155	164	155	204	159
81	76	91	100	144	74	98	89	83	71	110	76
151	170	352	150	191	194	222	156	146	165	202	172
63	94	268	52	111	92	128	101	44	77	55	84
840	550	1,260	945	1,260	630	1,050	840	630	630	525	420
840	1,260	2,100	1,050	2,310	1,260	2,100	1,260	876	945	876	945
165	141	206	158	208	158	195	147	159	143	179	136
91	82	119	94	121	86	105	88	90	77	96	73
39	33	53	53	69	53	65	26	53	39	69	26
105	105	105	105	105	105	105	105	105	105	105	105
158	105	210	105	210	146	210	105	158	105	185	105
210	210	210	210	210	210	210	210	210	210	210	210
210	210	315	210	420	210	315	210	210	210	315	210
420	292	420	315	420	315	420	315	315	210	420	210
420	420	631	420	631	420	525	420	584	420	438	315
630	550	1,260	945	1,260	630	840	840	630	630	525	420
5.4	4.1	2.9	3.5	14.1	6.3	10.5	10.1	1.5	1.4	1.9	0
.6	.5	.5	.7	1.2	.4	.6	.5	.4	.3	.9	0

TABLE 10.06.--HOME-FRIED AND HASH-BROWED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	12.1	1.4	13.4	11.8	12.5
ON ONLY 1 OF 3 DAYS.....PCT.....	10.3	1.4	11.0	10.1	11.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.5	0	1.9	1.5	1.0
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.5	.2	.4
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER....	1.2	1.0	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.0	.6	.4	.5
MAXIMUM.....DO.....	5.0	1.0	4.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	51	19	30	35	42
STANDARD DEVIATION.....GM.....	41	18	27	28	30
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	13		7	7	13
25TH.....GM.....	28		14	19	26
50TH.....GM.....	43		28	28	28
75TH.....GM.....	57		28	43	57
90TH.....GM.....	103		57	57	80
95TH.....GM.....	113		61	85	113
99TH.....GM.....	207		198	113	142
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	130	58	71	85	108
STANDARD DEVIATION.....GM.....	85	54	42	53	65
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	130	0	91	96	129
STANDARD DEVIATION.....GM.....	78	0	68	49	71
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	151	0	52	125	103
STANDARD DEVIATION.....GM.....	97	0	57	87	36
MAXIMUM IN ANY 1 DAY.....GM.....	1,020	170	340	340	340
MAXIMUM OVER 3 DAYS.....GM.....	1,530	170	680	723	680
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	128	58	72	88	108
STANDARD DEVIATION.....GM.....	81	54	51	57	65
PERCENTILES: <u>1</u> /					
5TH.....GM.....	39		8	21	39
25TH.....GM.....	78		43	43	78
50TH.....GM.....	85		64	85	85
75TH.....GM.....	170		85	85	155
90TH.....GM.....	227		155	170	170
95TH.....GM.....	340		170	170	210
99TH.....GM.....	340		340	340	340
MAXIMUM.....GM.....	850	170	340	340	340
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER....	58.6	0	.8	1.0	.9
PROPORTION OF EATING OCCASIONS.....PCT.....	1.1	0	.5	.4	.3

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
12.2	11.2	12.1	10.8	15.1	10.0	15.1	11.2	17.8	8.8	10.1	9.4
10.6	9.9	9.6	9.1	12.5	8.8	12.9	9.6	14.8	7.9	7.9	8.2
1.4	1.3	2.0	1.6	2.1	1.1	1.9	1.5	2.5	.7	1.7	1.2
.2	0	.4	(2)	.5	.2	.4	.1	.4	.2	.5	0
1.2	1.1	1.3	1.2	1.2	1.2	1.2	1.2	1.2	1.1	1.3	1.2
.5	.4	.6	.4	.5	.5	.5	.5	.5	.4	.8	.4
5.0	3.0	4.0	3.0	4.0	5.0	5.0	5.0	5.0	3.0	5.0	2.0
50	47	66	55	66	48	58	44	57	41	66	42
40	34	43	45	50	33	43	33	49	24	72	24
14	13	23	13	22	12	14	7	14	14	17	13
26	26	28	28	28	28	28	26	28	26	35	26
43	28	57	52	57	43	57	37	52	39	57	28
57	57	85	57	78	57	57	57	57	57	57	57
95	85	113	113	113	85	113	85	103	57	108	63
113	113	155	113	170	113	120	113	140	85	170	103
255	170	227	283	255	174	227	155	227	113	433	113
128	122	166	141	164	126	151	117	137	112	134	113
87	72	89	109	105	83	91	73	78	55	61	59
154	140	142	134	152	117	142	109	159	117	143	106
94	82	72	64	75	59	78	75	117	61	55	44
86	0	175	340	201	122	150	143	187	62	308	0
59	0	54	0	84	42	117	50	90	21	167	0
850	340	510	850	1,020	816	680	1,020	850	340	510	340
850	680	808	1,020	1,083	816	1,445	1,530	1,445	510	1,300	340
129	124	157	140	160	121	143	112	142	110	146	107
82	76	86	102	95	77	83	69	82	58	83	56
35	39	59	35	53	34	43	21	43	39	68	21
78	78	85	78	85	78	85	70	85	70	85	78
85	85	155	128	155	85	128	85	140	85	140	85
170	170	170	170	170	170	170	170	170	170	170	170
233	174	340	255	340	170	255	170	255	170	170	170
310	340	340	340	340	255	340	255	310	170	340	170
510	340	340	340	510	340	340	340	425	340	465	255
510	340	510	850	680	816	510	510	510	340	465	255
5.3	5.2	2.4	3.9	11.3	2.7	8.4	12.8	.8	1.4	0	1.9
1.8	1.9	1.1	2.1	1.5	.4	.9	1.4	.3	.8	0	1.8

TABLE 10.07.--FRENCH FRIED POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	25.3	2.6	27.7	33.2	34.5
ON ONLY 1 OF 3 DAYS.....PCT.....	20.7	2.3	24.7	28.3	28.6
ON ONLY 2 OF 3 DAYS.....PCT.....	3.9	.2	2.7	4.6	5.2
ON ALL 3 OF 3 DAYS.....PCT.....	.7	.1	.3	.4	.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.2	1.1	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.7	.4	.5	.5
MAXIMUM.....DO.....	6.0	4.0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	27	8	16	21	23
STANDARD DEVIATION.....GM.....	22	4	13	15	17
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		5	5	5
25TH.....GM.....	14		10	10	10
50TH.....GM.....	19		10	19	19
75TH.....GM.....	36		19	23	28
90TH.....GM.....	48		29	38	45
95TH.....GM.....	68		45	45	57
99TH.....GM.....	113		87	80	100
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	66	21	42	53	57
STANDARD DEVIATION.....GM.....	49	8	34	36	37
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	67	28	47	53	57
STANDARD DEVIATION.....GM.....	40	0	29	28	31
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	73	13	32	49	77
STANDARD DEVIATION.....GM.....	38	0	5	9	29
MAXIMUM IN ANY 1 DAY.....GM.....	816	34	285	272	300
MAXIMUM OVER 3 DAYS.....GM.....	816	57	285	329	402
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	64	20	42	51	56
STANDARD DEVIATION.....GM.....	45	9	32	32	34
PERCENTILES: <u>1/</u>					
5TH.....GM.....	14		14	14	14
25TH.....GM.....	29		29	29	29
50TH.....GM.....	57		29	57	57
75TH.....GM.....	70		57	57	60
90TH.....GM.....	114		70	86	114
95TH.....GM.....	136		102	120	136
99TH.....GM.....	268		136	136	136
MAXIMUM.....GM.....	680	34	285	272	272
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	213.6	0	4.2	7.6	14.9
PROPORTION OF EATING OCCASIONS.....PCT.....	1.8	0	1.3	1.1	1.9

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
35.6	33.8	38.6	34.9	37.1	26.7	23.1	16.3	12.3	7.6	8.2	6.5
28.9	27.9	27.5	26.8	27.4	22.5	19.2	14.5	11.0	7.3	7.9	5.3
6.0	5.0	8.5	6.4	8.0	3.8	3.4	1.7	1.3	.2	.3	1.0
.7	1.0	2.7	1.7	1.7	.5	.5	.1	0	.1	0	.2
1.3	1.2	1.4	1.4	1.4	1.2	1.2	1.1	1.2	1.1	1.0	1.2
.5	.5	.7	.7	.7	.5	.5	.4	.4	.3	.2	.5
4.0	5.0	5.0	6.0	5.0	5.0	4.0	4.0	3.0	3.0	2.0	3.0
28	26	36	29	34	26	27	23	23	22	24	25
21	21	30	24	28	19	22	18	15	15	17	26
7	5	10	7	10	7	7	6	10	5		5
17	12	19	17	19	16	17	11	14	10		10
19	19	29	23	23	19	19	19	19	19		19
38	34	44	38	38	34	38	23	29	23		19
50	45	68	52	67	45	48	42	42	45		69
67	62	91	76	91	60	68	48	56	61		96
95	101	159	136	148	95	131	110	76	74		108
71	64	85	68	80	66	69	59	65	62	68	45
53	45	81	47	59	43	53	35	42	40	50	18
61	62	78	66	71	68	69	67	57	75	82	100
33	39	41	39	49	33	41	43	20	25	0	36
73	72	69	75	81	61	69	80	0	74	0	108
27	26	31	43	41	39	29	69	0	0	0	0
544	303	816	408	680	544	680	408	272	204	285	152
544	544	816	565	734	710	680	587	272	222	285	325
65	62	75	64	74	64	68	59	61	62	69	63
45	43	57	44	54	40	50	39	38	39	50	37
22	15	22	14	22	22	22	14	22	14		14
29	29	57	29	43	35	43	29	38	29		29
57	57	57	57	57	57	57	57	57	57		57
70	70	86	70	86	70	70	68	68	70		67
114	114	134	114	134	114	114	113	91	134		136
136	136	150	136	170	136	136	136	136	136		136
204	272	327	272	272	204	272	227	227	204		152
544	303	544	340	680	544	680	408	272	204	285	152
25.5	15.4	12.1	15.7	28.3	20.2	32.6	17.5	5.2	11.0	1.2	2.2
2.7	1.7	1.6	2.2	1.4	1.2	2.4	1.3	3.3	7.8	2.6	2.9

TABLE 10.08.--POTATO SALAD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.4	.2	4.8	4.9	4.6
ON ONLY 1 OF 3 DAYS.....PCT.....	5.5	0	4.0	4.2	4.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	.2	.7	.7	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	2.0	1.2	1.2	1.1
STANDARD DEVIATION.....DO.....	.5	.0	.5	.4	.3
MAXIMUM.....DO.....	5.0	2.0	3.0	3.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	72	26	33	42	53
STANDARD DEVIATION.....GM.....	55	0	26	33	33
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	10		3	5	10
25TH.....GM.....	42		21	21	28
50TH.....GM.....	63		28	42	42
75TH.....GM.....	83		42	42	83
90TH.....GM.....	167		63	83	83
95TH.....GM.....	167		104	83	83
99TH.....GM.....	250		125	167	167
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	187	0	81	111	153
STANDARD DEVIATION.....GM.....	125	0	50	88	99
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	203	39	91	111	128
STANDARD DEVIATION.....GM.....	127	0	54	56	6
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	134	0	111	0	0
STANDARD DEVIATION.....GM.....	41	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,250	63	250	500	500
MAXIMUM OVER 3 DAYS.....GM.....	2,000	78	375	500	500
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	177	39	81	106	146
STANDARD DEVIATION.....GM.....	110	23	46	80	88
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31		16	16	30
25TH.....GM.....	125		63	63	83
50TH.....GM.....	125		63	94	125
75TH.....GM.....	250		125	125	250
90TH.....GM.....	250		125	155	250
95TH.....GM.....	375		125	250	250
99TH.....GM.....	500		250	500	500
MAXIMUM.....GM.....	750	63	250	500	500
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	17.6	0	0	0	2.1
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	0	2.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.8	5.0	6.2	5.5	7.5	6.6	8.0	7.6	5.9	5.6	3.6	3.9
5.5	4.6	5.9	5.0	6.5	5.6	6.7	6.6	5.0	4.3	3.4	3.6
.3	.4	.4	.5	.8	.9	1.2	.9	.8	1.0	.2	.3
0	(2)	0	0	.1	.1	(2)	(2)	.2	.2	0	0
1.1	1.2	1.2	1.1	1.2	1.2	1.2	1.2	1.3	1.4	1.2	1.3
.4	.5	.4	.4	.6	.5	.5	.5	.7	.7	.4	.5
3.0	4.0	3.0	3.0	4.0	4.0	3.0	4.0	4.0	5.0	2.0	2.0
62	68	93	63	90	71	85	66	68	63	87	49
45	45	56	44	72	47	60	53	46	41	65	33
10	21	32	10	21	10	21	10	5	10		
42	42	42	42	42	42	42	42	42	41		
42	42	83	42	83	63	83	42	42	42		
83	83	125	83	83	83	96	83	83	83		
104	125	167	89	167	125	167	146	167	125		
167	167	167	167	250	167	167	167	167	167		
208	229	250	250	292	250	313	292	167	167		
177	184	266	164	230	185	218	169	171	159	237	141
126	114	157	91	145	111	125	118	109	102	165	99
193	200	226	228	281	181	231	209	191	143	375	112
75	90	114	97	192	92	134	120	75	64	0	19
0	132	0	0	163	144	167	57	138	107	0	0
0	0	0	0	24	13	0	0	0	55	0	0
750	688	750	500	1,250	625	1,000	875	500	500	500	500
750	750	750	750	2,000	750	1,500	1,000	500	500	750	500
165	171	240	168	218	172	210	167	161	140	218	112
101	88	125	87	134	97	120	99	86	91	124	71
30	63	95	31	63	31	63	30	16	16		
125	125	125	125	125	125	125	125	125	63		
125	125	250	125	250	125	250	125	125	125		
250	250	250	250	250	250	250	250	250	250		
250	250	375	250	375	250	375	250	250	250		
375	250	500	250	500	250	500	375	250	250		
500	375	750	375	750	500	500	500	500	500		
500	625	750	500	750	500	750	500	500	500	500	250
1.4	0	1.1	0	1.5	3.4	.9	4.6	0	2.4	0	0
1.1	0	1.1	0	.4	.8	.2	.7	0	1.8	0	0

TABLE 10.09.--CABBAGE SALAD AND COLESLAW
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.7	0	3.4	4.9	8.5
ON ONLY 1 OF 3 DAYS.....PCT.....	8.5	0	3.1	4.4	7.7
ON ONLY 2 OF 3 DAYS.....PCT.....	1.1	0	.3	.4	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.1	0	0	(2)	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	0	.3	.4	.3
MAXIMUM.....DO.....	6.0	0	2.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	27	0	12	15	19
STANDARD DEVIATION.....GM.....	20	0	7	13	12
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	5			3	3
25TH.....GM.....	15			5	10
50TH.....GM.....	20			10	20
75TH.....GM.....	40			20	20
90TH.....GM.....	40			40	40
95TH.....GM.....	60			40	40
99TH.....GM.....	90			60	60
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	70	0	36	41	53
STANDARD DEVIATION.....GM.....	45	0	22	29	31
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	70	0	17	51	49
STANDARD DEVIATION.....GM.....	45	0	9	34	29
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	80	0	0	2	0
STANDARD DEVIATION.....GM.....	47	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,080	0	68	120	120
MAXIMUM OVER 3 DAYS.....GM.....	1,140	0	68	240	240
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	68	0	33	41	51
STANDARD DEVIATION.....GM.....	45	0	22	31	31
PERCENTILES: <u>1/</u>					
5TH.....GM.....	15			5	8
25TH.....GM.....	40			15	30
50TH.....GM.....	60			30	60
75TH.....GM.....	90			60	60
90TH.....GM.....	120			68	120
95TH.....GM.....	120			120	120
99TH.....GM.....	240			120	120
MAXIMUM.....GM.....	1,020	0	68	120	120
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	73.7	0	0	.8	2.6
PROPORTION OF EATING OCCASIONS.....PCT.....	1.7	0	0	.8	1.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
9.6	9.3	9.8	9.5	9.7	8.5	11.8	12.1	10.6	10.7	8.1	10.7
8.4	8.2	8.3	8.5	8.4	7.7	9.9	10.5	8.6	9.2	7.9	9.1
1.1	1.0	1.4	.9	1.1	.8	1.6	1.5	1.9	1.5	.3	1.3
.1	(2)	0	.1	.1	.1	.2	.1	.1	.1	0	.3
1.2	1.2	1.2	1.1	1.2	1.1	1.2	1.2	1.2	1.2	1.1	1.2
.5	.4	.5	.4	.5	.4	.6	.4	.5	.5	.3	.5
5.0	3.0	3.0	3.0	4.0	4.0	6.0	4.0	3.0	4.0	2.0	3.0
24	24	31	25	31	26	32	26	29	26	24	28
15	17	22	18	22	26	22	17	22	21	11	21
5	5	10	8	8	5	10	5	5	5		3
13	10	15	15	20	13	20	15	13	10		15
20	20	20	20	20	20	20	20	20	20		20
30	33	40	40	40	40	40	40	40	30		40
40	40	60	40	60	40	60	43	60	40		60
50	60	80	53	80	47	80	60	80	73		80
80	80	120	120	100	80	100	80	120	140		100
62	65	77	68	83	68	82	69	66	61	69	70
34	42	54	44	57	45	46	42	41	32	32	45
68	61	88	63	70	64	71	68	88	88	53	77
29	40	38	29	29	96	35	30	41	53	7	32
37	47	0	88	119	92	94	56	50	60	0	93
12	0	0	30	28	35	65	14	0	0	0	10
180	240	360	425	540	1,080	240	240	240	360	120	240
240	360	360	425	540	1,140	660	300	360	450	120	300
60	61	77	66	78	67	78	66	70	64	64	71
34	40	51	41	49	63	43	39	43	41	33	38
15	14	30	15	23	14	30	15	15	15		23
30	30	40	40	60	30	60	40	40	30		46
60	60	60	60	60	60	60	60	60	60		60
60	60	120	80	120	60	120	90	120	60		120
120	120	120	120	120	120	120	120	120	120		120
120	120	120	120	180	120	120	120	120	120		120
120	180	240	120	240	240	240	180	240	180		180
180	240	360	425	360	1,020	240	240	240	360	120	180
2.4	4.8	5.5	0	9.5	16.2	17.0	12.1	0	2.0	0	.9
1.0	2.0	3.4	0	2.1	3.1	2.4	1.2	0	.9	0	.7

TABLE 10.10.--RAW CARROTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.0	.8	3.4	5.4	9.8
ON ONLY 1 OF 3 DAYS.....PCT.....	4.0	.6	3.0	4.3	7.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	.2	.4	.9	1.5
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	.1	.2	.5
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.3	1.3	1.2	1.3	1.3
STANDARD DEVIATION.....DO.....	.6	.5	.8	.7	.8
MAXIMUM.....DO.....	7.0	2.0	6.0	6.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	18	16	11	17	17
STANDARD DEVIATION.....GM.....	19	11	13	18	22
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1			1	1
25TH.....GM.....	5			4	4
50TH.....GM.....	17			13	10
75TH.....GM.....	22			20	20
90TH.....GM.....	41			33	37
95TH.....GM.....	51			50	51
99TH.....GM.....	96			122	110
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	44	27	29	36	38
STANDARD DEVIATION.....GM.....	44	1	27	29	45
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	47	50	18	50	34
STANDARD DEVIATION.....GM.....	35	0	16	39	24
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	46	0	77	40	73
STANDARD DEVIATION.....GM.....	31	0	0	28	48
MAXIMUM IN ANY 1 DAY.....GM.....	500	50	112	183	300
MAXIMUM OVER 3 DAYS.....GM.....	500	100	232	366	495
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	43	37	28	38	38
STANDARD DEVIATION.....GM.....	40	12	25	33	41
PERCENTILES: <u>1/</u>					
5TH.....GM.....	4			4	4
25TH.....GM.....	13			12	10
50TH.....GM.....	31			31	28
75TH.....GM.....	55			54	50
90TH.....GM.....	100			61	72
95TH.....GM.....	122			84	110
99TH.....GM.....	183			183	165
MAXIMUM.....GM.....	500	50	100	183	300
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	7.0	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
8.6	6.5	4.5	5.5	3.4	5.2	4.0	4.9	3.8	3.9	.9	2.8
7.0	5.5	3.9	4.7	2.9	4.1	3.2	4.2	3.1	2.9	.9	1.8
1.3	.9	.6	.5	.5	1.0	.6	.5	.7	.8	0	.8
.2	.2	.1	.3	.1	.1	.2	.2	0	.2	0	.2
1.3	1.2	1.2	1.3	1.2	1.3	1.3	1.2	1.2	1.4	1.0	1.5
.6	.5	.4	.9	.5	.5	.7	.5	.4	.8	.0	.8
4.0	4.0	3.0	7.0	3.0	3.0	4.0	4.0	2.0	4.0	1.0	4.0
16	13	16	17	23	21	23	16	12	22	24	16
18	13	15	19	24	18	23	14	18	20	12	14
1	1	1	1	4	3	2	1		2		
3	4	5	4	9	9	8	5		9		
10	9	11	9	17	17	17	17		17		
22	18	20	19	33	30	33	20		33		
37	33	40	50	41	46	50	36		51		
48	37	50	58	81	51	74	41		61		
73	65	72	96	167	85	102	81		87		
38	33	43	33	61	52	57	41	29	48	72	37
40	31	42	29	64	45	54	35	35	43	36	42
45	40	35	77	59	53	52	42	36	57	0	34
39	28	19	38	50	29	35	30	46	37	0	17
49	17	71	51	38	42	53	32	0	51	0	21
16	10	0	24	27	5	37	16	0	0	0	0
300	200	216	150	500	300	305	200	165	183	150	110
400	200	216	288	500	300	324	268	300	261	150	110
39	33	42	39	57	51	53	39	30	48	72	33
36	31	39	35	60	40	48	32	33	37	36	32
4	4	4	4	7	8	7	4		4		
8	8	12	13	20	28	20	13		25		
28	25	30	25	50	50	50	28		50		
55	50	61	50	61	61	72	50		55		
100	72	72	72	122	100	100	72		100		
100	100	120	122	144	122	144	108		100		
150	144	216	150	244	200	300	150		183		
220	200	216	150	500	300	305	200	150	183	150	110
.9	.9	2.7	0	0	1.4	0	0	0	1.1	0	0
.4	.5	3.8	0	0	.4	0	0	0	1.2	0	0

TABLE 10.11.--RAW CELERY
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS ^{1/} (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.5	.2	1.8	3.1	5.4
ON ONLY 1 OF 3 DAYS.....PCT.....	3.6	.2	1.6	2.5	4.4
ON ONLY 2 OF 3 DAYS.....PCT.....	.7	0	.1	.3	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	.3	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE ^{1/}NUMBER....	1.3	1.0	1.1	1.4	1.3
STANDARD DEVIATION.....DO.....	.6	.0	.3	.8	.6
MAXIMUM.....DO.....	5.0	1.0	2.0	5.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY ^{1/}GM.....	14	3	8	13	12
STANDARD DEVIATION.....GM.....	13	0	6	14	9
AVERAGE PER DAY AT SPECIFIED PERCENTILES: ^{1/}					
5TH.....GM.....	3			2	3
25TH.....GM.....	7			6	6
50TH.....GM.....	10			9	9
75TH.....GM.....	19			13	13
90TH.....GM.....	28			20	20
95TH.....GM.....	40			37	28
99TH.....GM.....	65			76	59
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	34	10	25	27	29
STANDARD DEVIATION.....GM.....	26	0	18	17	19
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	36	0	14	20	29
STANDARD DEVIATION.....GM.....	25	0	6	7	23
USED ON ALL 3 OF 3 DAYS:					
AVERAGE ^{1/}GM.....	38	0	0	51	25
STANDARD DEVIATION.....GM.....	17	0	0	22	2
MAXIMUM IN ANY 1 DAY.....GM.....	280	10	84	112	120
MAXIMUM OVER 3 DAYS.....GM.....	330	10	84	228	176
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE ^{1/}GM.....	33	10	22	29	28
STANDARD DEVIATION.....GM.....	24	0	14	18	20
PERCENTILES: ^{1/}					
5TH.....GM.....	8			4	8
25TH.....GM.....	17			15	14
50TH.....GM.....	28			28	25
75TH.....GM.....	40			40	30
90TH.....GM.....	60			56	51
95TH.....GM.....	80			60	60
99TH.....GM.....	120			60	120
MAXIMUM.....GM.....	204	10	60	84	120
QUANTITIES NOT SPECIFIED BY USERS: ^{1/}					
EATING OCCASIONS.....NUMBER....	58.7	0	0	2.3	4.8
PROPORTION OF EATING OCCASIONS.....PCT.....	2.6	0	0	3.2	3.9

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.7	4.8	3.0	4.1	3.0	4.0	4.9	5.7	7.2	5.5	4.3	5.4
5.2	4.1	2.5	3.5	2.4	3.4	3.6	4.4	5.1	4.2	3.2	3.9
.4	.7	.4	.4	.5	.5	.9	1.1	1.6	1.0	1.1	1.1
.1	(2)	.1	.3	.1	(2)	.4	.3	.4	.3	0	.3
1.1	1.2	1.2	1.3	1.3	1.3	1.4	1.4	1.4	1.3	1.3	1.4
.4	.4	.4	.7	.6	.5	.7	.7	.7	.7	.4	.7
3.0	3.0	3.0	4.0	4.0	5.0	4.0	5.0	5.0	5.0	2.0	4.0
12	11	12	16	14	14	18	14	17	15	18	14
10	7	10	15	10	12	16	14	14	10	12	11
3	3		2	3	3	3	3	3	5		3
6	6		6	7	7	7	6	9	9		6
9	9		13	13	10	13	10	11	11		10
13	13		19	19	19	23	19	19	19		19
23	20		33	28	27	39	30	37	28		33
33	27		40	37	43	45	40	51	40		36
40	37		110	40	65	100	77	67	41		45
30	27	30	37	39	37	40	32	37	37	41	33
22	16	26	27	25	31	28	29	24	22	28	25
32	30	31	35	28	35	48	38	36	33	47	28
9	12	16	16	15	19	33	28	30	14	11	18
51	30	30	48	41	62	35	33	39	40	0	35
17	0	0	29	19	4	15	14	3	9	0	2
120	90	153	240	120	280	200	204	168	120	120	100
200	112	153	330	204	280	300	280	200	177	120	136
31	27	30	37	35	34	39	32	36	35	44	30
22	16	24	26	22	24	29	27	25	19	25	18
10	8		10	8	8	8	8	7	14		8
17	17		20	17	17	20	15	20	20		17
28	28		30	28	28	30	28	28	30		28
34	31		50	50	40	56	34	40	40		40
56	50		60	56	60	68	60	60	56		60
84	56		100	70	80	85	84	80	60		68
102	80		120	112	136	136	168	120	120		80
120	90	153	120	120	140	200	204	168	120	120	80
1.1	4.8	2.2	.9	5.4	10.6	5.3	14.3	1.8	2.5	2.7	0
.8	4.0	4.4	1.2	3.6	4.0	1.6	2.6	1.6	2.0	9.3	0

TABLE 10.12.--RAW CUCUMBERS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	5.6	.6	1.6	3.5	4.1
ON ONLY 1 OF 3 DAYS.....PCT.....	4.1	.4	1.3	2.4	3.1
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	.3	.3	1.2	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	0	0	.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.4	1.4	1.2	1.4	1.3
STANDARD DEVIATION.....DO.....	.7	.5	.4	.6	.6
MAXIMUM.....DO.....	6.0	2.0	2.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	37	30	16	27	30
STANDARD DEVIATION.....GM.....	45	26	14	32	43
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	3			1	3
25TH.....GM.....	12			7	5
50TH.....GM.....	23			15	18
75TH.....GM.....	47			37	37
90TH.....GM.....	79			53	75
95TH.....GM.....	105			105	105
99TH.....GM.....	220			179	330
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	77	24	46	50	66
STANDARD DEVIATION.....GM.....	72	16	40	46	70
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	88	89	30	72	52
STANDARD DEVIATION.....GM.....	80	0	20	68	51
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	105	0	0	0	138
STANDARD DEVIATION.....GM.....	99	0	0	0	117
MAXIMUM IN ANY 1 DAY.....GM.....	840	158	110	316	440
MAXIMUM OVER 3 DAYS.....GM.....	1,580	178	110	536	990
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	80	63	40	58	68
STANDARD DEVIATION.....GM.....	76	63	36	50	73
PERCENTILES: <u>1/</u>					
5TH.....GM.....	8			8	8
25TH.....GM.....	24			17	18
50TH.....GM.....	70			46	40
75TH.....GM.....	110			79	79
90TH.....GM.....	158			140	140
95TH.....GM.....	220			158	220
99TH.....GM.....	316			220	316
MAXIMUM.....GM.....	840	158	110	220	440
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	22.3	0	0	0	1.0
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	0	0	0	1.0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.2	4.6	3.9	6.3	4.2	6.0	6.7	8.0	6.1	7.2	6.4	3.3
2.2	3.7	3.5	5.0	3.5	4.7	4.5	5.8	4.5	4.6	3.3	2.1
.9	.7	.2	.9	.7	1.1	1.8	1.8	1.2	1.4	2.0	1.0
.1	.2	.2	.3	.1	.2	.4	.4	.3	1.2	1.0	.2
1.4	1.3	1.2	1.3	1.2	1.3	1.5	1.4	1.4	1.6	1.8	1.7
.6	.6	.5	.8	.5	.6	.7	.7	.6	1.0	.8	1.0
4.0	4.0	3.0	6.0	3.0	3.0	5.0	4.0	3.0	5.0	3.0	4.0
34	31	31	27	38	36	47	36	33	45	54	28
28	38	30	32	54	39	53	39	33	74	37	20
4	3	3	3	3	4	4	3	8	5		
12	7	12	7	9	12	16	12	12	12		
23	18	18	18	23	23	29	23	23	26		
47	37	47	37	37	53	70	47	37	53		
73	73	78	68	93	73	97	76	92	73		
76	100	93	73	108	93	146	105	93	110		
147	220	140	158	373	220	293	220	167	526		
77	69	75	63	80	84	84	76	82	80	89	86
64	90	70	72	85	68	71	71	57	65	54	59
76	99	77	74	132	80	109	86	78	70	99	29
52	80	34	77	140	72	87	72	83	49	47	19
59	57	110	63	149	118	137	87	42	119	110	70
20	23	20	21	92	92	102	75	38	155	0	0
220	660	280	474	840	440	500	632	280	632	220	220
440	660	420	660	1,120	948	1,220	880	500	1,580	330	220
75	72	76	62	96	83	96	78	71	83	93	50
58	82	64	64	115	72	80	67	64	84	50	52
12	8	8	8	8	12	12	9	12	9		
35	20	32	17	24	32	35	28	20	17		
55	46	55	55	55	70	70	70	53	53		
110	105	110	70	110	110	110	110	110	110		
158	158	140	140	220	185	220	158	158	220		
220	220	220	220	280	220	280	220	220	316		
220	280	280	280	840	316	316	280	280	316		
220	660	280	474	840	440	440	440	280	316	220	220
1.2	.8	1.8	0	1.9	3.1	2.1	4.7	1.8	2.0	0	1.8
1.3	.6	2.7	0	1.0	.8	.4	.6	1.8	1.0	0	3.2

TABLE 10.13.--LETTUCE AND TOSSED SALADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS 1/ (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	50.7	0	16.6	30.4	42.8
ON ONLY 1 OF 3 DAYS.....PCT.....	26.6	0	12.7	20.5	26.4
ON ONLY 2 OF 3 DAYS.....PCT.....	16.4	0	3.1	7.5	12.4
ON ALL 3 OF 3 DAYS.....PCT.....	7.8	0	.8	2.3	4.0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE 1/NUMBER....	1.8	0	1.4	1.5	1.6
STANDARD DEVIATION.....DO.....	1.0	0	.8	.8	.9
MAXIMUM.....DO.....	8.0	0	5.0	5.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY 1/GM.....	40	0	14	17	23
STANDARD DEVIATION.....GM.....	39	0	15	15	21
AVERAGE PER DAY AT SPECIFIED PERCENTILES: 1/					
5TH.....GM.....	3		2	2	3
25TH.....GM.....	13		4	5	9
50TH.....GM.....	31		9	13	18
75TH.....GM.....	54		18	23	31
90TH.....GM.....	90		32	36	47
95TH.....GM.....	113		47	43	67
99TH.....GM.....	186		65	71	104
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE 1/GM.....	67	0	29	34	43
STANDARD DEVIATION.....GM.....	62	0	26	29	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE 1/GM.....	73	0	34	39	48
STANDARD DEVIATION.....GM.....	54	0	27	23	31
USED ON ALL 3 OF 3 DAYS:					
AVERAGE 1/GM.....	80	0	44	35	49
STANDARD DEVIATION.....GM.....	55	0	25	21	32
MAXIMUM IN ANY 1 DAY.....GM.....	1,080	0	186	186	216
MAXIMUM OVER 3 DAYS.....GM.....	1,296	0	270	326	432
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE 1/GM.....	65	0	30	34	43
STANDARD DEVIATION.....GM.....	59	0	29	26	33
PERCENTILES: 1/					
5TH.....GM.....	10		5	7	10
25TH.....GM.....	20		10	14	14
50TH.....GM.....	55		23	28	30
75TH.....GM.....	93		47	47	55
90TH.....GM.....	140		55	67	93
95TH.....GM.....	186		93	93	108
99TH.....GM.....	270		135	108	140
MAXIMUM.....GM.....	1,080	0	186	156	216
QUANTITIES NOT SPECIFIED BY USERS: 1/					
EATING OCCASIONS.....NUMBER....	569.9	0	.7	4.4	19.4
PROPORTION OF EATING OCCASIONS.....PCT.....	1.6	0	.3	.6	1.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
45.8	47.5	47.7	49.0	54.2	56.5	59.3	60.9	49.5	49.4	33.1	40.7
28.1	27.7	25.9	27.4	29.9	30.3	27.7	28.3	23.8	24.8	15.8	19.1
13.6	15.5	15.5	15.2	16.0	18.9	20.4	21.1	15.4	16.0	10.1	12.4
4.1	4.4	6.3	6.4	8.3	7.3	11.1	11.4	10.3	8.6	7.3	9.2
1.6	1.7	1.8	1.7	1.8	1.8	2.0	2.0	1.9	1.9	1.9	1.9
.8	.9	1.0	.9	1.0	1.0	1.1	1.1	1.0	1.1	1.0	1.0
5.0	5.0	6.0	5.0	6.0	6.0	8.0	6.0	6.0	6.0	6.0	6.0
29	28	37	33	41	43	47	46	41	39	39	37
28	26	34	31	41	44	42	43	40	37	36	36
3	3	3	3	3	3	3	3	3	3	3	3
9	9	13	9	12	14	18	17	13	15	14	12
20	21	31	25	31	31	36	36	31	31	31	27
37	37	50	45	60	55	64	62	56	49	55	49
65	62	79	72	93	98	101	95	90	78	86	90
82	75	95	93	119	136	126	126	126	108	109	102
124	121	160	144	196	216	193	216	183	198	186	145
55	53	66	60	75	75	80	73	66	70	59	54
49	45	55	52	68	71	70	63	60	55	42	41
61	59	75	59	79	79	82	77	70	72	64	69
43	43	59	39	59	64	54	52	45	58	40	45
63	56	65	73	78	95	84	88	79	68	78	64
40	36	37	45	58	60	54	58	58	48	47	50
349	486	558	372	595	755	1,080	594	450	432	229	632
660	702	882	666	972	1,082	1,296	1,116	837	807	558	744
54	51	61	57	68	74	72	70	65	62	61	58
47	43	56	49	65	68	63	58	54	51	47	53
10	10	10	9	10	10	10	10	10	10	10	10
20	15	20	15	20	20	20	20	20	20	20	20
47	41	54	51	55	55	55	55	55	55	55	54
81	70	93	93	108	108	108	104	93	93	93	93
108	108	110	108	162	162	153	140	135	110	113	108
135	110	186	142	186	204	186	186	162	150	162	140
216	216	270	216	279	306	234	270	279	225	204	204
330	280	558	325	595	755	1,080	558	450	324	216	539
17.5	29.7	13.9	21.5	71.2	65.4	116.0	138.9	20.0	32.0	9.1	10.3
1.1	1.7	1.2	1.7	1.8	1.2	2.0	1.6	1.9	1.9	2.7	1.3

TABLE 10.14.--LETTUCE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	33.2	0	9.4	18.2	26.3
ON ONLY 1 OF 3 DAYS.....PCT.....	21.2	0	8.4	13.9	19.0
ON ONLY 2 OF 3 DAYS.....PCT.....	8.8	0	.9	3.3	5.7
ON ALL 3 OF 3 DAYS.....PCT.....	3.1	0	.1	1.0	1.6
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	1.6	0	1.2	1.3	1.4
STANDARD DEVIATION.....DO.....	.9	0	.5	.7	.7
MAXIMUM.....DO.....	8.0	0	4.0	5.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	19	0	8	9	11
STANDARD DEVIATION.....GM.....	22	0	10	9	10
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	3		1	2	2
25TH.....GM.....	5		3	3	3
50TH.....GM.....	11		5	7	9
75TH.....GM.....	23		9	10	17
90TH.....GM.....	43		18	20	28
95TH.....GM.....	60		30	28	32
99TH.....GM.....	110		60	45	45
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	36	0	21	19	24
STANDARD DEVIATION.....GM.....	42	0	27	16	19
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	40	0	25	25	29
STANDARD DEVIATION.....GM.....	34	0	18	17	18
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	45	0	30	21	24
STANDARD DEVIATION.....GM.....	36	0	0	10	12
MAXIMUM IN ANY 1 DAY.....GM.....	755	0	180	130	135
MAXIMUM OVER 3 DAYS.....GM.....	809	0	180	157	230
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	36	0	21	21	24
STANDARD DEVIATION.....GM.....	38	0	25	16	18
PERCENTILES: <u>1</u> /					
5TH.....GM.....	8		4	5	5
25TH.....GM.....	10		10	10	10
50TH.....GM.....	28		14	14	20
75TH.....GM.....	55		28	28	28
90TH.....GM.....	83		41	40	55
95TH.....GM.....	110		55	55	55
99TH.....GM.....	165		180	72	67
MAXIMUM.....GM.....	755	0	180	110	135
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	225.2	0	0	3.4	7.5
PROPORTION OF EATING OCCASIONS.....PCT.....	1.1	0	0	.8	1.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
29.4	29.8	32.7	32.8	37.7	36.0	38.1	40.1	34.4	32.7	20.9	29.3
21.3	20.6	21.4	20.6	23.7	23.8	22.8	24.6	20.4	19.6	12.0	16.9
6.6	7.1	8.5	9.9	9.9	10.2	10.8	11.5	8.4	8.4	5.7	7.7
1.6	2.1	2.7	2.3	4.1	2.0	4.5	4.0	5.6	4.7	3.2	4.7
1.4	1.5	1.6	1.6	1.6	1.5	1.7	1.6	1.7	1.7	1.8	1.7
.7	.8	.9	.8	.9	.8	1.0	.9	.9	1.0	1.0	1.0
5.0	5.0	6.0	5.0	6.0	6.0	8.0	6.0	5.0	6.0	6.0	6.0
14	15	18	17	19	20	21	21	25	20	19	22
16	16	19	19	21	26	23	24	27	23	14	25
2	2	3	3	3	3	3	3	3	3	2	3
5	5	7	5	7	5	7	7	7	6	9	7
9	9	10	10	13	13	14	13	16	13	17	17
18	18	23	20	26	25	28	28	30	28	27	28
31	33	43	37	45	45	48	47	48	45	37	45
40	41	55	55	62	63	64	68	90	57	43	57
89	67	90	90	95	136	110	110	135	150	55	180
30	29	36	31	39	42	38	38	46	35	39	48
29	27	36	32	40	58	34	42	53	34	33	75
35	36	38	36	40	42	43	43	43	42	35	41
30	33	27	27	32	38	36	35	33	39	20	27
38	29	43	46	39	58	49	49	53	43	32	33
27	17	30	38	35	45	35	39	40	32	12	27
280	280	275	270	290	755	360	360	450	270	180	539
550	550	380	549	559	755	590	809	450	539	180	539
31	30	35	33	36	40	38	38	44	36	32	38
29	29	33	32	36	50	36	38	42	36	20	49
7	7	10	7	10	7	10	10	10	10	7	8
10	10	10	10	10	10	10	10	14	10	18	14
24	20	20	20	20	28	28	28	28	28	28	28
41	40	55	55	55	55	55	55	55	55	55	55
55	55	67	65	83	96	83	83	110	67	55	65
83	83	110	90	110	120	110	110	135	110	55	110
135	135	135	135	165	220	136	165	138	165	90	165
270	275	275	270	270	755	270	280	450	270	110	539
5.5	13.5	5.0	9.7	23.8	27.0	48.5	40.0	8.7	15.4	9.1	8.2
.6	1.4	.7	1.3	1.0	.9	1.5	.9	1.3	1.6	4.6	1.6

TABLE 10.15.--TOSSED SALADS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	26.2	0	9.1	15.4	21.8
ON ONLY 1 OF 3 DAYS.....PCT.....	17.3	0	7.4	11.6	15.5
ON ONLY 2 OF 3 DAYS.....PCT.....	6.8	0	1.3	3.3	5.0
ON ALL 3 OF 3 DAYS.....PCT.....	2.1	0	.3	.5	1.2
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.5	0	1.3	1.3	1.4
STANDARD DEVIATION.....DO.....	.8	0	.7	.6	.7
MAXIMUM.....DO.....	6.0	0	5.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	50	0	16	22	30
STANDARD DEVIATION.....GM.....	39	0	15	15	23
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11		2	4	6
25TH.....GM.....	27		8	12	16
50TH.....GM.....	36		15	18	25
75TH.....GM.....	62		18	31	36
90TH.....GM.....	94		31	36	62
95TH.....GM.....	124		54	49	72
99TH.....GM.....	201		90	72	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	102	0	38	52	63
STANDARD DEVIATION.....GM.....	60	0	25	30	35
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	107	0	41	50	70
STANDARD DEVIATION.....GM.....	55	0	29	26	34
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	117	0	54	50	76
STANDARD DEVIATION.....GM.....	59	0	32	22	32
MAXIMUM IN ANY 1 DAY.....GM.....	1,080	0	186	186	216
MAXIMUM OVER 3 DAYS.....GM.....	1,296	0	270	326	432
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	101	0	39	49	64
STANDARD DEVIATION.....GM.....	56	0	29	27	34
PERCENTILES: <u>1/</u>					
5TH.....GM.....	30		6	13	16
25TH.....GM.....	60		15	28	41
50TH.....GM.....	93		33	47	54
75TH.....GM.....	108		54	54	93
90TH.....GM.....	186		70	93	108
95TH.....GM.....	216		93	108	108
99TH.....GM.....	279		108	108	162
MAXIMUM.....GM.....	1,080	0	186	147	216
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	339.0	0	.7	1.0	11.9
PROPORTION OF EATING OCCASIONS.....PCT.....	2.3	0	.6	.3	2.1

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
23.0	24.9	23.3	23.5	25.0	29.5	33.0	33.2	23.1	24.8	15.8	16.7
16.8	17.7	15.4	16.8	17.4	20.0	20.4	20.5	15.1	16.2	8.7	9.6
5.3	6.4	6.8	5.4	5.9	7.0	9.6	9.3	5.9	6.7	4.6	4.6
.9	.8	1.2	1.3	1.8	2.5	3.0	3.4	2.1	1.9	2.5	2.4
1.4	1.4	1.5	1.4	1.4	1.5	1.5	1.6	1.5	1.5	1.6	1.6
.6	.6	.7	.7	.7	.8	.8	.8	.8	.8	.8	.8
5.0	5.0	5.0	4.0	5.0	6.0	6.0	6.0	5.0	5.0	3.0	4.0
38	36	48	43	57	54	57	56	52	49	54	50
27	26	34	30	41	42	41	42	44	37	37	32
9	9	16	11	16	13	16	16	13	16	11	16
18	18	31	23	31	31	31	31	25	31	31	31
31	31	36	36	47	36	47	44	36	36	43	43
50	43	62	54	72	67	72	72	62	62	71	67
72	67	87	78	108	108	109	103	95	93	98	93
83	83	121	101	124	144	134	134	148	108	117	111
144	134	180	165	216	216	196	220	279	201	186	124
85	79	105	97	123	107	118	108	97	100	94	89
51	47	48	51	67	61	70	56	51	46	39	37
89	85	102	95	121	122	114	111	99	107	99	112
46	45	65	43	58	63	49	52	41	67	32	34
84	76	99	100	136	121	120	123	152	106	113	85
40	33	42	49	54	60	62	61	61	41	44	35
330	486	558	372	558	558	1,080	594	366	432	216	366
660	702	882	666	960	915	1,296	1,116	837	804	558	744
83	78	98	92	120	109	112	106	101	97	100	94
48	42	57	46	61	62	60	52	54	47	43	38
27	23	33	33	47	30	38	41	33	33	30	47
49	47	55	55	83	65	75	70	65	65	75	65
81	70	93	93	108	93	98	93	93	93	93	93
108	98	108	108	162	140	140	113	108	108	108	108
130	108	186	162	216	186	186	186	186	150	186	140
186	162	216	186	216	216	216	216	206	186	186	186
216	216	324	216	324	324	279	279	279	270	216	225
330	279	558	325	558	558	1,080	558	324	324	216	270
12.0	16.3	8.9	10.8	47.4	36.4	64.8	99.0	11.3	16.5	0	2.1
1.8	2.2	1.9	2.2	3.4	1.6	2.6	2.7	2.9	2.5	0	.8

TABLE 10.16.--RAW MATURE ONIONS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	8.5	0	1.4	3.1	3.9
ON ONLY 1 OF 3 DAYS.....PCT.....	7.1	0	1.1	2.6	3.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.2	0	.3	.6	.7
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	0	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	0	.4	.5	.4
MAXIMUM.....DO.....	5.0	0	2.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	13	0	9	8	8
STANDARD DEVIATION.....GM.....	16	0	8	11	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1			(2)	(2)
25TH.....GM.....	6			3	3
50TH.....GM.....	7			6	6
75TH.....GM.....	15			6	9
90TH.....GM.....	24			12	18
95TH.....GM.....	37			28	23
99TH.....GM.....	73			73	37
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	31	0	20	20	22
STANDARD DEVIATION.....GM.....	33	0	17	35	19
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	34	0	27	16	16
STANDARD DEVIATION.....GM.....	44	0	13	15	16
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	31	0	0	0	0
STANDARD DEVIATION.....GM.....	22	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	660	0	70	220	110
MAXIMUM OVER 3 DAYS.....GM.....	990	0	90	220	110
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	31	0	22	19	20
STANDARD DEVIATION.....GM.....	33	0	18	30	19
PERCENTILES: <u>1/</u>					
5TH.....GM.....	3			1	1
25TH.....GM.....	17			4	9
50TH.....GM.....	18			17	18
75TH.....GM.....	36			18	20
90TH.....GM.....	57			35	36
95TH.....GM.....	72			36	70
99TH.....GM.....	180			220	110
MAXIMUM.....GM.....	350	0	70	220	110
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	210.7	0	0	7.4	7.2
PROPORTION OF EATING OCCASIONS.....PCT.....	5.3	0	0	11.2	8.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
6.0	5.3	9.9	7.9	12.4	9.8	11.7	10.4	7.8	6.9	5.6	3.9
5.5	4.4	7.9	6.7	10.4	8.5	9.0	8.7	6.5	5.7	4.9	2.6
.4	.8	1.5	1.0	1.6	1.2	2.2	1.3	1.1	1.0	.7	1.2
(2)	.1	.5	.1	.3	.1	.6	.3	.3	.2	0	0
1.1	1.2	1.3	1.2	1.2	1.2	1.4	1.2	1.3	1.3	1.1	1.4
.3	.5	.7	.5	.5	.4	.6	.6	.6	.5	.3	.6
3.0	3.0	5.0	3.0	4.0	4.0	4.0	4.0	3.0	3.0	2.0	3.0
10	10	13	10	13	10	16	11	22	16	19	14
9	11	14	10	12	11	18	13	45	32	16	14
1	1	1	1	1	1	2	1	1	1		
6	6	6	3	6	5	6	6	6	6		
7	6	7	6	10	6	12	6	12	12		
12	12	14	12	18	12	18	12	23	18		
18	21	24	23	28	18	37	23	31	24		
24	37	42	37	37	25	47	37	46	35		
47	55	73	53	63	63	100	73	330	257		
26	26	32	27	35	27	38	28	41	31	53	29
20	30	37	26	31	29	37	34	38	31	48	31
35	30	29	25	32	27	37	30	99	71	47	36
24	21	26	22	28	22	34	21	160	105	10	24
18	13	27	22	29	19	43	25	38	26	0	0
0	6	16	5	10	5	29	18	7	11	0	0
110	220	274	140	220	220	360	350	660	440	210	140
140	220	274	158	280	220	369	350	990	770	210	158
27	26	29	25	33	26	36	27	50	39	52	31
20	27	29	26	29	27	35	31	68	53	43	27
3	3	3	3	5	3	3	3	3	5		
15	10	16	10	18	10	18	14	18	18		
18	18	18	18	20	18	25	18	35	18		
35	35	36	33	36	35	38	35	55	36		
54	54	55	54	70	55	70	55	70	55		
70	70	70	70	110	70	110	70	220	220		
110	110	220	110	140	150	180	175	330	330		
110	220	220	140	220	220	280	350	330	330	210	110
9.3	3.8	20.5	7.8	23.7	36.7	42.0	37.6	3.5	9.7	0	1.4
6.7	2.8	11.2	5.6	4.0	6.0	5.4	4.1	3.1	6.4	0	2.7

TABLE 10.17.--RAW TOMATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	27.8	.3	10.6	15.7	18.3
ON ONLY 1 OF 3 DAYS.....PCT.....	17.4	.3	8.5	11.6	14.2
ON ONLY 2 OF 3 DAYS.....PCT.....	7.4	0	1.8	3.3	3.3
ON ALL 3 OF 3 DAYS.....PCT.....	2.9	0	.3	.8	.8
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.6	1.0	1.3	1.4	1.3
STANDARD DEVIATION.....DO.....	1.0	.0	.6	.7	.7
MAXIMUM.....DO.....	8.0	1.0	4.0	7.0	6.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	44	7	20	24	24
STANDARD DEVIATION.....GM.....	42	2	17	23	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	10		4	5	7
25TH.....GM.....	20		10	10	10
50TH.....GM.....	30		15	20	20
75TH.....GM.....	60		29	30	30
90TH.....GM.....	91		46	51	45
95TH.....GM.....	123		60	71	61
99TH.....GM.....	205		80	111	120
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	78	21	43	54	53
STANDARD DEVIATION.....GM.....	59	7	32	55	34
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	91	0	61	54	66
STANDARD DEVIATION.....GM.....	56	0	23	32	43
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	110	0	57	64	59
STANDARD DEVIATION.....GM.....	62	0	19	32	18
MAXIMUM IN ANY 1 DAY.....GM.....	907	30	182	615	308
MAXIMUM OVER 3 DAYS.....GM.....	1,638	30	240	615	490
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	81	21	46	52	55
STANDARD DEVIATION.....GM.....	55	7	32	44	33
PERCENTILES: <u>1/</u>					
5TH.....GM.....	30		11	15	15
25TH.....GM.....	45		30	30	30
50TH.....GM.....	62		31	31	56
75TH.....GM.....	113		60	62	62
90TH.....GM.....	123		91	91	91
95TH.....GM.....	182		120	123	123
99TH.....GM.....	246		182	182	180
MAXIMUM.....GM.....	728	30	182	615	246
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	68.6	0	.8	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.4	0	.6	0	0

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
20.1	21.0	24.4	24.3	29.6	29.5	32.4	35.9	30.9	35.8	27.9	25.0
14.9	15.6	17.4	16.4	20.3	19.3	18.2	21.2	15.1	18.9	14.4	12.6
4.2	4.1	5.5	6.3	7.0	8.1	9.7	10.5	9.5	10.3	7.2	8.3
.9	1.3	1.5	1.6	2.3	2.2	4.4	4.2	6.3	6.6	6.2	4.1
1.4	1.4	1.5	1.5	1.5	1.5	1.8	1.7	1.9	1.9	2.0	1.9
.8	.8	.8	.8	.8	.8	1.1	1.0	1.1	1.1	1.3	1.1
5.0	6.0	5.0	5.0	7.0	7.0	8.0	6.0	6.0	6.0	8.0	6.0
35	33	37	33	41	39	55	48	60	54	60	56
36	30	37	29	33	35	53	42	56	43	56	44
8	10	10	10	10	10	10	10	10	10	10	10
15	15	15	15	20	20	21	20	30	21	21	21
21	21	21	21	30	30	40	36	41	41	40	41
41	41	41	41	58	50	66	61	81	71	81	81
71	62	80	62	82	81	121	99	123	122	121	112
100	82	120	101	102	102	164	132	143	140	143	143
212	133	205	150	171	164	262	205	204	205	273	241
73	73	73	66	83	75	87	83	92	85	76	89
59	54	61	47	67	58	57	60	55	58	36	56
84	75	83	72	92	85	104	95	100	92	104	96
59	57	54	37	44	60	67	56	51	40	45	38
110	86	107	94	93	95	129	110	121	116	129	118
75	38	64	42	42	42	78	58	82	45	70	59
735	519	615	364	675	907	735	677	546	615	455	362
1,014	822	733	546	675	1,361	1,599	1,107	1,638	738	818	800
74	71	75	66	81	75	93	84	93	86	92	89
58	49	56	44	54	49	63	57	65	44	43	48
23	30	30	28	30	28	30	30	30	30	30	30
30	31	30	30	45	34	60	56	60	60	60	60
60	60	60	60	62	62	90	62	90	90	90	90
90	90	90	90	120	91	123	120	123	123	120	123
123	123	123	123	123	123	180	123	180	123	180	135
180	180	180	180	182	180	182	182	182	153	182	182
364	246	270	182	246	246	360	273	270	182	182	246
675	369	492	364	675	546	728	675	546	615	273	270
4.0	4.1	0	2.6	9.3	12.5	13.8	12.7	1.8	5.0	0	2.0
.7	.6	0	.5	.5	.5	.5	.3	.3	.4	0	.4

TABLE 10.18.--COOKED BROCCOLI
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	6.2	1.0	5.7	3.8	5.6
ON ONLY 1 OF 3 DAYS.....PCT.....	5.7	1.0	5.3	3.6	5.3
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	0	.5	.2	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.0	1.1	1.1	1.0
STANDARD DEVIATION.....DO.....	.3	.0	.3	.2	.2
MAXIMUM.....DO.....	3.0	1.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	41	14	20	23	29
STANDARD DEVIATION.....GM.....	27	9	13	16	18
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	12		4	3	7
25TH.....GM.....	26		13	14	15
50TH.....GM.....	31		15	22	26
75TH.....GM.....	57		28	29	31
90TH.....GM.....	62		32	31	59
95TH.....GM.....	99		50	58	62
99TH.....GM.....	137		62	93	85
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	114	42	56	65	82
STANDARD DEVIATION.....GM.....	69	27	36	43	52
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	108	0	55	61	88
STANDARD DEVIATION.....GM.....	63	0	27	44	2
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	100	0	0	0	0
STANDARD DEVIATION.....GM.....	57	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	680	93	185	278	255
MAXIMUM OVER 3 DAYS.....GM.....	930	93	185	278	255
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	112	42	55	65	83
STANDARD DEVIATION.....GM.....	68	27	33	43	50
PERCENTILES: <u>1/</u>					
5TH.....GM.....	30		11	10	22
25TH.....GM.....	78		39	43	47
50TH.....GM.....	90		46	57	80
75TH.....GM.....	155		85	88	90
90TH.....GM.....	185		90	93	175
95TH.....GM.....	190		93	175	185
99TH.....GM.....	350		185	278	255
MAXIMUM.....GM.....	680	93	185	278	255
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	9.0	0	0	0	1.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.3	0	0	0	1.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.6	5.1	4.3	4.1	6.4	6.4	7.1	8.1	7.1	6.9	4.0	7.1
4.4	4.8	3.9	3.8	6.0	5.7	6.6	7.3	6.4	5.6	3.4	6.0
.2	.3	.4	.3	.4	.7	.5	.7	.7	1.3	.7	1.1
0	0	0	0	0	(2)	(2)	(2)	0	0	0	0
1.0	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.2	1.2
.2	.2	.3	.3	.3	.3	.3	.3	.3	.5	.4	.4
2.0	2.0	2.0	2.0	2.0	3.0	3.0	3.0	2.0	3.0	2.0	2.0
33	31	36	38	54	42	48	42	41	41	38	40
25	21	20	20	34	21	34	26	24	24	22	27
8	7	15	13	14	15	14	13	13	14		12
16	16	22	27	29	28	29	28	27	27		26
28	28	30	31	52	35	43	31	31	31		29
44	32	52	53	62	58	58	57	52	57		57
58	57	58	62	110	62	62	63	73	62		62
62	62	62	77	113	74	110	103	103	85		71
155	113	117	117	146	123	227	123	117	129		183
97	87	101	107	151	117	132	118	114	104	96	103
74	55	46	57	96	56	76	63	64	52	46	51
76	97	94	96	155	93	152	102	109	103	100	104
32	49	55	21	40	44	95	62	35	46	41	68
0	0	0	0	0	66	55	133	0	0	0	0
0	0	0	0	0	0	7	57	0	0	0	0
465	370	185	350	620	340	680	465	320	255	185	285
465	370	350	350	620	475	930	620	350	388	340	548
96	88	100	106	150	112	134	115	110	100	97	98
72	55	48	55	89	56	81	64	51	48	45	51
19	21	30	39	44	30	44	35	39	39		39
47	51	62	78	88	78	85	78	85	78		70
85	85	88	93	140	93	116	93	93	85		85
127	93	155	131	175	155	175	160	155	131		106
175	170	175	175	278	185	185	185	170	175		175
185	175	175	185	340	190	310	190	185	185		185
465	310	185	350	425	320	465	350	310	255		285
465	370	185	350	620	340	680	465	310	255	185	285
2.2	1.0	0	0	0	0	0	4.0	0	0	0	0
2.2	.8	0	0	0	0	0	.6	0	0	0	0

TABLE 10.19.--COOKED CABBAGE
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.7	.4	3.2	3.3	3.8
ON ONLY 1 OF 3 DAYS.....PCT.....	4.2	.3	2.7	3.0	3.5
ON ONLY 2 OF 3 DAYS.....PCT.....	.5	.2	.5	.3	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.4	1.2	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.5	.4	.5	.4
MAXIMUM.....DO.....	4.0	2.0	3.0	4.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	50	35	23	31	35
STANDARD DEVIATION.....GM.....	39	32	19	21	22
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	13			5	6
25TH.....GM.....	25			13	19
50TH.....GM.....	50			25	25
75TH.....GM.....	50			50	50
90TH.....GM.....	100			50	50
95TH.....GM.....	102			75	75
99TH.....GM.....	200			100	100
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	133	35	61	91	95
STANDARD DEVIATION.....GM.....	84	32	52	62	59
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	137	112	53	52	96
STANDARD DEVIATION.....GM.....	83	0	36	33	38
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	206	0	0	0	0
STANDARD DEVIATION.....GM.....	139	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	750	150	218	300	300
MAXIMUM OVER 3 DAYS.....GM.....	1,220	225	228	300	300
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	128	77	57	77	92
STANDARD DEVIATION.....GM.....	83	52	48	51	54
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28			9	18
25TH.....GM.....	75			38	38
50TH.....GM.....	145			75	75
75TH.....GM.....	150			93	150
90TH.....GM.....	225			150	150
95TH.....GM.....	300			150	150
99TH.....GM.....	450			150	225
MAXIMUM.....GM.....	610	150	218	300	300
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	32.4	0	.8	1.4	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.6	0	2.0	2.0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.9	4.5	4.5	4.3	3.1	3.9	5.3	6.0	5.9	6.7	10.1	6.5
3.8	4.3	4.3	4.0	2.7	3.4	4.8	5.6	5.0	5.2	9.0	4.7
.1	.2	.2	.3	.4	.4	.5	.4	.6	1.4	1.1	1.9
0	0	0	0	0	(2)	(2)	.1	.3	.1	0	0
1.1	1.1	1.1	1.1	1.2	1.2	1.2	1.1	1.3	1.4	1.2	1.3
.3	.3	.3	.3	.4	.4	.4	.4	.5	.7	.6	.5
3.0	3.0	2.0	2.0	3.0	4.0	3.0	4.0	3.0	4.0	4.0	2.0
42	43	49	45	58	48	56	50	50	65	58	57
28	36	22	30	39	33	32	48	37	52	36	32
7	9	13	13	13	13	13	9	19	19	25	24
25	25	38	25	30	25	36	25	25	38	25	25
38	25	50	48	50	50	50	48	50	50	50	50
50	50	50	50	75	50	75	50	50	100	75	75
100	75	75	75	100	100	100	100	100	102	100	100
100	141	100	100	145	102	117	102	100	125	138	102
150	150	102	150	200	140	150	250	250	407	200	150
124	119	146	126	153	131	157	129	126	153	148	133
84	88	63	85	86	77	88	92	62	81	70	70
110	176	73	123	168	115	137	151	185	153	198	134
37	115	52	41	84	60	55	82	99	119	59	42
0	0	0	0	0	237	133	312	43	200	0	0
0	0	0	0	0	0	0	140	0	33	0	0
450	450	305	450	435	375	600	750	450	610	413	305
450	750	305	450	600	712	600	1,220	750	1,220	600	450
117	121	129	119	150	125	146	132	118	142	144	129
79	91	65	81	84	75	82	94	73	94	55	68
22	26	38	38	38	28	38	28	36	38	75	38
75	75	75	75	75	75	75	75	75	75	75	75
75	75	150	75	150	145	150	132	113	150	150	150
150	150	150	150	150	150	150	150	150	150	150	150
150	225	200	218	300	225	300	300	150	300	225	150
300	300	300	300	300	300	300	305	300	300	225	300
450	450	305	450	435	305	450	450	450	610	300	305
450	450	305	450	435	305	600	610	450	610	300	305
2.0	1.9	0	2.1	2.3	4.1	9.7	2.8	0	1.1	0	4.2
2.3	1.9	0	2.9	1.7	1.7	3.2	.6	0	.7	0	4.8

TABLE 10.20.--COOKED CARROTS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.8	21.7	11.7	8.0	8.7
ON ONLY 1 OF 3 DAYS.....PCT.....	8.6	16.4	10.2	7.5	7.9
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	4.1	1.4	.5	.7
ON ALL 3 OF 3 DAYS.....PCT.....	.2	1.2	.2	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.4	1.2	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.7	.4	.3	.4
MAXIMUM.....DO.....	6.0	4.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	30	34	22	18	22
STANDARD DEVIATION.....GM.....	21	23	19	12	15
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	6	8	5	3	6
25TH.....GM.....	15	19	12	13	13
50TH.....GM.....	25	33	13	13	20
75TH.....GM.....	39	43	26	25	26
90TH.....GM.....	52	57	50	32	39
95TH.....GM.....	71	85	68	46	50
99TH.....GM.....	103	132	85	63	75
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	81	88	56	49	58
STANDARD DEVIATION.....GM.....	53	61	45	31	33
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	80	67	64	54	75
STANDARD DEVIATION.....GM.....	41	36	41	24	35
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	73	57	40	52	45
STANDARD DEVIATION.....GM.....	34	31	11	9	0
MAXIMUM IN ANY 1 DAY.....GM.....	736	425	255	155	192
MAXIMUM OVER 3 DAYS.....GM.....	736	425	300	233	300
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	79	71	54	49	59
STANDARD DEVIATION.....GM.....	50	41	38	31	33
PERCENTILES: <u>1/</u>					
5TH.....GM.....	19	14	10	10	18
25TH.....GM.....	46	46	38	38	38
50TH.....GM.....	75	57	39	39	50
75TH.....GM.....	92	98	71	59	75
90TH.....GM.....	150	128	113	78	92
95TH.....GM.....	155	128	150	113	144
99TH.....GM.....	276	213	192	155	155
MAXIMUM.....GM.....	736	213	213	155	192
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	25.2	0	0	1.4	3.8
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	.9	2.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
8.5	8.8	8.5	7.0	7.9	7.9	10.8	10.7	11.6	12.7	12.9	13.5
7.8	8.0	7.9	6.5	7.1	7.4	9.5	9.7	9.8	9.5	11.4	11.0
.7	.8	.6	.5	.6	.5	1.3	.9	1.5	2.6	1.2	1.5
.1	.1	0	0	.1	.1	.1	.1	.4	.6	.2	1.1
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.3	1.2	1.3
.5	.5	.3	.2	.4	.3	.4	.3	.5	.6	.5	.6
6.0	5.0	2.0	2.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
30	28	31	25	35	28	36	28	36	38	31	37
22	20	17	16	24	17	27	18	23	24	16	22
3	8	8	6	11	5	13	6	13	8	13	13
15	13	16	13	17	15	25	15	19	25	17	25
25	25	26	25	26	25	26	25	26	31	26	31
32	32	50	26	50	33	50	32	50	50	46	50
52	52	52	50	57	50	52	50	71	64	50	75
75	76	52	52	78	52	92	52	96	88	51	77
125	103	100	100	125	91	153	103	101	102	90	128
79	74	87	74	95	77	97	78	89	88	85	96
53	46	50	48	61	46	69	46	54	48	40	43
88	85	74	47	94	83	91	67	100	87	81	82
34	45	19	16	51	24	56	33	28	31	27	37
125	88	0	0	88	52	73	106	73	83	46	67
0	17	0	0	30	5	18	38	21	30	0	37
375	300	300	300	400	300	736	310	300	300	155	225
375	388	300	300	450	300	736	464	303	414	271	383
79	75	86	71	94	75	94	76	88	84	78	86
48	46	48	46	60	43	65	44	49	41	36	44
10	23	23	14	29	18	28	18	38	23	38	18
39	39	48	39	48	46	50	46	57	50	48	66
75	75	75	75	78	75	75	75	75	75	75	78
92	78	113	78	136	78	138	92	113	113	93	113
150	150	155	150	155	150	155	150	150	150	150	150
155	155	155	150	192	150	184	155	155	150	150	150
225	233	300	300	310	233	310	276	288	162	155	225
300	300	300	300	400	300	736	310	300	225	155	225
1.0	1.2	3.1	0	0	0	5.3	4.0	0	2.9	0	2.5
.5	.6	2.4	0	0	0	.9	.5	0	1.0	0	1.4

TABLE 10.21.--COOKED WHOLE KERNEL CORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	23.9	3.2	25.8	30.1	34.6
ON ONLY 1 OF 3 DAYS.....PCT.....	20.4	2.2	22.2	25.4	29.2
ON ONLY 2 OF 3 DAYS.....PCT.....	3.2	1.0	3.3	4.5	4.7
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.3	.3	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.4	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.6	.5	.4	.5
MAXIMUM.....DO.....	6.0	3.0	5.0	4.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	38	10	22	27	32
STANDARD DEVIATION.....GM.....	27	6	16	19	22
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7		4	7	7
25TH.....GM.....	26		13	14	18
50TH.....GM.....	28		19	26	28
75TH.....GM.....	51		28	28	41
90TH.....GM.....	57		42	55	57
95TH.....GM.....	85		55	57	62
99TH.....GM.....	133		85	110	112
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	97	28	58	72	81
STANDARD DEVIATION.....GM.....	59	21	41	48	44
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	96	16	56	62	75
STANDARD DEVIATION.....GM.....	52	7	28	37	39
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	110	0	49	68	100
STANDARD DEVIATION.....GM.....	53	0	15	41	41
MAXIMUM IN ANY 1 DAY.....GM.....	850	85	255	510	340
MAXIMUM OVER 3 DAYS.....GM.....	869	85	267	510	680
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	95	22	56	68	78
STANDARD DEVIATION.....GM.....	56	17	40	45	41
PERCENTILES: <u>1/</u>					
5TH.....GM.....	21		10	11	20
25TH.....GM.....	65		31	41	43
50TH.....GM.....	83		43	77	83
75TH.....GM.....	123		83	85	85
90TH.....GM.....	170		85	123	154
95TH.....GM.....	170		154	165	170
99TH.....GM.....	330		170	170	170
MAXIMUM.....GM.....	850	85	255	510	340
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	49.3	0	2.7	0	5.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.9	0	.7

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
32.0	31.0	28.8	24.5	26.7	24.3	23.5	18.9	17.4	15.0	16.6	13.1
26.3	26.1	23.5	21.5	23.2	21.4	19.9	16.1	14.5	12.5	14.6	11.2
5.0	4.5	4.9	2.8	3.3	2.6	3.2	2.6	2.3	2.3	1.5	1.7
.7	.5	.4	.2	.2	.3	.4	.3	.6	.1	.5	.1
1.2	1.2	1.2	1.2	1.2	1.1	1.2	1.2	1.3	1.2	1.3	1.2
.5	.5	.5	.4	.4	.4	.5	.5	.5	.5	.8	.4
5.0	4.0	4.0	4.0	4.0	3.0	5.0	4.0	3.0	3.0	6.0	3.0
39	33	48	36	46	35	44	37	41	38	43	35
30	22	33	28	30	22	30	26	27	25	31	19
11	7	14	10	14	10	14	7	14	7	7	14
26	19	28	26	28	26	28	26	26	26	26	26
28	28	41	28	41	28	28	28	28	28	28	28
55	43	57	51	57	51	57	51	55	51	55	51
77	57	84	57	83	57	82	57	83	57	83	57
99	77	113	95	110	77	104	84	89	85	128	69
161	113	170	113	154	113	154	128	154	117	145	103
98	86	122	95	120	94	110	92	105	96	115	93
66	50	73	61	66	50	58	54	60	58	76	43
96	84	111	96	122	90	118	97	95	95	79	95
56	43	58	54	59	38	56	50	37	46	48	30
113	77	150	142	128	121	122	109	111	108	139	39
47	44	80	81	56	29	52	55	37	0	8	0
850	340	510	454	495	340	492	369	369	350	425	206
850	510	863	869	825	510	680	738	462	462	434	308
95	83	116	94	118	93	110	92	99	93	98	89
62	47	70	59	66	48	58	52	54	51	54	39
21	21	41	39	41	22	41	21	28	21	21	39
77	43	83	63	83	77	77	63	77	77	77	77
83	83	85	83	85	83	85	83	85	83	83	83
85	85	165	85	165	123	154	115	145	123	128	85
170	165	170	170	170	165	170	165	165	165	170	165
170	170	246	170	246	170	231	170	170	170	231	170
340	248	340	340	340	231	308	252	330	340	255	170
850	340	510	454	495	330	492	369	369	350	255	170
5.0	5.2	0	0	4.1	12.0	9.8	1.9	1.2	2.0	0	0
.6	.6	0	0	.3	.8	.7	.1	.5	.6	0	0

TABLE 10.22.--CREAM-STYLE CORN
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.3	8.5	5.6	4.7	5.1
ON ONLY 1 OF 3 DAYS.....PCT.....	3.9	6.2	5.0	4.0	4.6
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	1.0	.6	.7	.5
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	1.3	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.6	1.1	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.9	.3	.4	.3
MAXIMUM.....DO.....	5.0	5.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	53	39	36	40	46
STANDARD DEVIATION.....GM.....	38	24	29	34	28
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11		2	5	11
25TH.....GM.....	28		16	21	21
50TH.....GM.....	43		21	32	43
75TH.....GM.....	85		43	43	43
90TH.....GM.....	85		85	85	85
95TH.....GM.....	128		85	128	85
99TH.....GM.....	171		128	171	149
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	144	99	86	98	128
STANDARD DEVIATION.....GM.....	88	50	62	68	72
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	151	58	137	131	115
STANDARD DEVIATION.....GM.....	91	13	34	74	60
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	117	70	0	0	0
STANDARD DEVIATION.....GM.....	88	36	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	768	213	256	256	384
MAXIMUM OVER 3 DAYS.....GM.....	1,024	383	384	512	448
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	141	75	95	104	123
STANDARD DEVIATION.....GM.....	86	46	65	70	71
PERCENTILES: <u>1/</u>					
5TH.....GM.....	31		8	16	16
25TH.....GM.....	71		57	64	64
50TH.....GM.....	128		64	84	128
75TH.....GM.....	192		128	128	128
90TH.....GM.....	256		213	256	256
95TH.....GM.....	256		256	256	256
99TH.....GM.....	512		256	256	256
MAXIMUM.....GM.....	768	213	256	256	384
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	11.9	.6	0	0	1.5
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	.9	0	0	1.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
4.6	4.5	4.2	3.9	4.4	4.0	4.6	3.9	4.7	4.7	1.9	2.7
4.3	4.2	4.0	3.8	4.2	3.7	4.1	3.5	3.9	4.1	1.4	2.3
.2	.3	.1	.1	.2	.4	.5	.3	.8	.6	.5	.4
0	0	0	0	0	0	(2)	(2)	0	.1	0	0
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.2	1.6	1.2
.3	.3	.2	.2	.2	.3	.4	.4	.5	.5	1.1	.6
2.0	2.0	2.0	2.0	2.0	2.0	3.0	4.0	3.0	4.0	4.0	3.0
50	54	60	46	63	47	63	51	62	56	123	56
38	33	42	33	41	31	41	35	32	32	105	44
11	10	11	4	13	7	21	10	21	21		
28	43	43	21	43	27	43	28	43	43		
43	43	43	43	43	43	43	43	43	43		
64	85	85	43	85	43	85	85	85	85		
85	85	128	85	85	85	128	85	128	85		
85	107	171	85	128	85	171	107	128	96		
171	171	213	161	256	213	213	171	149	213		
136	154	179	128	180	129	166	139	166	150	210	132
76	87	126	79	108	68	95	80	74	71	103	90
210	148	117	241	188	129	182	134	142	126	424	190
143	74	72	0	99	91	72	76	65	48	122	64
0	0	0	0	0	0	167	175	0	213	0	0
0	0	0	0	0	0	96	117	0	0	0	0
512	512	640	384	768	512	512	512	448	289	512	512
1,024	512	640	483	768	640	768	896	448	640	1,024	512
140	148	172	131	179	129	165	137	150	136	234	138
86	80	111	75	109	80	90	77	71	62	74	81
32	16	32	32	38	21	64	32	48	49		
64	84	128	64	128	64	128	64	128	128		
128	128	128	128	128	128	128	128	128	128		
128	192	256	128	256	128	256	192	192	128		
256	256	256	256	256	256	256	256	256	256		
256	256	384	256	287	256	384	256	256	256		
512	256	512	256	513	512	512	279	256	256		
512	512	512	256	768	512	512	384	256	256	384	512
0	1.2	0	0	0	1.3	1.8	4.6	0	.9	0	0
0	1.1	0	0	0	.5	.7	1.5	0	.9	0	0

TABLE 10.23.--COOKED LIMA BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.8	1.0	2.4	1.9	1.9
ON ONLY 1 OF 3 DAYS.....PCT.....	2.5	1.0	2.1	1.5	1.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	.3	.4	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.0	1.2	1.3	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.4	.4	.3
MAXIMUM.....DO.....	3.0	1.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	43	24	21	21	29
STANDARD DEVIATION.....GM.....	33	22	21	15	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7				
25TH.....GM.....	28				
50TH.....GM.....	29				
75TH.....GM.....	58				
90TH.....GM.....	85				
95TH.....GM.....	113				
99TH.....GM.....	170				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	114	71	52	56	84
STANDARD DEVIATION.....GM.....	82	67	30	34	45
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	120	0	77	40	64
STANDARD DEVIATION.....GM.....	77	0	64	33	66
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	101	0	0	0	0
STANDARD DEVIATION.....GM.....	65	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	875	175	175	175	175
MAXIMUM OVER 3 DAYS.....GM.....	875	175	350	175	350
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	110	71	54	49	79
STANDARD DEVIATION.....GM.....	75	67	38	31	47
PERCENTILES: <u>1/</u>					
5TH.....GM.....	21				
25TH.....GM.....	67				
50TH.....GM.....	88				
75TH.....GM.....	170				
90TH.....GM.....	175				
95TH.....GM.....	219				
99TH.....GM.....	350				
MAXIMUM.....GM.....	875	175	175	88	175
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	11.4	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.9	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.8	2.3	2.6	1.8	2.7	2.5	3.5	3.2	4.5	4.3	2.9	3.7
1.6	2.2	2.4	1.8	2.3	2.3	3.1	2.8	4.3	3.7	2.7	2.8
.2	.1	.2	.1	.4	.2	.4	.4	.3	.6	.2	.8
0	0	0	0	0	0	0	(2)	0	0	0	.1
1.1	1.1	1.2	1.0	1.2	1.1	1.1	1.2	1.1	1.2	1.3	1.3
.3	.3	.4	.2	.4	.3	.4	.4	.3	.4	.5	.5
2.0	3.0	3.0	2.0	3.0	2.0	3.0	3.0	2.0	2.0	2.0	3.0
43	30	54	31	54	39	49	41	52	40	49	52
50	15	45	26	35	24	37	30	41	31	37	38
	7			7	7	15	7	15	7		
	15			29	28	29	28	29	22		
	29			57	29	29	29	44	29		
	29			58	58	58	58	58	58		
	58			113	58	113	58	88	68		
	58			117	85	117	88	131	88		
	58			170	117	248	170	233	175		
112	91	146	96	150	113	125	109	138	98	130	123
148	46	101	78	105	68	74	73	82	62	95	81
132	59	205	11	119	89	169	105	214	129	175	142
34	26	125	0	32	47	92	61	132	64	0	61
0	0	0	0	0	0	0	146	0	0	0	7
0	0	0	0	0	0	0	0	0	0	0	0
875	175	700	340	510	350	394	510	394	350	263	350
875	175	744	340	510	350	744	525	700	525	350	510
114	86	141	91	137	106	129	103	138	101	111	123
133	45	94	78	83	61	79	59	87	61	60	83
	22			43	21	44	22	44	21		
	44			88	58	88	67	88	77		
	88			131	88	88	88	88	88		
	88			175	170	175	170	175	117		
	175			175	175	196	175	350	175		
	175			340	175	340	175	350	175		
	175			454	350	350	263	350	350		
875	175	350	340	454	350	350	340	350	350	175	350
.7	0	.9	1.0	1.5	1.4	3.2	2.7	0	0	0	0
1.7	0	2.2	3.5	1.2	.9	1.6	1.0	0	0	0	0

TABLE 10.24.--COOKED MIXED VEGETABLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.4	11.4	3.7	3.1	4.0
ON ONLY 1 OF 3 DAYS.....PCT.....	3.0	7.7	3.3	2.7	3.8
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	3.0	.3	.3	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.1	.7	.1	0	(2)
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.5	1.3	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.9	.7	.3	.3
MAXIMUM.....DO.....	4.0	4.0	4.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	46	42	38	25	29
STANDARD DEVIATION.....GM.....	33	26	47	17	16
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11	9		7	7
25TH.....GM.....	30	21		16	16
50TH.....GM.....	31	38		20	31
75TH.....GM.....	62	45		31	31
90TH.....GM.....	78	81		39	61
95TH.....GM.....	123	104		62	62
99TH.....GM.....	187	115		94	78
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	118	99	86	66	83
STANDARD DEVIATION.....GM.....	70	57	83	38	44
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	126	79	97	79	101
STANDARD DEVIATION.....GM.....	63	37	26	38	12
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	138	96	256	0	43
STANDARD DEVIATION.....GM.....	77	18	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	454	344	454	187	187
MAXIMUM OVER 3 DAYS.....GM.....	935	344	767	281	233
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	117	81	89	69	82
STANDARD DEVIATION.....GM.....	69	47	78	40	44
PERCENTILES: <u>1/</u>					
5TH.....GM.....	28	23		22	22
25TH.....GM.....	91	57		47	47
50TH.....GM.....	94	64		60	94
75TH.....GM.....	182	115		94	94
90TH.....GM.....	187	128		94	140
95TH.....GM.....	187	135		182	187
99TH.....GM.....	374	344		187	187
MAXIMUM.....GM.....	454	344	454	187	187
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	11.3	0	0	0	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	0	2.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.7	3.4	2.7	1.8	2.7	2.9	3.1	3.3	4.7	4.5	4.6	6.4
3.5	3.3	2.7	1.7	2.5	2.6	2.8	2.7	4.2	3.7	2.9	5.3
.2	.1	0	.1	.2	.3	.2	.5	.4	.6	1.6	.7
0	0	0	0	.1	0	.1	(2)	0	.2	.1	.3
1.0	1.0	1.0	1.1	1.1	1.1	1.2	1.2	1.2	1.3	1.4	1.3
.2	.2	.0	.2	.4	.3	.5	.5	.4	.6	.5	.5
2.0	2.0	1.0	2.0	3.0	2.0	4.0	3.0	2.0	3.0	3.0	3.0
41	35	36	44	48	44	57	49	57	55	62	52
32	17	20	26	34	26	46	32	31	38	40	28
15	8			15	12	16	16	16	8		30
30	30			31	31	31	31	31	31		31
31	31			31	31	47	31	62	38		39
47	47			62	61	62	62	62	62		62
62	62			78	62	125	94	123	94		94
94	62			121	94	125	125	123	130		125
249	62			125	151	312	184	125	187		125
112	102	107	134	126	114	146	121	152	128	124	148
63	51	60	80	69	58	86	68	69	69	47	86
158	81	0	53	159	133	128	135	186	149	145	91
144	10	0	31	94	44	70	56	1	33	67	28
0	0	0	0	125	0	192	78	0	156	109	77
0	0	0	0	0	0	91	0	0	58	0	18
374	187	374	374	374	454	374	374	369	374	281	374
748	187	374	374	748	468	935	561	374	561	561	374
116	101	107	124	129	116	146	122	149	128	133	123
75	50	60	80	73	58	85	66	54	56	60	75
46	23			46	23	47	47	47	47		47
91	91			94	94	94	91	94	94		91
94	94			94	94	116	94	182	121		94
137	94			187	182	187	187	187	187		182
187	187			187	187	234	187	187	187		187
281	187			281	187	374	187	187	187		374
374	187			374	281	374	374	281	187		374
374	187	374	374	374	454	374	374	281	281	281	374
3.4	2.6	2.6	.9	0	0	0	0	0	0	0	0
4.2	3.5	6.9	3.2	0	0	0	0	0	0	0	0

TABLE 10.25.--COOKED COWPEAS, FIELD PEAS, AND BLACK EYE PEAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.9	.5	2.1	2.5	2.7
ON ONLY 1 OF 3 DAYS.....PCT.....	2.5	.5	1.8	2.2	2.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	.2	.3	(2)
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER.....	1.2	1.0	1.3	1.2	1.0
STANDARD DEVIATION.....DO.....	.5	.0	.9	.4	.2
MAXIMUM.....DO.....	5.0	1.0	5.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	53	42	28	32	34
STANDARD DEVIATION.....GM.....	42	21	25	22	21
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	11				7
25TH.....GM.....	29				28
50TH.....GM.....	44				29
75TH.....GM.....	58				44
90TH.....GM.....	117				58
95TH.....GM.....	117				58
99TH.....GM.....	233				117
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	140	127	73	93	98
STANDARD DEVIATION.....GM.....	101	64	70	68	56
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	136	0	56	63	175
STANDARD DEVIATION.....GM.....	90	0	38	17	0
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	193	0	69	0	0
STANDARD DEVIATION.....GM.....	89	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,050	175	350	263	350
MAXIMUM OVER 3 DAYS.....GM.....	1,050	175	350	263	350
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	131	127	63	84	97
STANDARD DEVIATION.....GM.....	88	64	50	60	57
PERCENTILES: <u>1</u> /					
5TH.....GM.....	22				22
25TH.....GM.....	88				85
50TH.....GM.....	88				88
75TH.....GM.....	175				131
90TH.....GM.....	196				175
95TH.....GM.....	350				175
99TH.....GM.....	350				350
MAXIMUM.....GM.....	700	175	175	263	350
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER.....	10.7	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
2.7	2.3	3.2	2.4	2.7	2.5	3.7	3.0	3.7	3.8	3.4	2.2
2.5	2.3	3.0	2.1	2.3	2.3	3.2	2.7	2.7	3.3	2.6	1.3
.1	(2)	.2	.3	.4	.2	.5	.3	1.0	.5	.9	.9
.1	0	0	0	.1	0	0	(2)	0	0	0	0
1.1	1.1	1.1	1.1	1.2	1.1	1.2	1.2	1.5	1.3	1.5	1.5
.4	.2	.3	.3	.5	.4	.4	.5	.6	.5	.6	.5
3.0	2.0	2.0	2.0	4.0	3.0	2.0	3.0	3.0	3.0	3.0	2.0
41	34	57	62	66	48	62	55	57	51	56	72
31	25	28	45	55	32	51	46	31	37	34	34
7	4			14	7	15	14		11		
29	19			29	28	29	29		29		
29	29			58	31	57	43		29		
58	39			88	58	59	58		58		
58	58			117	88	117	117		117		
88	88			117	117	143	175		117		
233	117			292	146	292	233		175		
109	97	161	162	153	135	165	143	183	140	144	188
57	69	76	101	98	94	127	104	95	108	93	109
174	175	159	181	179	113	157	145	68	112	122	130
115	0	21	103	89	36	111	107	38	55	42	33
87	0	0	0	280	0	0	233	0	0	0	0
0	0	0	0	14	0	0	0	0	0	0	0
350	350	350	350	525	438	1,050	875	438	525	350	438
700	350	350	700	875	438	1,050	875	438	525	350	438
109	96	151	163	161	126	157	136	115	113	110	143
60	67	63	100	93	83	106	94	75	68	46	83
22	11			44	22	43	22		22		
88	58			88	85	88	88		88		
88	88			175	88	131	88		88		
175	88			175	175	175	175		175		
175	175			350	263	340	263		175		
175	175			350	340	350	350		263		
350	350			525	350	526	525		350		
350	350	350	350	525	350	700	525	350	350	175	438
0	0	.6	0	1.4	3.1	4.3	1.4	0	0	0	0
0	0	1.1	0	1.1	2.0	2.0	.5	0	0	0	0

TABLE 10.26.--COOKED GREEN PEAS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	18.3	16.0	21.8	20.9	22.1
ON ONLY 1 OF 3 DAYS.....PCT.....	15.7	10.5	18.5	17.6	19.5
ON ONLY 2 OF 3 DAYS.....PCT.....	2.4	5.0	3.1	3.1	2.6
ON ALL 3 OF 3 DAYS.....PCT.....	.2	.5	.2	.1	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.5	1.2	1.2	1.1
STANDARD DEVIATION.....DO.....	.4	.7	.4	.4	.4
MAXIMUM.....DO.....	6.0	4.0	3.0	3.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	35	30	21	24	27
STANDARD DEVIATION.....GM.....	26	23	17	20	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	7	5	3	3	4
25TH.....GM.....	19	14	7	14	14
50TH.....GM.....	28	28	14	19	28
75TH.....GM.....	55	43	28	28	28
90TH.....GM.....	57	57	43	55	57
95TH.....GM.....	85	67	57	57	57
99TH.....GM.....	113	142	85	110	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	92	80	51	58	71
STANDARD DEVIATION.....GM.....	60	68	35	37	48
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	94	56	60	73	81
STANDARD DEVIATION.....GM.....	55	35	39	45	41
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	96	32	50	75	65
STANDARD DEVIATION.....GM.....	52	20	4	11	22
MAXIMUM IN ANY 1 DAY.....GM.....	850	425	255	340	340
MAXIMUM OVER 3 DAYS.....GM.....	1,190	425	340	425	425
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	90	61	53	61	72
STANDARD DEVIATION.....GM.....	57	45	36	42	46
PERCENTILES: <u>1/</u>					
5TH.....GM.....	20	7	10	10	11
25TH.....GM.....	43	29	21	41	43
50TH.....GM.....	85	57	43	43	83
75TH.....GM.....	85	80	85	85	85
90TH.....GM.....	170	128	85	85	165
95TH.....GM.....	170	135	128	165	170
99TH.....GM.....	330	213	170	170	170
MAXIMUM.....GM.....	680	255	170	330	340
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	41.2	0	2.1	1.6	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.5	0	.8	.4	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY, 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
20.9	19.4	18.1	16.9	17.6	15.7	19.9	17.6	18.9	17.5	15.0	15.9
18.0	16.7	15.2	14.8	15.5	14.1	17.1	15.2	15.6	14.4	11.1	12.9
2.7	2.6	2.8	2.1	2.1	1.6	2.5	2.3	2.8	2.7	3.4	2.9
.2	.1	.1	0	(2)	.1	.3	.1	.5	.3	.5	.1
1.2	1.2	1.2	1.1	1.1	1.1	1.2	1.2	1.3	1.2	1.4	1.2
.4	.4	.4	.4	.4	.3	.4	.4	.6	.5	.9	.5
3.0	4.0	3.0	3.0	3.0	3.0	3.0	4.0	5.0	3.0	6.0	3.0
34	32	44	36	41	34	42	35	41	39	45	34
23	23	39	28	27	23	30	24	29	27	35	22
7	7	7	7	14	7	11	7	11	14	14	7
19	14	28	18	28	21	28	19	28	28	28	19
28	28	28	28	28	28	28	28	28	28	28	28
43	43	57	55	57	55	57	43	57	55	57	43
57	57	83	57	57	57	71	57	83	57	110	57
78	83	110	83	85	57	110	85	98	85	113	57
113	113	170	113	113	113	125	113	170	140	158	128
87	82	109	93	109	92	109	92	99	98	96	93
54	52	66	58	66	54	66	59	53	68	69	56
90	94	129	111	108	97	107	86	111	95	117	63
48	43	106	68	53	52	62	42	51	34	41	38
77	88	100	0	112	33	109	114	111	101	127	127
8	20	8	0	28	0	63	63	43	31	42	0
495	425	850	510	680	413	680	340	340	680	340	340
495	425	1,190	850	680	510	1,020	850	598	680	475	383
86	83	112	96	107	91	106	88	97	94	96	82
52	46	73	62	64	55	61	52	50	59	60	48
20	20	32	28	32	20	31	21	21	31	21	21
43	43	83	54	83	56	83	54	83	83	56	43
85	83	85	83	85	85	85	83	85	85	85	85
85	85	165	128	165	85	165	85	124	85	114	85
170	170	170	170	170	170	170	170	170	170	170	165
170	170	255	170	170	170	170	170	170	170	170	170
255	170	340	340	340	330	340	255	170	255	330	170
495	255	510	510	680	413	495	340	340	680	330	340
4.4	2.6	0	.7	5.8	1.5	10.8	10.1	0	1.6	0	0
.9	.5	0	.3	.7	.2	.9	.7	0	.4	0	0

TABLE 10.27.--COOKED SPINACH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.5	.9	2.8	3.2	5.1
ON ONLY 1 OF 3 DAYS.....PCT.....	4.0	.9	2.6	3.2	5.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	0	.1	(2)	.2
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.1	1.1	1.1	1.0	1.0
STANDARD DEVIATION.....DO.....	.3	.3	.4	.2	.2
MAXIMUM.....DO.....	3.0	2.0	3.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	45	10	21	25	33
STANDARD DEVIATION.....GM.....	29	6	18	18	21
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	8			1	8
25TH.....GM.....	30			15	17
50TH.....GM.....	34			30	32
75TH.....GM.....	65			34	34
90TH.....GM.....	68			39	65
95TH.....GM.....	103			63	68
99TH.....GM.....	137			98	95
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	122	30	58	76	92
STANDARD DEVIATION.....GM.....	73	18	51	55	56
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	122	0	81	51	119
STANDARD DEVIATION.....GM.....	59	0	0	0	28
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	119	0	43	0	0
STANDARD DEVIATION.....GM.....	33	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	454	51	205	293	308
MAXIMUM OVER 3 DAYS.....GM.....	570	51	205	293	308
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	121	26	58	73	93
STANDARD DEVIATION.....GM.....	70	19	48	53	56
PERCENTILES: <u>1/</u>					
5TH.....GM.....	24			4	20
25TH.....GM.....	78			45	51
50TH.....GM.....	103			68	95
75TH.....GM.....	185			98	103
90TH.....GM.....	205			103	190
95TH.....GM.....	205			190	195
99TH.....GM.....	380			293	285
MAXIMUM.....GM.....	454	51	205	293	308
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	14.5	0	0	0	1.7
PROPORTION OF EATING OCCASIONS.....PCT.....	.8	0	0	0	1.7

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
5.2	3.6	4.5	3.0	4.0	4.0	4.7	4.9	6.3	6.3	5.5	5.5
4.9	3.4	4.0	2.8	3.9	3.7	4.1	4.5	5.5	4.8	3.8	4.2
.3	.1	.5	.2	.1	.4	.5	.5	.8	1.5	1.6	1.3
(2)	0	0	0	0	0	.1	(2)	0	0	0	0
1.1	1.1	1.1	1.1	1.0	1.1	1.2	1.1	1.2	1.3	1.3	1.2
.3	.2	.3	.2	.2	.3	.4	.3	.4	.5	.5	.4
3.0	2.0	2.0	2.0	2.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0
38	36	47	38	52	45	51	47	44	55	57	48
24	22	28	26	27	28	33	30	20	33	42	30
8	8	16		16	10	16	15	14	16		4
24	17	32		33	32	32	32	32	32		33
33	33	34		49	34	34	34	34	48		43
51	49	65		65	63	65	65	65	68		60
65	65	86		69	68	86	73	68	101		90
87	68	98		103	85	133	127	68	127		120
114	110	137		130	171	167	137	103	137		151
103	104	137	105	152	121	129	130	132	138	121	132
61	62	81	64	80	67	64	78	62	77	86	88
130	94	91	129	121	134	148	126	64	125	143	93
24	62	46	56	37	64	71	51	19	57	63	46
86	0	0	0	0	0	126	152	0	0	0	0
0	0	0	0	0	0	9	0	0	0	0	0
410	308	410	308	410	385	390	425	308	390	309	454
410	330	410	412	410	512	513	458	308	570	475	454
105	102	127	108	150	122	131	129	112	128	128	117
59	62	80	64	79	64	63	75	62	64	74	77
23	19	23		48	29	37	38	41	45		23
71	52	90		98	95	95	95	52	90		74
98	98	103		146	103	103	103	95	103		103
130	140	190		195	180	195	190	172	190		180
195	195	205		205	195	205	205	205	205		190
195	205	205		293	205	205	256	205	205		270
205	308	410		390	380	308	385	308	308		454
410	308	410	308	410	385	390	425	308	380	309	454
1.4	1.7	.8	.9	3.1	2.6	0	2.3	0	0	0	0
1.1	2.1	1.2	1.9	1.9	1.1	0	.6	0	0	0	0

TABLE 10.28.--COOKED STRING BEANS
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	27.3	19.7	25.1	25.4	31.6
ON ONLY 1 OF 3 DAYS.....PCT.....	22.7	13.0	20.5	21.5	26.7
ON ONLY 2 OF 3 DAYS.....PCT.....	4.1	5.7	4.0	3.8	4.3
ON ALL 3 OF 3 DAYS.....PCT.....	.5	1.1	.5	.2	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.6	1.2	1.2	1.2
STANDARD DEVIATION.....DO.....	.5	.8	.5	.4	.5
MAXIMUM.....DO.....	6.0	5.0	4.0	3.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	35	36	20	20	26
STANDARD DEVIATION.....GM.....	29	27	16	18	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	6	3	3	3	5
25TH.....GM.....	23	17	11	11	12
50TH.....GM.....	23	29	12	18	23
75TH.....GM.....	47	45	23	23	28
90TH.....GM.....	69	76	45	45	47
95TH.....GM.....	93	77	47	47	70
99TH.....GM.....	140	134	85	70	93
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	88	82	48	53	63
STANDARD DEVIATION.....GM.....	57	68	37	52	40
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	90	77	49	51	71
STANDARD DEVIATION.....GM.....	51	38	26	26	34
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	110	61	63	44	81
STANDARD DEVIATION.....GM.....	82	34	32	15	33
MAXIMUM IN ANY 1 DAY.....GM.....	840	344	220	700	280
MAXIMUM OVER 3 DAYS.....GM.....	1,820	401	304	700	420
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	86	69	48	51	64
STANDARD DEVIATION.....GM.....	54	47	33	46	38
PERCENTILES: <u>1/</u>					
5TH.....GM.....	18	9	8	8	16
25TH.....GM.....	67	35	34	34	35
50TH.....GM.....	70	57	35	35	68
75TH.....GM.....	135	107	68	70	70
90TH.....GM.....	140	128	70	70	135
95TH.....GM.....	140	135	101	135	140
99TH.....GM.....	280	213	140	140	210
MAXIMUM.....GM.....	840	344	220	700	280
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	91.4	2.0	0	4.2	8.2
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	1.3	0	.8	1.2

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
31.1	29.4	29.5	24.8	25.4	23.4	29.4	27.6	27.2	29.8	32.1	27.2
25.2	24.9	25.9	21.5	22.6	19.9	24.9	22.8	20.5	22.3	25.8	18.6
5.5	4.3	3.4	3.0	2.5	3.2	4.0	4.3	5.4	6.7	4.4	7.0
.4	.2	.2	.2	.3	.3	.5	.5	1.3	.8	1.8	1.6
1.2	1.2	1.2	1.2	1.1	1.2	1.2	1.2	1.4	1.4	1.3	1.5
.5	.5	.4	.4	.4	.4	.5	.5	.7	.7	.7	.8
4.0	5.0	3.0	4.0	4.0	3.0	6.0	6.0	4.0	5.0	5.0	6.0
31	30	36	33	38	33	42	37	45	42	38	42
31	25	29	22	24	22	37	27	36	33	30	29
5	6	9	11	11	6	12	11	11	11	11	9
18	15	23	23	23	23	23	23	23	23	23	23
23	23	23	23	34	23	35	23	41	35	23	35
45	45	47	45	47	47	47	47	47	47	47	47
47	47	68	47	70	49	90	70	93	93	77	90
70	70	90	79	93	76	93	93	117	93	93	93
117	135	140	117	123	113	175	140	228	187	161	140
75	76	91	89	100	85	103	92	110	94	99	94
47	57	51	58	56	48	59	54	86	57	73	55
76	79	114	79	101	87	111	99	90	97	66	91
42	52	79	36	40	44	54	52	43	57	29	48
138	98	99	73	114	95	186	91	124	140	101	80
160	19	24	13	41	36	142	51	51	50	43	39
840	700	630	420	420	338	700	560	700	420	420	280
1,820	700	770	420	560	560	1,820	700	700	595	514	560
75	74	93	83	99	85	106	91	99	92	87	86
54	55	58	51	53	47	60	52	60	47	55	47
17	17	27	26	34	18	34	26	28	17	17	18
35	35	68	68	68	68	70	68	70	68	68	68
70	70	70	70	70	70	87	70	70	70	70	70
70	70	140	105	140	135	140	135	140	140	135	135
140	140	140	140	140	140	140	140	140	140	140	140
140	140	140	140	147	140	210	140	140	140	140	140
270	280	280	280	280	270	280	280	280	210	280	280
840	700	560	420	420	280	560	420	454	280	397	280
4.2	4.8	5.1	5.2	13.2	5.9	12.0	18.5	0	6.0	0	2.1
.5	.6	1.1	1.2	1.2	.4	.7	.8	0	.8	0	.5

TABLE 10.29.--COOKED SUMMER SQUASH
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	2.8	.7	1.3	1.4	1.1
ON ONLY 1 OF 3 DAYS.....PCT.....	2.4	.7	1.0	1.4	1.0
ON ONLY 2 OF 3 DAYS.....PCT.....	.3	0	.2	.1	.1
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	0	0	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.2	1.0	1.3	1.0	1.1
STANDARD DEVIATION.....DO.....	.4	.0	.4	.2	.3
MAXIMUM.....DO.....	4.0	1.0	2.0	2.0	2.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	56	9	41	34	50
STANDARD DEVIATION.....GM.....	46	6	25	47	63
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	11				
25TH.....GM.....	36				
50TH.....GM.....	36				
75TH.....GM.....	72				
90TH.....GM.....	108				
95TH.....GM.....	143				
99TH.....GM.....	287				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	143	26	98	73	100
STANDARD DEVIATION.....GM.....	95	19	59	49	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	169	0	115	352	291
STANDARD DEVIATION.....GM.....	98	0	3	0	162
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	192	0	0	0	0
STANDARD DEVIATION.....GM.....	108	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	860	54	215	352	529
MAXIMUM OVER 3 DAYS.....GM.....	1,075	54	237	704	881
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	145	26	96	97	136
STANDARD DEVIATION.....GM.....	98	19	63	91	121
PERCENTILES: <u>1/</u>					
5TH.....GM.....	27				
25TH.....GM.....	105				
50TH.....GM.....	108				
75TH.....GM.....	215				
90TH.....GM.....	215				
95TH.....GM.....	352				
99TH.....GM.....	430				
MAXIMUM.....GM.....	860	54	215	352	529
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	12.3	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	1.0	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.2	1.7	2.1	1.2	2.3	2.6	3.6	3.4	5.5	5.8	4.4	4.3
1.0	1.6	1.9	1.2	2.1	2.4	3.2	3.0	4.3	5.2	3.7	3.3
.1	.1	.2	.1	.2	.2	.3	.3	1.3	.5	.6	.7
.1	0	0	0	(2)	0	.1	.1	0	.1	0	.3
1.3	1.1	1.1	1.1	1.2	1.1	1.1	1.2	1.2	1.1	1.1	1.4
.6	.2	.3	.2	.5	.3	.4	.5	.4	.4	.4	.7
3.0	2.0	2.0	2.0	3.0	3.0	4.0	4.0	2.0	3.0	2.0	4.0
43	36	55	43	60	64	61	52	61	60	45	69
30	19	30	29	46	53	51	42	31	47	22	65
				9	17	13	9	18	18		
				36	36	36	36	36	36		
				47	54	44	36	59	36		
				72	72	72	72	72	72		
				143	108	108	108	90	117		
				167	176	143	143	142	143		
				179	287	358	287	143	189		
100	107	155	119	152	175	156	130	161	151	124	137
60	56	79	82	98	144	92	79	64	90	66	67
139	60	154	134	215	180	162	193	127	194	97	131
26	40	53	0	88	107	101	105	65	86	9	55
107	0	0	0	167	0	263	89	0	324	0	288
0	0	0	0	0	0	122	33	0	0	0	0
215	215	430	430	529	860	430	430	352	704	215	352
323	215	430	430	744	860	1,075	860	430	973	215	865
103	102	155	121	154	173	158	132	146	160	117	143
50	56	76	78	92	138	97	85	69	106	59	87
				26	52	40	26	53	54		
				108	108	108	95	108	108		
				134	108	108	108	161	108		
				215	215	215	161	215	215		
				215	352	323	215	215	352		
				287	430	430	323	215	352		
				454	860	430	430	352	430		
215	215	430	430	529	860	430	430	352	704	215	352
0	0	0	0	2.7	0	3.1	5.1	0	1.3	0	0
0	0	0	0	2.5	0	1.5	1.8	0	1.1	0	0

TABLE 10.30.--COOKED SWEET POTATOES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1</u> / (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	4.1	10.8	3.8	3.1	3.2
ON ONLY 1 OF 3 DAYS.....PCT.....	3.7	8.0	3.7	3.1	2.9
ON ONLY 2 OF 3 DAYS.....PCT.....	.4	1.7	.1	0	.3
ON ALL 3 OF 3 DAYS.....PCT.....	(2)	1.1	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1</u> /NUMBER....	1.2	1.5	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.4	.7	.4	.2	.4
MAXIMUM.....DO.....	4.0	3.0	3.0	2.0	3.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1</u> /GM.....	52	40	36	34	38
STANDARD DEVIATION.....GM.....	38	24	35	19	25
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1</u> /					
5TH.....GM.....	14	9		7	8
25TH.....GM.....	34	21		20	19
50TH.....GM.....	38	38		34	34
75TH.....GM.....	68	45		38	40
90TH.....GM.....	85	76		68	75
95TH.....GM.....	132	81		70	100
99TH.....GM.....	200	147		113	113
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	141	103	94	102	103
STANDARD DEVIATION.....GM.....	92	48	77	57	61
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	145	86	78	0	102
STANDARD DEVIATION.....GM.....	77	60	0	0	57
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1</u> /GM.....	123	55	193	0	0
STANDARD DEVIATION.....GM.....	81	19	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,020	229	450	338	300
MAXIMUM OVER 3 DAYS.....GM.....	1,020	442	579	338	339
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1</u> /GM.....	136	82	97	96	99
STANDARD DEVIATION.....GM.....	87	47	70	50	62
PERCENTILES: <u>1</u> /					
5TH.....GM.....	38	19		18	26
25TH.....GM.....	86	57		60	56
50TH.....GM.....	114	64		103	86
75TH.....GM.....	185	115		113	115
90TH.....GM.....	225	128		150	225
95TH.....GM.....	238	135		205	225
99TH.....GM.....	450	229		225	225
MAXIMUM.....GM.....	1,020	229	338	225	300
QUANTITIES NOT SPECIFIED BY USERS: <u>1</u> /					
EATING OCCASIONS.....NUMBER....	13.4	0	0	0	1.0
PROPORTION OF EATING OCCASIONS.....PCT.....	.7	0	0	0	1.5

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
3.4	2.1	3.2	3.3	2.9	3.3	3.6	4.9	7.8	7.2	8.5	8.1
3.1	1.9	2.9	3.0	2.6	3.2	3.4	4.4	6.4	6.0	7.5	6.7
.3	.2	.4	.3	.3	.1	.2	.4	1.4	1.2	.9	1.3
0	0	0	0	0	(2)	(2)	(2)	0	.1	0	.2
1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	1.3	1.2	1.3
.3	.3	.4	.3	.4	.3	.4	.4	.4	.5	.6	.6
2.0	2.0	3.0	2.0	2.0	4.0	4.0	3.0	2.0	4.0	3.0	3.0
52	50	57	61	70	51	61	49	62	48	59	47
31	34	32	33	50	42	38	33	48	39	38	28
17				22	9	20	17	19	11		17
35				38	34	38	33	34	22		34
38				62	38	50	38	40	38		38
70				75	68	75	67	75	60		56
77				133	78	103	76	123	98		77
119				150	137	150	113	153	113		114
150				340	178	205	200	265	300		160
142	139	153	170	188	140	177	132	146	118	161	119
82	98	72	88	137	92	112	76	89	73	108	51
157	113	153	160	198	194	147	142	188	135	158	113
37	45	69	46	71	65	46	71	97	93	24	59
0	0	0	0	0	333	103	146	0	98	0	114
0	0	0	0	0	0	0	47	0	0	0	0
510	510	420	476	1,020	700	800	630	615	450	456	304
510	510	453	476	1,020	1,000	800	630	795	900	456	481
144	134	150	166	183	144	168	130	157	114	145	113
79	92	75	84	129	92	100	71	104	57	89	49
38				56	28	57	42	57	33		46
105				113	103	114	99	103	76		100
114				168	114	130	114	119	113		114
205				225	200	210	180	205	130		119
225				255	228	238	225	228	200		191
225				450	400	410	225	410	210		225
450				1,020	450	450	315	615	357		255
510	510	420	476	1,020	510	800	630	615	357	456	281
2.1	0	.8	0	0	0	2.0	1.2	1.3	2.3	0	2.8
2.7	0	1.5	0	0	0	1.0	.3	1.2	1.4	0	2.7

TABLE 10.31.--TOMATO JUICES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	3.9	0	.8	.9	.9
ON ONLY 1 OF 3 DAYS.....PCT.....	2.6	0	.7	.8	.7
ON ONLY 2 OF 3 DAYS.....PCT.....	.8	0	0	(2)	.2
ON ALL 3 OF 3 DAYS.....PCT.....	.4	0	.1	0	0
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.5	0	1.6	1.2	1.3
STANDARD DEVIATION.....DO.....	.8	0	1.1	.5	.9
MAXIMUM.....DO.....	7.0	0	4.0	3.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	94	0	77	60	59
STANDARD DEVIATION.....GM.....	72	0	71	47	43
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	30				
25TH.....GM.....	45				
50TH.....GM.....	81				
75TH.....GM.....	121				
90TH.....GM.....	182				
95TH.....GM.....	243				
99TH.....GM.....	354				
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	191	0	211	184	135
STANDARD DEVIATION.....GM.....	105	0	222	143	41
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	208	0	0	61	177
STANDARD DEVIATION.....GM.....	104	0	0	0	98
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	204	0	121	0	0
STANDARD DEVIATION.....GM.....	99	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	1,879	0	790	729	456
MAXIMUM OVER 3 DAYS.....GM.....	2,844	0	790	729	699
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	190	0	147	156	133
STANDARD DEVIATION.....GM.....	85	0	73	61	48
PERCENTILES: <u>1/</u>					
5TH.....GM.....	91				
25TH.....GM.....	122				
50TH.....GM.....	182				
75TH.....GM.....	243				
90TH.....GM.....	243				
95TH.....GM.....	363				
99TH.....GM.....	486				
MAXIMUM.....GM.....	1,150	0	243	243	243
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	13.6	0	0	0	0
PROPORTION OF EATING OCCASIONS.....PCT.....	.6	0	0	0	0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
1.2	1.0	2.1	2.2	3.7	3.5	6.3	5.7	7.4	5.0	4.7	7.2
1.1	.9	1.4	1.6	2.7	2.5	3.8	3.8	5.2	3.8	2.8	5.1
.1	.1	.5	.2	.5	.8	1.8	1.3	.9	.8	1.8	1.3
0	0	.3	.4	.5	.3	.7	.6	1.3	.4	.2	.9
1.1	1.2	1.5	1.5	1.5	1.4	1.6	1.5	1.6	1.3	1.5	1.4
.2	.4	.7	.8	.8	.7	.9	.8	.9	.7	.7	.7
2.0	2.0	3.0	3.0	4.0	3.0	7.0	6.0	4.0	4.0	4.0	3.0
57	73	93	99	108	99	107	96	80	69	99	68
20	44	67	67	84	64	86	73	48	37	106	33
				40	40	35	31	10	20		40
				61	61	61	61	41	41		41
				81	81	81	81	61	61		61
				122	141	151	121	121	81		81
				243	182	182	182	162	121		122
				324	243	243	243	172	182		122
				405	303	384	354	182	182		161
169	187	181	217	204	214	206	187	170	165	117	155
58	103	79	111	87	114	156	68	97	60	34	58
93	199	209	183	273	228	205	210	193	141	219	146
42	64	135	43	131	102	74	127	40	53	56	43
0	0	188	195	238	195	225	219	136	154	567	121
0	0	9	76	88	52	140	73	35	30	0	(2)
304	547	486	605	547	729	1,879	1,573	486	304	972	363
304	547	972	1,089	1,214	1,094	2,844	2,541	547	547	1,701	484
159	183	191	194	222	211	198	189	153	154	199	144
63	95	94	84	96	93	99	67	68	55	104	49
				121	121	92	92	30	91		121
				182	151	122	122	121	122		122
				182	182	182	182	122	122		122
				243	243	243	243	182	182		182
				365	363	243	243	243	243		182
				484	365	365	273	243	243		242
				514	607	486	484	365	304		363
304	547	486	484	545	608	1,150	486	365	304	486	363
0	0	0	1.6	0	0	4.4	1.3	2.2	0	0	4.0
0	0	0	3.2	0	0	.9	.2	1.7	0	0	3.9

TABLE 10.32.--CATSUP
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	13.7	.4	13.4	21.8	21.3
ON ONLY 1 OF 3 DAYS.....PCT.....	11.0	.2	11.2	16.8	16.5
ON ONLY 2 OF 3 DAYS.....PCT.....	2.4	.2	2.2	4.6	4.6
ON ALL 3 OF 3 DAYS.....PCT.....	.3	0	.1	.4	.3
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.3	1.5	1.2	1.3	1.3
STANDARD DEVIATION.....DO.....	.6	.5	.5	.6	.6
MAXIMUM.....DO.....	5.0	2.0	3.0	4.0	5.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	9	5	5	8	7
STANDARD DEVIATION.....GM.....	11	(2)	7	10	7
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2		1	1	2
25TH.....GM.....	2		2	2	2
50TH.....GM.....	5		4	5	5
75TH.....GM.....	10		5	10	10
90TH.....GM.....	20		10	20	15
95TH.....GM.....	25		15	25	23
99TH.....GM.....	51		34	46	46
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	15	12	18	17
STANDARD DEVIATION.....GM.....	28	0	10	24	17
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	21	9	14	20	18
STANDARD DEVIATION.....GM.....	22	0	11	18	15
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	22	0	73	17	14
STANDARD DEVIATION.....GM.....	21	0	0	5	7
MAXIMUM IN ANY 1 DAY.....GM.....	495	15	83	273	137
MAXIMUM OVER 3 DAYS.....GM.....	495	17	220	273	137
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	20	11	13	17	16
STANDARD DEVIATION.....GM.....	25	4	13	22	16
PERCENTILES: <u>1/</u>					
5TH.....GM.....	3		3	3	5
25TH.....GM.....	6		6	6	6
50TH.....GM.....	15		11	15	15
75TH.....GM.....	28		15	15	15
90TH.....GM.....	38		30	30	30
95TH.....GM.....	68		33	68	45
99TH.....GM.....	137		68	68	68
MAXIMUM.....GM.....	495	15	83	273	137
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	706.3	0	16.5	57.1	51.7
PROPORTION OF EATING OCCASIONS.....PCT.....	10.6	0	9.5	11.6	10.0

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
22.5	20.4	21.1	18.5	19.4	14.5	11.2	7.3	5.5	5.5	4.3	3.5
17.3	16.1	16.0	13.3	15.8	12.1	8.8	6.4	4.1	4.8	3.8	3.0
4.1	3.5	4.5	4.4	3.1	2.2	2.2	.8	1.3	.7	.3	.3
1.0	.7	.6	.8	.5	.3	.2	.1	.1	0	.2	.2
1.3	1.3	1.4	1.4	1.3	1.2	1.3	1.2	1.3	1.1	1.2	1.2
.6	.6	.7	.7	.6	.5	.5	.4	.7	.4	.5	.5
4.0	5.0	5.0	5.0	5.0	4.0	4.0	3.0	5.0	2.0	3.0	3.0
10	8	12	9	11	7	9	7	7	6	6	5
15	11	15	10	12	8	11	13	7	6	4	5
2	2	2	2	2	1	2	1	1	1		
4	3	5	4	4	2	4	2	3	2		
5	5	9	5	5	5	5	5	5	5		
10	10	15	10	14	10	10	7	10	5		
20	15	25	23	23	15	18	11	10	10		
27	23	35	28	35	23	23	20	23	15		
91	51	91	47	60	46	51	46	46	40		
23	18	27	22	25	18	23	20	18	16	17	14
39	21	33	22	26	18	30	41	23	13	14	14
26	19	34	22	30	17	20	11	11	16	9	3
26	15	31	23	31	14	18	6	5	18	0	0
24	30	18	15	26	32	17	12	27	0	6	15
25	32	8	9	14	26	14	7	0	0	0	0
454	303	360	273	273	137	341	495	137	68	60	68
454	333	360	288	375	273	341	495	137	120	60	68
23	18	26	20	25	18	21	17	16	15	14	13
34	24	29	23	25	18	26	36	18	15	9	13
4	5	6	3	6	4	6	3	3	3		
6	6	11	6	11	6	6	6	6	6		
15	15	15	15	15	15	15	15	15	15		
30	15	30	28	30	17	30	15	15	15		
45	30	57	45	60	33	33	30	30	30		
68	45	68	68	68	60	60	45	33	60		
137	137	137	120	137	102	137	137	137	68		
454	273	273	273	205	137	341	495	137	68	33	68
62.6	72.2	52.2	62.8	89.0	83.5	78.5	63.9	1.9	11.2	0	3.3
9.9	12.5	12.7	16.5	9.0	8.9	11.2	10.6	2.3	10.2	0	8.0

TABLE 10.33.--MUSTARD
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER.....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	14.6	0	6.1	10.6	14.4
ON ONLY 1 OF 3 DAYS.....PCT.....	10.6	0	5.2	8.0	10.2
ON ONLY 2 OF 3 DAYS.....PCT.....	3.1	0	.3	2.2	3.5
ON ALL 3 OF 3 DAYS.....PCT.....	.9	0	.6	.4	.7
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER.....	1.4	0	1.3	1.4	1.4
STANDARD DEVIATION.....DO.....	.7	0	.6	.6	.7
MAXIMUM.....DO.....	6.0	0	3.0	3.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	4	0	2	3	3
STANDARD DEVIATION.....GM.....	4	0	2	3	3
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	1		(2)	(2)	(2)
25TH.....GM.....	2		1	2	2
50TH.....GM.....	2		2	2	2
75TH.....GM.....	5		2	3	3
90TH.....GM.....	7		3	7	5
95TH.....GM.....	10		5	8	7
99TH.....GM.....	20		10	10	12
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	8	0	5	7	6
STANDARD DEVIATION.....GM.....	7	0	4	7	5
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	8	0	3	6	7
STANDARD DEVIATION.....GM.....	8	0	1	4	6
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	8	0	5	6	6
STANDARD DEVIATION.....GM.....	7	0	2	3	2
MAXIMUM IN ANY 1 DAY.....GM.....	120	0	30	60	65
MAXIMUM OVER 3 DAYS.....GM.....	165	0	30	60	70
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	8	0	5	6	6
STANDARD DEVIATION.....GM.....	7	0	3	6	5
PERCENTILES: <u>1/</u>					
5TH.....GM.....	3		1	1	1
25TH.....GM.....	5		3	3	5
50TH.....GM.....	5		5	5	5
75TH.....GM.....	10		5	5	5
90TH.....GM.....	15		5	15	10
95TH.....GM.....	20		15	15	15
99TH.....GM.....	30		15	30	28
MAXIMUM.....GM.....	120	0	15	60	60
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER.....	811.0	0	6.8	24.3	39.0
PROPORTION OF EATING OCCASIONS.....PCT.....	10.5	0	8.4	9.8	10.2

1/ SEE TABLE NOTES IN APPENDIX.

2/ MORE THAN 0 BUT LESS THAN 0.5 IF ROUNDED TO WHOLE NUMBER; LESS THAN 0.05 IF ROUNDED TO 1 DECIMAL PLACE.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
17.9	16.6	20.4	16.6	21.1	16.6	18.2	11.5	10.7	7.5	5.5	3.9
12.9	11.7	14.9	11.9	14.1	12.8	12.1	9.5	8.4	5.8	3.4	2.2
3.6	4.0	4.1	3.7	4.9	3.1	4.7	1.7	2.0	1.6	2.2	1.5
1.5	.9	1.5	1.0	2.0	.7	1.5	.3	.4	.1	0	.2
1.5	1.4	1.4	1.5	1.6	1.3	1.5	1.2	1.3	1.3	1.4	1.6
.8	.7	.8	.7	.9	.6	.8	.5	.5	.5	.5	.7
5.0	5.0	4.0	5.0	6.0	5.0	6.0	3.0	3.0	3.0	2.0	3.0
4	3	5	3	5	3	4	3	3	2	3	3
4	3	7	3	6	3	4	2	3	2	3	2
1	1	1	1	1	1	1	1	1	1		
2	2	2	2	2	2	2	2	2	2		
2	2	3	2	3	2	3	2	2	2		
5	3	7	3	5	3	5	3	3	3		
8	6	10	7	10	5	10	5	5	5		
12	10	17	8	15	9	15	7	10	5		
19	15	40	13	28	15	20	10	18	10		
8	7	11	8	9	7	9	7	7	5	4	9
8	6	11	6	8	6	9	6	6	3	1	9
7	7	13	7	11	8	9	7	6	6	9	3
5	6	11	4	11	7	8	4	4	3	3	1
9	6	13	6	11	8	8	5	12	5	0	1
5	3	13	5	9	7	4	2	6	0	0	0
65	60	120	45	90	120	75	57	45	15	15	29
65	90	150	75	165	125	90	57	55	30	30	29
7	7	11	7	9	7	8	6	7	6	7	5
7	6	11	5	9	7	8	5	7	3	4	6
1	3	3	3	3	3	3	1	1	3		
5	5	5	5	5	5	5	5	5	5		
5	5	5	5	5	5	5	5	5	5		
10	5	15	5	15	10	10	5	5	5		
15	15	29	15	15	15	15	15	15	10		
15	15	30	15	30	15	30	15	15	15		
30	30	60	30	45	30	30	30	30	15		
60	60	60	45	60	120	60	57	45	15	15	29
50.8	52.7	43.3	40.1	119.0	107.4	175.5	112.1	13.6	23.1	3.4	0
9.3	10.4	10.6	11.2	9.3	9.2	12.9	11.1	9.0	14.0	8.2	0

TABLE 10.34.--CUCUMBER PICKLES
CONSUMED IN 3 DAYS BY INDIVIDUALS, 48 STATES, 1977-78

MEASURE AND UNIT	TOTAL	AGE (YEARS) AND SEX OF INDIVIDUAL			
		UNDER 1	1-2	3-5	6-8
		MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE	MALE AND FEMALE
TOTAL INDIVIDUALS <u>1/</u> (WEIGHTED).....NUMBER....	37,874	498	1,045	1,719	1,841
TOTAL INDIVIDUALS USING FOOD:					
AT LEAST ONCE IN 3 DAYS.....PCT.....	9.2	.2	4.6	6.2	8.1
ON ONLY 1 OF 3 DAYS.....PCT.....	7.9	.2	4.4	5.6	7.2
ON ONLY 2 OF 3 DAYS.....PCT.....	1.0	0	.3	.5	.8
ON ALL 3 OF 3 DAYS.....PCT.....	.2	0	0	0	.1
EATING OCCASIONS AT WHICH ITEM WAS CONSUMED IN 3 DAYS BY USERS:					
AVERAGE <u>1/</u>NUMBER....	1.2	1.0	1.1	1.1	1.1
STANDARD DEVIATION.....DO.....	.5	.0	.4	.3	.4
MAXIMUM.....DO.....	5.0	1.0	3.0	2.0	4.0
QUANTITY CONSUMED BY USERS:					
USED AT LEAST ONCE IN 3 DAYS:					
AVERAGE PER DAY <u>1/</u>GM.....	18	2	12	14	17
STANDARD DEVIATION.....GM.....	20	0	9	14	20
AVERAGE PER DAY AT SPECIFIED PERCENTILES: <u>1/</u>					
5TH.....GM.....	2			2	2
25TH.....GM.....	6			5	5
50TH.....GM.....	11			10	10
75TH.....GM.....	22			22	22
90TH.....GM.....	43			29	43
95TH.....GM.....	50			45	64
99TH.....GM.....	90			72	108
USED ON ONLY 1 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	45	6	33	41	42
STANDARD DEVIATION.....GM.....	47	0	28	42	47
USED ON ONLY 2 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	48	0	30	30	60
STANDARD DEVIATION.....GM.....	44	0	14	16	49
USED ON ALL 3 OF 3 DAYS:					
AVERAGE <u>1/</u>GM.....	47	0	0	0	55
STANDARD DEVIATION.....GM.....	34	0	0	0	0
MAXIMUM IN ANY 1 DAY.....GM.....	810	7	130	215	270
MAXIMUM OVER 3 DAYS.....GM.....	945	7	130	215	370
QUANTITY CONSUMED PER EATING OCCASION:					
AVERAGE <u>1/</u>GM.....	45	6	32	38	45
STANDARD DEVIATION.....GM.....	45	0	26	36	46
PERCENTILES: <u>1/</u>					
5TH.....GM.....	7			7	7
25TH.....GM.....	16			13	15
50TH.....GM.....	30			26	30
75TH.....GM.....	65			60	65
90TH.....GM.....	90			74	130
95TH.....GM.....	130			135	135
99TH.....GM.....	222			155	222
MAXIMUM.....GM.....	455	7	130	195	260
QUANTITIES NOT SPECIFIED BY USERS: <u>1/</u>					
EATING OCCASIONS.....NUMBER....	217.1	0	.8	8.3	11.0
PROPORTION OF EATING OCCASIONS.....PCT.....	5.3	0	1.5	7.0	6.5

1/ SEE TABLE NOTES IN APPENDIX.

SOURCE: USDA NATIONWIDE FOOD CONSUMPTION SURVEY 1977-78, 48 CONTERMINOUS STATES.

AGE (YEARS) AND SEX
OF INDIVIDUAL--CONTINUED

9-14		15-18		19-34		35-64		65-74		75 AND OVER	
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE
2,089	2,158	1,394	1,473	3,928	5,346	4,929	7,069	1,118	1,738	536	993
8.6	9.1	9.9	8.5	11.7	11.0	10.7	9.1	7.8	6.6	7.6	6.3
7.7	8.1	8.6	7.3	10.0	9.8	8.7	7.8	6.4	5.9	6.2	5.2
.6	.8	.9	1.0	1.4	1.1	1.5	1.1	1.4	.6	.9	.9
.2	.2	.5	.1	.3	.1	.5	.2	0	.1	.5	.2
1.1	1.2	1.2	1.2	1.2	1.2	1.3	1.2	1.2	1.1	1.3	1.2
.4	.5	.6	.5	.5	.4	.6	.5	.4	.4	.7	.5
3.0	4.0	4.0	4.0	4.0	4.0	5.0	4.0	2.0	3.0	3.0	3.0
18	19	18	23	19	15	21	17	15	15	18	17
19	28	19	28	18	16	21	19	13	16	24	14
3	2	2	2	3	2	2	2	2	2		5
7	5	7	7	7	6	8	6	4	5		7
10	10	11	12	12	10	13	11	10	10		12
22	22	22	25	22	22	24	22	22	19		22
43	43	43	49	43	35	45	39	26	28		43
65	65	65	87	51	45	60	45	43	50		49
87	108	90	127	87	87	90	90	65	90		49
49	46	45	61	47	43	52	42	36	36	47	40
51	51	49	76	43	46	50	42	29	34	62	32
48	77	65	54	54	37	49	45	42	60	16	55
38	99	39	51	38	31	40	40	24	48	6	19
29	48	32	74	48	27	52	62	0	14	63	19
34	28	17	51	17	17	34	48	0	0	32	8
378	810	270	455	270	430	540	405	135	195	325	135
378	945	270	455	405	430	540	518	195	270	325	148
47	50	45	58	48	40	50	42	37	38	42	42
50	59	46	71	40	41	45	43	29	34	53	36
7	7	7	7	10	7	7	7	7	6		13
15	13	20	14	20	15	20	16	13	15		16
30	30	30	30	34	30	34	30	30	26		30
65	65	60	65	65	60	65	65	60	57		60
130	130	85	135	111	74	130	74	70	85		130
150	195	130	195	130	130	135	130	78	130		135
195	325	270	325	195	200	195	195	135	135		135
378	405	270	455	270	390	270	405	135	135	325	135
15.9	10.2	10.4	6.8	25.5	52.1	39.8	27.4	2.6	6.4	0	0
7.9	4.4	6.1	4.6	4.6	7.6	6.0	3.6	2.5	4.9	0	0

APPENDIX A

System for Classifying Foods Into Food Groups Based on the First Three Digits of Seven-digit Food Codes

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
1 MILK AND MILK PRODUCTS	11 Milk and Milk Drinks	111 Milk, fluid--pasteurized, filled, buttermilk, and dry reconstituted 112 Milk, concentrated fluid (canned) 113 Milk, imitation 114 Yogurt 115 Chocolate, malted, shakes, and other flavored milk drinks 116 Meal replacements with milk 117 Baby formulas 118 Milk, dry and powdered mixtures with dry milk, not reconstituted
	12 Cream and Cream Substitutes	121 Sweet dairy cream (fluid whipped and dry) 122 Sweet cream and whipped cream substitutes 123 Sour dairy cream
	13 Milk Desserts	131 Milk desserts, frozen 132 Milk desserts, not frozen 133 Baby foods with milk 134 Milk sauces and gravies 135 Other milk products
	14 Cheeses	141 Natural cheese 142 Cottage cheese 143 Cream cheese 144 Processed cheese and cheese spreads 145 Imitation cheese 146 Cheese mixtures 147 Cheese soups
2 MEAT, POULTRY, FISH, AND MIXTURES	21 Beef	210 Meat, NFS ¹ , and beef, NFS ¹ 211 Beef steaks with bone 212 Beef steaks without bone 213 Beef cuts with bone, not steaks 214 Beef slices and chunks 215 Ground beef patties and meat balls 216 Other beef items 217 Baby and junior beef
	22 Pork	220 Pork, NFS ¹ 221 Pork chops 222 Pork steaks and cutlets 223 Ham 224 Pork roasts, not ham 225 Canadian bacon 226 Bacon, salt pork, and fat back

¹NFS - not further specified

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
		227 Miscellaneous pork cuts
		228 Baby and junior pork, ham, and meat sticks
	23 Lamb, Veal, Game, and Other Carcass Meat	230 Lamb, NFS ¹
		231 Lamb
		232 Veal
		233 Game
		234 Baby veal and lamb
	24 Poultry	241 Chicken
		242 Turkey
		243 Duck
		244 Rock cornish game hen and other poultry
		247 Baby and junior poultry
	25 Organ Meats, Sausages, Luncheon Meats, and Spreads	251 Organ meats and mixtures
		252 Frankfurters, sausages, luncheon meats, and meat spreads
	26 Fish and Shellfish	261 Finfish
		262 Other seafood
		263 Shellfish
	27 Meat, Poultry, or Fish Combined With Nonmeat Items	271 Meat, poultry, or fish in gravy, sauce, or creamed
		272 Meat, poultry, or fish combined with starch
		273 Meat, poultry, or fish with starch and vegetable
		274 Meat, poultry, or fish with vegetable, excluding white potatoes
		275 Sandwiches (reported as a unit) with meat, poultry, or fish
		276 Baby foods, combination with meat, poultry, or fish
	28 Plate Meals, Soups and Gravies with Meat, Poultry or Fish Base and Plain Gelatin Drinks	281 Frozen plate meals and soups
		283 Soups, broths, and extracts from meat, poultry, or fish base
		284 Gelatin drink, plain
		285 Gravies, meat or poultry base, made with water

¹NFS - not further specified

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
3 EGGS, MIXTURES, AND SUBSTITUTES	31 Eggs	311 Chicken eggs 312 Other poultry eggs
	32 Egg Mixtures	321 Egg dishes 322 Egg sandwiches 323 Egg soups 324 Meringues
	33 Egg Substitutes	330 Substitutes, NFS ¹ 331 Made from powdered mixtures 332 Made from frozen mixtures 333 Made from liquid mixtures
	34 Baby-food Eggs	341 Egg yolk
	41 Legumes	411 Cooked or canned dried beans 412 Cooked or canned dried bean mixtures 413 Cooked dried peas and lentils and mixtures 414 Soybean dried products 415 Frozen meals with dried beans or peas as main course 416 Soups, mainly legumes 417 Baby-food legumes 418 Meat substitutes 419 Meat substitute sandwiches
		421 Nuts
		422 Nut butters
		423 Nut butter sandwiches
		424 Coconut beverages and mixtures
		425 Nut mixtures
		431 Seeds and seed mixtures
		441 Carob powders
		442 Carob chips
		500 Flour and dry mixes
		510 Bread, rolls, NFS ¹
	51 Yeast Breads and Rolls	511 White bread and rolls 512 Whole wheat bread and rolls 513 Cracked wheat bread and rolls 514 Rye bread and rolls 515 Oatmeal bread
4 DRY LEGUMES, NUTS, AND SEEDS	42 Nuts and Nut Butters and Mixtures	
	43 Seeds and Seed Mixtures	
	44 Carob and Carob Products	
	50 Flour	
5 GRAIN PRODUCTS		

¹NFS - Not further specified

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
		516 Multigrain bread
		517 Cottonseed bread
		518 Other breads
	52 Quick Breads	521 Biscuits
		522 Cornbread and corn muffins
		523 Other muffins and popovers
		524 Quickbreads excluding cornbread and muffins
	53 Cakes, Cookies, Pies, and Pastries	531 Cakes
		532 Cookies
		533 Pies
		534 Cobblers, eclairs, turnovers, and other pastries
		535 Danish, breakfast pastries, bars, and doughnuts
		536 Coffee cakes, not yeast type
	54 Crackers and Salty Snacks from Grain Products	541 Sweet crackers
		542 Low sodium (dietary) crackers
		543 Nonsweet crackers
		544 Salty snack products from grain products
	55 Pancakes, Waffels, and French Toast	551 Pancakes
		552 Waffles
		553 French toast
		554 Crepes
		555 Flour-water patties
		556 Flour-milk patties
		557 Rice flour cakes
	56 Cooked Pastes and Cooked Cereals	561 Pastes
		562 Cooked cereals
	57 Ready-to-eat Cereals	571 Cereal, ready-to-eat, NFS ¹
		571 Buckwheat cereals
		572 Bran cereals
		573 Corn cereals
		574 Oat cereals
		575 Rice cereals
		576 Wheat cereals
		577 Multigrain cereals
		578 Baby cereals
		579 Other cereals

¹NFS - not further specified

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
	58 Mixtures, Mostly Grain	581 Mixtures with animal protein 582 Mixtures without animal protein 583 Frozen plate meals 584 Soups with grain product as main ingredient 585 Baby food mixtures with animal protein
6 FRUITS	61 Citrus Fruits, and Juices	611 Citrus fruits 612 Citrus fruit juices
	62 Dried Fruits	621 Dried fruits
	63 Fruits and Berries (exclude citrus, dried)	631 Fruit excluding berries 632 Berries 633 Mixtures of two or more fruits 634 Mixtures of fruits, berries, and nonfruit items
	64 Fruit Juices and Nectars (exclude citrus)	641 Juices 642 Nectars
	67 Baby or Strained and Junior Fruits and Juices	671 Fruits and fruit mixtures 672 Juices 673 Fruits with cereal 674 Fruit desserts and puddings
7 VEGETABLES	71 White Potatoes	710 White potatoes, NFS ¹ 711 Baked, boiled, and canned 712 Chips and sticks 713 Creamed, scalloped, and au gratin 714 Fried 715 Mashed, stuffed, and puffs 716 Salad 717 Special recipes 718 Soups 719 Puerto Rican starchy vegetables
	72 Dark-green Vegetables	721 Dark-green leafy vegetables 722 Dark-green nonleafy vegetables 723 Dark-green vegetable soups
	73 Deep-yellow Vegetables	731 Carrots 732 Pumpkin 733 Squash 734 Sweetpotatoes 735 Deep-yellow vegetable soups

¹NFS - not further specified

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
	74 Tomatoes	741 Raw tomatoes 742 Cooked tomatoes 743 Tomato juice 744 Tomato sauces 745 Tomato mixtures 746 Tomato soups 747 Tomato sandwiches
	75 Other Vegetables and Vegetable Mixtures	751 Raw vegetables 752 Cooked or canned vegetables 753 Cooked vegetables, mixture of two or more vegetables (including nuts) 754 Cooked vegetables with sauces, batters, casseroles 755 Olives, pickles, and relishes (excluding tomatoes) 756 Vegetable soups
	76 Baby or Strained and Junior Vegetables and Mixtures, Mostly Vegetables	761 Dark-green vegetables 762 Deep-yellow vegetables 764 Vegetables other than dark green, deep-yellow, and tomato 765 Potato mixtures with meat 766 Vegetable mixtures and meat with cereal 767 Vegetable mixtures and liver with cereal
	77 Vegetable Mixtures with Animal Protein	771 White potato mixtures 772 Puerto Rican starchy vegetable (viandas) mixtures 773 Other vegetable mixtures 775 Puerto Rican stews and soups with starchy vegetables (viandas)
8 FATS, OILS, AND SALAD DRESSINGS	81 Fats	811 Table fats 812 Cooking fats 813 Other fats
	82 Oils	821 Vegetable oils
	83 Salad Dressings	831 Regular-type salad dressings 832 Low-calorie-type salad dressing
9 SUGAR, SWEETS, AND BEVERAGES	91 Sugar and Sweets	911 Sugars 912 Sugar replacements and substitutes 913 Sirups, honey, and molasses 914 Jellies, jams, and preserves 915 Gelatin desserts and salads 916 Ices and popsicles

Food Group Codes--Continued

<u>Major food groups</u>	<u>Major food subgroups</u>	<u>Minor food subgroups</u>
		917 Candies
		918 Chewing gum and cough drops
	92 Nonalcoholic Beverages	921 Coffee
		922 Coffee substitutes
		923 Tea
		924 Soft drinks
		925 Fruit ades and drinks
		926 Nonfruit beverages
		929 Sugar concentrates with vitamin C, powdered, not reconstituted
	93 Alcoholic Beverages	931 Beers and ales
		932 Cordials and liqueurs
		933 Cocktails
		934 Wines
		935 Distilled liquors
0 FOOD ITEMS FOR WHICH NO NUTRITIVE VALUES ARE ASSIGNED	00 Blank	001 Artificial sweeteners
		002 Extracts, flavors, and vinegar
		003 Seasonings, spices, and herbs
		004 Vitamins, minerals, and supplements
		009 Cooking oils, sprays, and sticks

APPENDIX B

Table Notes

Data in tables were derived as follows:

1. Total individuals include only those individuals with 3-day diet records.
2. Average number of eating occasions at which food was consumed in 3 days by users =

$$\frac{\text{Total number of eating occasions at which food was consumed}}{\text{Number of individuals using food at least once in 3 days}}$$

3. Average quantity per day consumed by individuals who used food at least once in 3 days =

$$\frac{\text{Total grams consumed by users divided by 3}}{\text{Number of individuals using at least once in 3 days}}$$

4. Percentiles are omitted when fewer than 50 individuals in a sex-age group reported the food.

5. Average quantity per day consumed by individuals who used food on only 1 of 3 days =

$$\frac{\text{Total grams consumed by 1-day users}}{\text{Number of individuals using on only 1 of 3 days}}$$

6. Average quantity per day consumed by individuals who used food on only 2 of 3 days =

$$\frac{\text{Total grams consumed by 2-day users divided by 2}}{\text{Number of individuals using on only 2 of 3 days}}$$

7. Average quantity per day consumed by individuals who used food on all 3 of 3 days =

$$\frac{\text{Total grams consumed by 3-day users divided by 3}}{\text{Number of individuals using on all 3 of 3 days}}$$

8. Average quantity consumed per eating occasion =

$$\frac{\text{Total grams consumed}}{\text{Total number of eating occasions}}$$

9. See number 4 above.

10. Quantities not specified by users are included in "quantity consumed by users" but excluded in "quantity consumed per eating occasion."

United States
Department of Agriculture

Washington, D C
20250

OFFICIAL BUSINESS
Penalty for Private Use \$300

Postage and Fees Paid
U S Department of Agriculture
AGR-101



FOURTH CLASS MAIL